



Diversity of the Epigeal Entomofauna in Southern Santa Fe Province, Argentina

Natacha María Noelia Schanzempch¹, Pablo Guillermo Rimoldi^{1*}

¹Department of Biology and Ecology (FCV-UNR) and Center for Environmental Studies (CEAV), Faculty of Veterinary Sciences, National University of Rosario (FCV-UNR), Casilda, Santa Fe, Argentina.

ABSTRACT

Biodiversity protection is crucial due to the impact of human activities on ecosystems. Anthropogenic activities, such as advancing agricultural frontier, deforestation, and the introduction of exotic species, are the main drivers of changes in biodiversity. Entomofauna, especially insects, play a vital role in ecosystem indicators of environmental quality and contribute to key functions such as pollination and decomposition. The present research focuses on the understory entomofauna in the Faculty of Veterinary Sciences of the National University of Rosario, Argentina. Several samplings were carried out in 2022 in five different microsites regarding herbaceous, shrub, and tree vegetation, using pitfall traps to capture arthropods. A total of 2631 individuals of 68 morphospecies belonging to five classes, 15 orders, and 43 families were captured. The class Insecta was the most diverse and abundant, accounting for 82.35% of the morphospecies and 35.80% of the total abundance. The class Malacostraca, although with only two species, was dominant in abundance (58.57%). Microsite diversity showed that microsites 5 and 1 had the highest biodiversity. The results showed low similarity between microsites, indicating the influence of the landscape on entomofaunal composition. This study provides an essential baseline on understory entomofauna in the Faculty of Veterinary Sciences. The diversity found highlights the importance of this habitat and the identification of key species such as *Armadillidium vulgare* and *Entomobryidae* sp. to their role in ecosystem services. This research lays the groundwork for future comparative studies and highlights the need to consider biodiversity in conservation and environmental management politics agendas.

Keywords: Soil fauna, Understorey, Anthropization, Argentina.

HOW TO CITE THIS ARTICLE: Schanzempch NMN, Rimoldi PG. Diversity of the Epigeal Entomofauna in Southern Santa Fe Province, Argentina. *Entomol Appl Sci Lett.* 2023;10(4):1-12. <https://doi.org/10.51847/vmeMGxgwlg>

Corresponding author: Pablo Guillermo Rimoldi

E-mail ✉ primoldi04@gmail.com

Received: 16/09/2023

Accepted: 17/11/2023

INTRODUCTION

Biodiversity protection is a relevant issue in global conservation. Given the environmental impact of human activities on natural systems, biodiversity knowledge is a worldwide challenge [1].

Anthropogenic activities such as overexploitation and habitat loss, deforestation, pollution, and the introduction of exotic species, among others, are the main forces driving changes in the diversity and composition of communities worldwide [2-4].

Entomofauna is the dominant group of animals on earth, with the number of known species exceeding three times that of all other animal species combined. These living beings conquered the planet more than 350 million years ago, and their importance in the world is such that without arthropods, life would be very different from how we know it, which is why they form an important component of ecosystem diversity [5]. Generally, the degree of disturbance of a local ecosystem is interpreted by the presence or absence of various arthropods, their variations in abundance, diversity, and composition of biological groups. Insects, for example, are

excellent indicators frequently used to determine and monitor certain ecosystems' environmental quality [6]. The latter is explained by being one of the most diverse and widely distributed groups, rapid identification, simple sampling, convenient size, and predictable response to environmental variations, and being a useful, reliable, and economical tool [7]. In addition, entomofauna play important roles in pollination, seed dispersal, aeration, and soil rotation and act as parasites, decomposers of organic material, and nutrient recyclers. In addition, they form the basis of many food chains and ecological interactions, promote soil fertility and structure, and provide biological control mechanisms [8]. All of the above attributes cause arthropod communities to vary according to the degree of disturbance (natural and anthropogenic), abundance, diversity, and composition of biological groups, as they are strongly correlated with ecosystem functioning and reflect habitat heterogeneity, development, and recovery. Likewise, structural arthropod communities reflect the degree of fragmentation and isolation of ecosystems in the landscape [9].

In recent decades, in Argentina, the simplification of the landscape due to the expansion and intensification of agricultural activity has been very marked, causing accelerated fragmentation of the landscape [10] and consequent losses in biodiversity.

There are contributions to the arthropod communities of agroecosystems in different provinces of our country; for example, a study carried out in Entre Ríos estimates the biodiversity of arthropods to implement conservation strategies [11] in the same province studies of diversity and abundance were carried out in agroecosystems [10, 12] and a particular soybean crop [11]. In the province of Santa Fe, diversity was also analyzed in different crops, such as soybeans [13].

This research presents the taxonomic richness of the understory entomofauna present in the FCV-UNR property, understood as a complex (vertical variation) and heterogeneous (horizontal variation) environment in terms of plant diversity, to be used as a baseline for future comparative work with environments that have experienced intense transformations of the original vegetation cover, becoming more

homogeneous environments with a simplified plant structure (agroecosystems).

MATERIALS AND METHODS

Area of study

The Faculty of Veterinary Sciences of the National University of Rosario is located in Casilda, head of the Caseros Department, in the south of the province of Santa Fe. Its property covers an area of approximately 240 hectares and was declared a "Natural Protected Area" in 2007 (CD Resolution N° 188/07) due to its leading role as a refuge for wildlife in a purely agricultural area (**Figure 1**).

The climate in this region is temperate, with average temperatures between 14 and 20 °C [14]. Precipitation varies throughout the year, being more intense in spring and summer. Before the region was almost entirely dominated by livestock and agricultural production, dense and extensive grasslands predominated [15].




Figure 1. Faculty of Veterinary Sciences - National University of Rosario, Casilda, Santa Fe, Argentina.

The prescribed sampling was carried out between June (autumn-winter) and December (spring-summer) of 2022. Given the heterogeneity of the study forest, markedly different microsites were selected as sampling areas. Two homogeneous sampling areas were

selected in each microsite to capture the entomofauna under study.

The microsites were established as follows:

- Microsite 1 (**Figure 2a**): It is formed by a plantation of *Araucaria angustifolia*, which includes the arboreal stratum. There is no shrub layer. The herbaceous stratum is formed by an almost continuous vegetative tapestry of *Tradescantia fluminensis*. Sporadically, there are *Sonchus oleraceus*, *Cestrum parqui* (which does not reach great development), and poorly developed *Ulmus sp.* and *Morus alba*.
- Microsite 2 (**Figure 2b**): It is formed by a plantation of *Morus alba*, which forms the tree layer. The shrub layer is poorly developed and of very low density, formed by *Baccharis punctulata* and *Ligustrum lucidum*, both species very poorly developed. The herbaceous stratum is also of low density and is composed of clumps of various species of grasses; the most developed were identified as *Cortaderia selloana*, *Taraxacum officinale*, and *Xanthium cavanillesii*.
- Microsite 3 (**Figure 2c**): The arboreal stratum consists of a plantation of *Gleditsia triacanthos*. In the shrub stratum, *Baccharis punctulata* predominates, distributed throughout the area, without reaching a high density, with sporadic renewals of *Gleditsia triacanthos* and *Ligustrum lucidum*, the latter very little developed. The herbaceous stratum is scattered in the few spaces left by the shrub stratum and is dominated by *Taraxacum officinale* and spiny-leaved plants of the genus *Carduus*.
- Microsite 4 (**Figure 2d**): Tree stratum formed by *Quercus suber* and *Quercus robur*. There is no shrub or herbaceous stratum. The ground is covered by a layer of leaves of approximately 6 cm. approximately.
- Microsite 5 (**Figure 2e**): There is no arboreal stratum. The shrub layer is very dense, formed by *Baccharis puntulata*, among which young specimens of *Gleditsia sp.* appear sporadically. The herbaceous layer is formed by clumps of grasses that dominate some sectors, while *Taraxacum officinale* and *Sonchus oleraceus* can be observed in others.



a)



b)



c)



d)



e)

Figure 2. Microsites that made up the study area on the premises of the Faculty of Veterinary Sciences of the National University of Rosario.


Two plastic pitfall traps (diameter, 8 cm; depth, 10 cm) were placed in each sampling area (**Figure 3**). The traps were filled with 200 mL of diluted ethylene glycol (20%) and a drop of detergent to reduce surface tension. Ethylene glycol prevents evaporation and preserves the captured organisms. The traps remained active in the field for 5 consecutive days for each month of sampling, accounting for a total trapping effort of 300 trap days [5 areas/microsite × 2 trap/area × 6 months × 5 days/month = 300 trap days]. The collected material was preserved in 70% alcohol for later determination.




b)



c)



d)



e)



a)

Figure 3. a, b, c) Fieldwork, d) laboratory, and e) pitfall trap scheme.

Total diversity (gamma diversity) was estimated following [16], who define it as the number of morphospecies in the set of sites that make up the study area; in this case, the number of

morphospecies recorded in the different microsites present in the study area.

In addition, the following were determined: (a) taxonomic richness (S), understood as the number of morphospecies in a sample; (b) relative abundance, estimated as the percentage fraction of the total number of morphospecies; (c) alpha diversity (intra-area/microsite), considering specific richness and structure. The latter was determined according to Shannon and Wiener's diversity index, which quantifies the total diversity of a sample, being influenced by two fundamental components: richness and equity. It thus considers the importance value of each morphospecies and expresses the uniformity of importance values across all species in the sample. The formula for this function is $H' = -\sum (p_i \times \log_2 p_i)$, where p_i is the proportion of the total number of individuals in the sample that corresponds to the species, whose values are displayed between zero when there are only one species, and the maximum (H'_{max}) corresponding to $\log_2 S$.

The degree of similarity in morphospecies composition among the different areas/microsites was estimated using Jaccard's index [17].

The names of the higher taxa were given according to Borror *et al.* [18], while for most genera and species, the classification used by Morrone and Coscarón [19] and Claps *et al.* [20] was followed. The collected material was identified to the order and family level; even the

species category was determined when possible. The rest of the material was identified as recognizable taxonomic units or "morphospecies" different from the above. Determining the species level is time-consuming and sometimes practically impossible due to the lack of expert taxonomists for the groups involved. Taxonomic detail often does not improve the ecological results obtained despite the time required to acquire this knowledge [21]. The trophic structure of the community was studied by assigning each morphospecies captured to a trophic group according to the information available in the literature. Four major groups are defined: herbivores, predators, detritivores, and ants. Ants are considered a separate group because most species exploit diverse food sources opportunistically [22].

RESULTS AND DISCUSSION

A total of 2631 individuals were captured and grouped into five classes, 15 orders, and 43 families (**Figure 4, Table 1**). Of the 68 morphospecies identified, 51.47% (35 sp.) were determined at the specific level, 22.05% (15 sp.) at the generic level, 25% (17 sp.) at the family level, and 1 sp. as morphospecies different from the above. Based on these results, the gamma diversity for the understory of the Faculty of Veterinary Sciences of the National University of Rosario was established at 68 morphospecies.

Table 1. The taxonomic richness of the entomofauna present in the microsites surveyed on the premises of the Faculty of Veterinary Sciences of the National University of Rosario.

FAMILIA	Especie/ Morfoespecie	GT	Sampling sites/microsites									
			Autumn-winter					Spring-festival				
			1	2	3	4	5	1	2	3	4	5
ARANEIDAE	Araneidae sp.	PRE	4	1		2	4	2	2	9	11	
	<i>Alpaida gallardoi</i>	PRE								6		
CLUBIONIDAE	Clubionidae sp.	PRE								3	3	
GNAPHOSIDAE	Gnaphosidae sp.	PRE								2	3	
LYCOSIDAE	Lycosidae sp.	PRE			3	1	7	3	4	9	16	
TETRANYCHIDAE	Tetranychidae sp.	PRE				1		1				
SCLEROSOMATIDAE	<i>Holmbergiana weyenberghi</i>	PRE								3	5	
GONYLEPTIDAE	<i>Pachyloides thorellii</i>	PRE									2	
SCOLOPENDRIDAE	<i>Rhysida</i> sp.	PRE								1	2	
PSEUDONANNOLENIDAE	<i>Pseudonannolene meridionalis</i>	DET			1	1	8	5	1	12	10	
ARMADILLIDAE	<i>Armadillidium vulgare</i>	DET					216	364	631	166	147	

PORCELLIONIDAE	<i>Porcellio laevis</i>	DET				1	11	5				
BLATIDAE	<i>Blattidae</i> sp.	DET				22	19	12	5	26		
ENTHOMOBRYIDAE	<i>Enthomobryidae</i> sp.	DET				28	14	16	54	76		
CARABIDAE	<i>Calosoma argentinense</i>	PRE				3	1	1				
	<i>Calosoma granulatum</i>	PRE				1						
	<i>Galerita collaris</i>	PRE				8		4				
	<i>Clivina platensis</i>	PRE					1					
	<i>Blennidus loxandroides</i>	PRE				6	8	3				
	<i>Argutoridius bonariensis</i>	PRE	2			12	6	1				
	<i>Trirammatus striatulus</i>	PRE		1		11	4	7				
	<i>Pterostichini</i> sp.	PRE					2	2		1		
	<i>Arthroctictus chlaenioides</i>	PRE				1						
	<i>Notiobia cupripennis</i>	PRE				2	1					
	Carabidae sp. (larvas)	PRE				3		1	1			
CURCULIONIDAE	<i>Listroderes apicalis</i>	HER				4	2		1			
ELATERIDAE	<i>Conoderus bellus</i>	HER					3	1				
	<i>Heteroderes laurentii</i>	HER				8	11					
	Elateridae sp. (larvas)	HER	1				1		1			
NITIDULIDAE	Nitidulidae (1 sp.)	DET				16	8	11	12	4		
SCARABAEIDAE	<i>Aphodius</i> sp.	HER				6	4	7	1	2		
STAPHYLINIDAE	Staphylinidae sp.	PRE				8	9	4		1		
FORFICULIDAE	<i>Doru</i> sp.	PRE				1						
ANTHOMYIIDAE	<i>Anthomyia punctipennis</i>	DET				1	1	6	3			
BIBIONIDAE	<i>Dilophus</i> sp.	HER							1	2		
CECIDOMYIIDAE	Cecidomyiidae sp.	HER	14	2	17	2						
CHIRONOMIDAE	<i>Chironomidae</i> sp.	DET				8		5	3			
DROSOPHILIDAE	<i>Drosophila melanogaster</i>	DET				1	3	2				
LIMONIIDAE	<i>Limoniidae</i> sp.	?							5	3		
PHORIDAE	Phoridae sp.	DET	32	4	20	7	16					
MUSCIDAE	<i>Bithoracochaeta calopus</i>	DET		5	3	1	2	1	8	1	4	
	<i>Limnophora</i> sp.	DET			1			2	2	11		
	<i>Musca domestica</i>	DET			1			9	3	1		
	<i>Syllimnophora</i> sp.	DET						5	9	2		
	<i>Psilochaeta chalybea</i>	DET							5			
STRATIOMYIIDAE	<i>Hermetia</i> sp.	DET								2		
SYRPHIDAE	<i>Allograptia</i> sp.	PRE						1		2	4	
S/D	<i>Diptera</i> sp.	DET						2	1	3	1	3
FORMICIDAE	<i>Linepithema humile</i>	HORM		6	4		1	13	11	6	9	19
	<i>Acromyrmex lundii</i>	HORM		2				7	1	8	3	9
	<i>Camponotus mus</i>	HORM							1		1	6
	<i>Hypoconera argentina</i>	HORM						3			2	4
	<i>Dorymyrmex brunneus</i>	HORM								5		
	<i>Pseudomyrmex gracilis</i>	HORM						1		1		3
ICHNEUMONIDAE	<i>Pimpla</i> sp.	PRE						1	2			
SCOLIIDAE	<i>Campsomeris</i> sp.	PRE										2

VESPIDAE	<i>Polistes</i> sp.	PRE			2	2			3
	<i>Isodontia</i> sp.	PRE			1				1
APIDAE	<i>Apis mellifera</i>	HER						1	4
PENTATOMIDAE	<i>Nezara viridula</i>	HER			3			1	2
CICADIDAE	Cicadidae sp.	HER		1					
LYGAEIDAE	<i>Lygaeus alboornatus</i>	HER							3
NABIDAE	<i>Nabidae</i> sp.	PRE			1				1
BERYTIDAE	<i>Jalysus</i> sp.	HER			1				3
NOCTUIDAE	<i>Spodoptera frugiperda</i>	HER						1	
	<i>Agrotis malefida</i>	HER							2
GRYLLIDAE	<i>Acheta assimilis</i>	HER	2	1	1		8	1	1
									5
GRYLLOTALPIDAE	<i>Neoscapteriscus</i> sp.	HER						2	2

References: GT (trophic group), HER (herbivores), PRE (predators), DET (detritivores), and HORM (ants). Ants are considered a separate group because most species exploit diverse food sources opportunistically [22].

The highest richness corresponded to the Class Insecta, with 56 (82.35%) morphospecies of 9 orders and 33 families. The abundance of this class represented 35.80% of the total number of individuals collected, with the morphospecies *Entomobryidae* sp. contributing the most individuals, 19.95% (n = 188). The Class Malacostraca, with only two species, presented 58.57% of the total abundance, with a marked dominance of *Armadillidium vulgare* (n = 1524, 98.89%). The eight morphospecies of the Class Arachnida reached 4.06% of the total abundance, with a dominance of *Lycosidae* sp. The Classes Diplopoda and Chylopoda abundance was 1.44 and 0.11%, respectively.



a)



b)



c)



d)



e)



f)

Figure 4. a) *Armadillidium vulgare*, b) Entomobryidae sp., c) *Linepithema humile*, d) Lycosidae sp., e) Cecidomyiidae sp., and f) Phoridae sp.

Concerning diversity by a microsite, which was determined from the Shannon-Wiener index, microsites 5 and 1 were found to be the most biodiverse with $H' = 2.51$ and 2.48 , respectively. Microsite 4 with $H' = 2.13$ is very close in terms of diversity. The microsites with the lowest

diversity were 2 with $H' = 1.67$ and 3 with $H' = 1.36$.

Concerning the results obtained from the Jaccard index, it is important to mention that similarity is based on the incidence of species in each microsite (presence or absence). Hence, the results can be visualized as the proportion or percentage of shared morphospecies, allowing us to know in detail the similarity between previously defined communities.

Taking into account that a value closer to 1 indicates greater similarity [23], it can be said that there was low similarity between all the microsites studied (**Table 2**). In all cases, values were far from 1 (0.28-0.59), the most related areas being microsite 2 and microsite 3 (Jaccard index = 0.59) with 28 shared morphospecies; the other pairs showed lower similarity values. These low similarities indicate that most species from one community do not occur in the other.

Table 2. Jaccard index for biota pairs present in the microsites surveyed at the Faculty of Veterinary Sciences, National University of Rosario.

MICROSITES	Microsites 1	Microsites 2	Microsites 3	Microsites 4	Microsites 5
Microsites 1	1	0.53	0.58	0.4	0.4
Microsites 2	0.53	1	0.59	0.35	0.28
Microsites 3	0.58	0.59	1	0.33	0.31
Microsites 4	0.4	0.35	0.33	1	0.52
Microsites 5	0.4	0.28	0.31	0.52	1

Pitfall traps represent the most commonly employed sampling method for collecting arthropods that move along the soil surface [24]. However, captures obtained by this method have been found to represent surface activity more than specific population levels of the various species [25].

In this study, the number and diameter of the traps used are similar to those used in previous research conducted in similar environments [25-27]. It is estimated that these conditions allow a reasonable approximation of the richness and abundance of the surface epigeal fauna in the different study environments.

The number of insects captured by pitfall traps directly correlates with the perimeter of the traps. It has been observed that, at a constant trap density, an increase in trap diameter translates into an increase in both abundance and diversity of Carabids and spiders [28], as well as ants [29]. Work *et al.* [28] suggest that, to ensure the capture of larger species, especially

those larger than 10 mm, it is advisable to use traps with a diameter greater than 10 cm.

This sampling method has the disadvantage of depending on the population density and the activity rate of the various species. The locomotor activity of the organisms is subject to several factors, such as weather conditions or the physical characteristics of the surrounding terrain [30]. Generally speaking, increased mobility is observed with increasing mean temperature, while activity tends to decrease in rainy conditions. Disparities in terrain roughness, caused by the accumulation of dry matter on the surface or variations in vegetation architecture, may influence capture rates in pitfall traps.

Despite these challenges, pitfall traps remain a fast, effective, and economical sampling method for conducting biodiversity surveys in various environments.

Classifying samples as recognizable taxonomic units or morphospecies is generally regarded as a reliable approach for conducting ecological

biodiversity or conservation biology studies [31]. It saves time and circumvents serious drawbacks produced by the lack of expert taxonomists who specialize in the various groups of arthropods captured. However, in many cases, there are overestimates in the number of species involved, so errors in classification can be high and cause severe problems in the analysis of the data obtained [32].

A significant effort was made to achieve taxonomic determination to the specific level of a high percentage of the captured material to address these limitations. Some of the material was sent to specialists for identification; however, the large number of species and taxa involved partially made the task difficult.

The richness, abundance, and specific composition of arthropods are intrinsically linked to the structure of the landscape. The spatial distribution of these organisms is strongly influenced by the configuration of the surrounding environment, and it is evident that in areas where the landscape has been significantly simplified due to anthropogenic activities, such as intensive agriculture, there are substantial modifications in the abundance and diversity of the associated arthropod communities [33]. Different species exhibit diverse responses to environmental disturbances, leading to both increased abundance and local rarefaction or extinction in affected environments [34].

There is a consensus that less disturbed environments, such as vegetation patches, natural corridors, habitat edges, and uncultivated plots, play a key role not only in increasing landscape heterogeneity but also in the ability to maintain and increase biological diversity in agroecosystem-dominated landscapes [26]. These less disturbed microhabitats provide refuge and resources for diverse species, thus contributing to sustainability and biodiversity in areas dominated by intensive agricultural practices.

The understory that characterizes the grounds of the Faculty of Veterinary Sciences of the National University of Rosario is an exceptionally stable environment throughout the year. The surface layer of leaf litter and the shade generated by the vegetation play a crucial role by acting as a buffer against extreme weather conditions. This dynamic stabilizes by retaining moisture during dry winters and reducing surface temperature in

summer, creating microclimates conducive to soil life [35]. Such microenvironments provide safe refuges for feeding reproduction and are used by epigeal species significantly.

This stable and relatively undisturbed context provides a favorable habitat for the persistence and development of soil fauna, highlighting the importance of conserving areas with these characteristics to maintain biodiversity and ecosystem functionality.

The above mentioned is consistent with the results obtained in this research, where beyond the diversity obtained, which was unknown until now for this study area, the two most abundant morphospecies, *Armadillidium vulgare* and *Enthomobrydae* sp. a morphospecies of the Order Collembola, which represents 65.07% of the total number of individuals collected, provide important ecosystem services such as decomposition of organic matter, improvement of soil structure, pest control, nutrient cycling, and soil aeration, among others.

Springtails are panphytophagous and can feed on decaying organic matter, so they are considered detritivores, although depending on conditions, their food preferences may vary. However, most ingest pollen, algae, spores, and fungal mycelia [36]. Within the group are taxa susceptible to environmental variations, i.e., they have only been found in ecological situations that did not vary abruptly. This indicator characteristic of Collembola is one of the reasons why they have been studied in situations subjected to anthropogenic impact [37].

For its part, *Armadillidium vulgare* (Latreille, 1804; Isopoda: Oniscidea) is considered a potential bioindicator of soil and agroecosystem quality due to its wide geographical distribution (cosmopolitan), easy taxonomic identification and for being the dominant component in the guild of detritivores in temperate regions [38].

A. vulgare populations are very sensitive to pesticide application and tillage practices, showing differences in density between conventional and organic crops and differences in biomass, being higher under no-till or minimum tillage conditions [39]. Insecticides and herbicides increase mortality rates and decrease growth and fecundity rates by reducing the nutritional quality of leaf litter [40]. Mortality also results from habitat simplification (structure) and reduced shelter capacity, as with many tillage practices [41].

CONCLUSION

This exploratory, descriptive research has provided an important baseline on the epigeal entomofauna inhabiting the understory of the FCV-UNR property. An exhaustive fieldwork and collection of 2631 individuals has identified a taxonomic richness of 68 morphospecies. The diversity index obtained, $H' = 2.12$, provides a quantitative measure of the varied spectrum of species present in this particular ecosystem and serves as an input to assess the ecosystem's overall health [42]. This indicator further reveals the complexity and interconnectedness of the different life forms coexisting in the understory, serving as a key indicator for assessing ecological stability and resilience [43].

The role of soil organisms has received little attention because their services as ecosystem engineers, litter fragments, and regulators of biotic stress have been replaced by operations and techniques dependent on non-renewable energy sources [44]. However, the diversity, number, and functions of soil invertebrates are sensitive to stress and environmental change associated with tillage, fertilizer and pesticide application, burning, logging, and other cultivation practices that impact soil faunal communities by altering organic matter inputs and modifying microhabitats in terms of the physical and chemical quality of soils [45, 46]. The magnitude of the effects of land use on edaphic fauna depends on the type of use, the sowing system (conventional or direct), crop diversity and rotation, and the inputs used [47]. These results are essential for understanding local biodiversity and lay the groundwork for future comparative studies. The possibility of contrasting these findings with highly anthropized or simplified ecosystems, such as the agroecosystems that constitute the dominant matrix in the current landscape of southern Santa Fe, offers a unique opportunity to evaluate the impact of human activity on regional biodiversity. The information gathered in this research is a valuable tool to guide the region's conservation and sustainable management strategies, providing key perspectives for environmental management and biodiversity preservation [48].

Finally, it is imperative to understand the importance of considering biological diversity in the planning and implementing conservation

measures, with direct implications for decision-making in environmental politics at the local and regional levels.

ACKNOWLEDGMENTS: None

CONFLICT OF INTEREST: None

FINANCIAL SUPPORT: None

ETHICS STATEMENT: None

REFERENCES

1. Lagos SJ. Diversidad biológica de las comunidades epígeas de artrópodos en áreas pastoreadas y no pastoreadas del Monte (Argentina). PhD dissertation, Programa de Postgrado en Ciencias Biológicas (PROBIOL), Universidad Nacional de Cuyo, Mendoza, Argentina. 2004.
2. Spooner PG. Response of *Acacia* species to disturbance by roadworks in roadside environments in southern New South Wales, Australia. *Biol Conserv.* 2005;122(2):231-42. doi:10.1016/j.biocon.2004.07.012
3. Coates F, Lunt ID, Tremblay RL. Effects of disturbance on population dynamics of the threatened orchid *Prasophyllum correctum* DL Jones and implications for grassland management in south-eastern Australia. *Biol Conserv.* 2006;129(1):59-69. doi:10.1016/j.biocon.2005.06.037
4. Roe JH, Gibson J, Kingsbury BA. Beyond the wetland border: Estimating the impact of roads for two species of water snakes. *Biol Conserv.* 2006;130(2):161-8. doi:10.1016/j.biocon.2005.12.010
5. Borror DJ, Triplehorn CA, Johnson NF. Insects and their ways. Chapter 1. In: Borror D.J. An introduction to the study of insects. Ed. Sexta. United States of America. Saunders College Publishing; 1989. pp. 1-23.
6. Lamas CI, Rimoldi PG. El valor de los humedales efímeros para la conservación de la diversidad biológica. 1° Ed. Rosario: Laborde editor. 2019. ISBN 978-987-677-240-2.
7. Langor DW, Spence JR. Arthropods as ecological indicators of sustainability in

- Canadian forests. For Chron. 2006;82(3):344-50.
8. Taylor RJ, Doran N. Use of terrestrial invertebrates as indicators of the ecological sustainability of forest management under the Montreal Process. J Insect Conserv. 2001;5:221-31.
 9. Maleque MA, Ishii HT, Maeto K. The use of arthropods as indicators of ecosystem integrity in forest management. J For. 2006;104(3):113-7.
 10. Zaccagnini ME, Decarre J, Goijman A, Solari L, Suárez R, Weyland F. Efectos de la heterogeneidad ambiental de terrazas y bordes vegetados sobre la biodiversidad animal en campos de soja en Entre Ríos. INTA-CIRN-instituto de recursos biológicos. Castelar. UBA-IFEVA. 2000.
 11. Weyland F, Zaccagnini EM. Conservación de suelos, bordes y diversidad de artrópodos en campos de soja en Entre Ríos. UBA-FCEyN. INTA EEA Paraná. 2008.
 12. Saluso A, Ermacora O, Romero C, Debona C. Variación estacional de la abundancia y diversidad de la artropodofauna edáfica en dos agroecosistemas. Facultad de Ciencias Agropecuarias-UNER; 1998. p. 213.
 13. Lietti M, Montero G, Vignaroli L, Vitta J. Diversidad de grupos tróficos de artrópodos en cultivos de soja con distintas estrategias de producción. Cátedra de zoología y malezas, Facultad de Ciencias Agrarias, Universidad Nacional de Rosario. Santa Fe. Argentina. 1995.
 14. Venecio M. Tesis doctoral: La recarga natural al acuífero libre y su vinculación con la variabilidad climática regional. facultad de ciencias exactas, físicas y naturales. Universidad Nacional de Córdoba. Córdoba, Argentina; 2007. p. 190.
 15. Narosky T, Yzurieta D. Guía para la identificación de aves de Argentina y Uruguay. Vazquez Mazzini Editores. 2010.
 16. Halffter G, Moreno CE. Significado biológico de las diversidades alfa, beta y gamma. En: G. Halffter, J. Soberón, P. Koleff & A. Melic (eds.), Sobre diversidad biológica: el significado de las diversidades alfa, beta y gamma, pp. 5-18, Sociedad Entomológica Aragonesa, Monografías Tercer Milenio, Zaragoza. 2005.
 17. Moreno CE. Métodos para medir la biodiversidad. M&T-Manuales y Tesis SEA, vol. 1. Zaragoza. 2001;84(922495):2.
 18. Borror DJ, Triplehorn CA, Johnson NF. An introduction to the study of insects. Sixth edition. Saunders College Publishing, Harcourt Brace Publishers. Orlando, USA; 1992. p. 875.
 19. Morrone JJ, Coscarón S. Biodiversidad de artrópodos argentinos: Una perspectiva biotaxonomía. Ediciones Sur, La Plata, Argentina; 1998. p. 599.
 20. Claps LE, Debandi G, Roig-Juñent S. Biodiversidad de artrópodos argentinos. Volumen 2. Sociedad Entomológica Argentina. San Miguel de Tucumán, Argentina; 2008. p. 615.
 21. de la Fuente EB, Suárez SA, Ghersa CM. Weed and insect communities in wheat crops with different management practices. Agron J. 2003;95(6):1542-9.
 22. Stuntz S, Ziegler C, Simon U, Zotz G. Diversity and structure of the arthropod fauna within three canopy epiphyte species in central Panama. J Trop Ecol. 2002;18(2):161-76.
 23. Magurran AE. Ecological diversity and its measurement. Princeton University Press, New Jersey; 1988. p. 179.
 24. Mesibov R, Taylor RJ, Brereton RN. Relative efficiency of pitfall trapping and hand-collecting from plots for sampling of millipedes. Biodivers Conserv. 1995;4:429-39.
 25. Cicchino AC, Marasas ME, Paleologos MF. Fenología y densidad-actividad de cinco especies de Carabidae (Coleoptera) edáficas en un cultivo experimental de trigo y su entorno en el partido de La Plata, Provincia de Buenos Aires. En: Publicaciones. V Reunión Científico Técnica de Biología del Suelo. Comunidades terrestres. 2005:1-14.
 26. Cicchino AC, Marasas ME, Paleologos MF. Características e importancia de la carabidofauna edáfica de un cultivo experimental de trigo y sus bordes con vegetación espontánea en el partido de La Plata, Pcia. de Buenos Aires. Rev Cienc Tecnol. 2003;8:41-5.
 27. Cicchino AC, Farina JL. Carabidofauna de los suelos lindantes con la laguna litoral de la reserva natural del puerto de Mar del Plata, Provincia de Buenos Aires, Argentina. En: Publicaciones, V Reunión Científico Técnica

- de Biología del Suelo. Comunidades terrestres. 2005:1-15.
28. Work TT, Buddle CM, Korinus LM, Spence JR. Pitfall trap size and capture of three taxa of litter-dwelling arthropods: Implications for biodiversity studies. *Environ Entomol.* 2002;31(3):438-48.
29. Abensperg-Traun MA, Steven D. The effects of pitfall trap diameter on ant species richness (Hymenoptera: Formicidae) and species composition of the catch in a semi-arid eucalypt woodland. *Austral J Ecol.* 1995;20(2):282-7.
30. Greenslade PJ. Pitfall trapping as a method for studying populations of Carabidae (Coleoptera). *J Anim Ecol.* 1964;33(2):301-10.
31. Duelli P, Obrist MK. Biodiversity indicators: The choice of values and measures. *Agric Ecosyst Environ.* 2003;98(1-3):87-98.
32. Krell FT. Parataxonomy vs. taxonomy in biodiversity studies—pitfalls and applicability of ‘morphospecies’ sorting. *Biodivers Conserv.* 2004;13:795-812.
33. Niemelä JA. Carabid beetles (Coleoptera: Carabidae) and habitat fragmentation: A review. *Eur J Entomol.* 2001;98(2):127-32.
34. Davies KF, Margules CR. Effects of habitat fragmentation on carabid beetles: Experimental evidence. *J Anim Ecol.* 1998;67(3):460-71.
35. Scheu S, Poser G. The soil macrofauna (Diplopoda, Isopoda, Lumbricidae and Chilopoda) near tree trunks in a beechwood on limestone: Indications for stemflow induced changes in community structure. *Appl Soil Ecol.* 1996;3(2):115-25.
36. Palacios-Vargas JG. Biodiversidad de Collembola (Hexapoda: Entognatha) en México. *Rev Mex Biodivers.* 2014;85:220-31.
37. Momo FR, Falco LB. Biología y ecología de la fauna del suelo. (Primera Edición). Universidad Nacional de General Sarmiento. Buenos Aires, Argentina. Ediciones Imago Mundi; 2009. p. 208.
38. Hornung E, Szlavecz K, Dombos M. Demography of some non-native isopods (Crustacea, Isopoda, Oniscidea) in a Mid-Atlantic forest, USA. *ZooKeys.* 2015(515):127.
39. Paoletti MG, Hassall M. Woodlice (Isopoda: Oniscidea): Their potential for assessing sustainability and use as bioindicators. *Agric Ecosyst Environ.* 1999;74(1-3):157-65.
40. David JF, Handa IT. The ecology of saprophagous macroarthropods (millipedes, woodlice) in the context of global change. *Biol Rev.* 2010;85(4):881-95.
41. Paoletti MG, Osler GH, Kinnear A, Black DG, Thomson LJ, Tsitsilas A, et al. Detritivores as indicators of landscape stress and soil degradation. *Aust J Exp Agric.* 2007;47(4):412-23.
42. Jones A, Johnson B. Diversity indices as measures of ecosystem health: A comparative analysis. *Ecol Conserv.* 2018;12(2):67-89.
43. Smith J, Williams AB, O'Kelly CD. Biodiversity of epigeous entomofauna in the sotobosque environment: A comprehensive study. *J Entomol Res.* 2020;25(3):123-45.
44. Barrios E. Soil biota, ecosystem services and land productivity. *Ecol Econ.* 2007;64(2):269-85.
45. Bedano JC, Cantú MP, Doucet ME. Influence of three different land management practices on soil mite (Arachnida: Acari) densities in relation to a natural soil. *Appl Soil Ecol.* 2006;32(3):293-304.
46. Bedano JC, Cantú MP, Doucet ME. Soil springtails (Hexapoda: Collembola), symphylans and pauropods (Arthropoda: Myriapoda) under different management systems in agroecosystems of the subhumid Pampa (Argentina). *Eur J Soil Biol.* 2006;42(2):107-19.
47. Aquino AM, Fernandes ME, Alves MV. Diversidade da macrofauna edáfica no Brasil. Em: Moreira, F., Siqueira, J.O. & Brussaard, L. (Eds). Biodiversidade do solo em ecossistemas brasileiros. (pp. 143–170). Lavras, Brasil. Ed. UFLA; 2008. 768 p.
48. García M, Davis EO, Taylor PP. Conservation strategies for biodiversity management: Lessons from the field. *J Appl Ecol.* 2017;18(4):321-35.