

Prevalence and Associated Environmental Risk Factors of Malaria: Dansha Town and its Surroundings, Tsegedie District, Ethiopia

Abaynew Jemal Jenber^{1*}, Ahmed Abafita Abagero²

¹Department of Plant Sciences, Bahir Dar University, Bahir Dar, Ethiopia

²Tsegedie District Education Office, Dansha, Ethiopia.

ABSTRACT

Malaria is the main public health issue in Ethiopia since it infects more than 75% of the country. This study's goal was to evaluate the risk factors for malaria in Dansha town and the area around it as well as the prevalence of the disease. Structured questionnaires and a Health facility-based study of confirmed malaria cases were conducted to determine the five-year (2012–2016) trend of malaria prevalence. In order to evaluate the data, SPSS version 20 was used. There were 28.4% more people who had malaria overall. The major plasmodium species identified was Plasmodium falciparum, followed by Plasmodium vivax. The prevalence of malaria parasite infections was significantly associated with age, while it was non-significantly associated with sex. Although the prevalence of Plasmodium falciparum was 3.413 times more likely to be higher than the prevalence of Plasmodium vivax among the age group under five years, the prevalence of Plasmodium falciparum was 0.793 times less likely to be higher than the prevalence of Plasmodium vivax among the age group from five to nine years. About 95% of the respondents had adequate knowledge about malaria. At the time of this study, 82.7% of the respondents had: a practice of using impregnated mosquito nets, 92.4% had risk factors for mosquito breeding sites near their homes, and 20.9% had headache symptoms. Health workers and responsible administrators should focus on giving health education about the prevention and control activities of malaria and how individual households use insecticide-treated nets correctly.

Keywords: Environmental factors, Knowledge, Malaria, Household, Plasmodium, Prevalence.

HOW TO CITE THIS ARTICLE: Jenber AJ, Abagero AA. Prevalence and Associated Environmental Risk Factors of Malaria: Dansha Town and its Surroundings, Tsegedie District, Ethiopia. Entomol Appl Sci Lett. 2023;10(4):13-25. <https://doi.org/10.51847/2LBWv13s30>

Corresponding author: Abaynew Jemal Jenber

E-mail ✉ abujemal900@gmail.com

Received: 23/08/2023

Accepted: 04/12/2023

INTRODUCTION

Malaria is a parasitic disease caused by single-celled organisms called Plasmodium. It is transmitted to humans through the bite of infected mosquitoes. Malaria is a major cause of illness and death worldwide [1]. Malaria is a serious disease that affects millions of people around the world. In 2012, an estimated 3.4 billion people were at risk of malaria, and there were about 207 million cases and 627,000 deaths. Most of these deaths occurred in Africa. However, thanks to increased prevention and control measures, the mortality rate from malaria has decreased by 42% in the world and 49% in Africa [2].

Death and morbidity from Malaria have enormous social and economic costs [3, 4]. It causes poverty. Thus, it has a significant and measurable direct and indirect cost and has recently been shown to be a major constraint to economic development. Malaria may account for 40% of public health expenditure, 30%–50% of inpatient admissions, and up to 50% of outpatient visits in areas with high malaria transmission [5, 6]. It is estimated to cost Africa more than 12 billion United States dollars (USD) every year [7]. Additionally, malaria in Ethiopia has compelled people to live in the higher, less productive agricultural regions [8, 9]. Since agriculture is the foundation of the nation's economy and the peak season for malaria

transmission occurs between planting and harvesting seasons, the disease has a significant negative influence on the economy [10].

Malaria is a major public health concern in Ethiopia, as it is in many other tropical countries [11, 12], 75% of Ethiopia's landmass is malarial, and 68% of its population is estimated to be at risk of malaria infection [13].

In Ethiopia, areas at altitudes between 1,600 and 2,000 m above sea level are epidemic-prone hypoendemic zones of malaria [14, 15]. Malaria prevalence has been found to vary among locations, possibly because of the country's heterogeneous topography and weather conditions [16].

All population groups are affected by the erratic and seasonal transmission of malaria in Ethiopia. The key harvesting seasons of September to December and April to May coincide with the biannual malaria transmission peaks, which have substantial effects on the nation's subsistence economy. About 30% of all disability-adjusted life threats lost are due to malaria [17]. In 2009, malaria was the most common reason for people to seek medical attention in Ethiopia. It accounted for 12% of all outpatient visits and nearly 8.55% of all hospital admissions as per the Ethiopia Federal Ministry of Health (FMOH). The FMOH estimate that there are five to ten million clinical malaria cases and approximately 70,000 people die of malaria each year [10, 18]. It has also been documented in the nationwide child survival study that malaria affects school attendance by 20% and contributes to 47% of child deaths in Ethiopia [16]. However, given that a sizeable section of the population lacks access to health care services, these numbers may significantly underestimate the real burden of malaria in the nation.

Significant epidemics occur every five to eight years, with focal epidemics being the most common form. Large-scale malaria outbreaks struck 3368 places in 211 Districts in 2003 from April to December, resulting in two million clinical and confirmed cases and 3000 fatalities [16].

According to the Presidential Malaria Intervention (PMI), *Plasmodium falciparum* and *Plasmodium vivax* parasite prevalence in Ethiopia was 0.7% and 0.3%, respectively. According to the 2011 PMI draft report, 4.5% of children under the age of five tested positive for malaria

using rapid diagnostic methods (RDTs), whereas 1.3% of children under the age of five tested positive using microscopy. *Plasmodium falciparum* constituted 77% of these infections. Thus, malaria is one of the serious public health problems that have a devastating impact on the socio-economic development of a country with a high mortality rate. People living in rural Ethiopia, especially in lowlands are most of the time infected by malarial disease with less knowledge and prevention methods.

Climate change is expected to have a direct impact on malaria by changing the behavior and distribution of mosquitoes, as well as the length of the parasite's life cycle. It is also expected to have an indirect impact by changing the ecological relationships between the organisms involved in malaria transmission (the vector, parasite, and host) [1]. At the Regional level, 75% of the area is potentially malarious [19].

Like in other parts of Ethiopia, malaria intervention and control activities have been scaled up. Malaria, however, continues to be one of the primary causes of morbidity in the area despite minor declines in prevalence [20].

According to the 2016/2009 health profile of the region, malaria is a leading cause of outpatient visits and admissions; approximately 58% of all suspected and confirmed cases of malaria occurred [21].

Malaria is one of the major public health issues in the research region as a result of these difficulties. In order to determine the prevalence of malaria and risk factors in Dansha town and its environs, this study was started.

MATERIALS AND METHODS

Description of the study area

The study was conducted at Dansha town and surrounding kebeles (lower administration units), Tsegedie District, Ethiopia. There are 25 kebeles in Tsegedie District. Tsegedie District has three climatic zones: *Dega* (highland) accounts for 33%, *Weinadega* (mid-land) constitutes 44% and *Kola* (low-land) accounts for the remaining 27% [22]. The area has two major rivers; Mekezo and Kaza. Dansha town and the surrounding kebeles such as Yekatit kebele, Alemgenet kebele, and Selam kebele have a total population of 27, 850 and 6328 households residing in an area of 3234 km² making the population density of 16

people /km². The town population accounts for 63.6% of the study area¹ population. The temperature in the research region ranges from 30 – 40 °C. There is a District Hospital, a Health Center, four private clinics, three health posts, and six rural pharmacies in the research region. Dansha town has a total population of 17,712 and 4025 households and the surrounding administrative kebeles; namely, Yekatit kebele a population of 4555 and 1035 households, Alemgenet kebele a population of 4676 and 1063 households, and Selam kebele with a population of 907 and 205 households [23].

Study design, data collection instrument and procedure

The study was carried out in Dansha town and surrounding kebele to estimate malaria prevalence and to identify risk factors associated with it. This was done through structured questionnaires and health institution based treatment records from 2012-2016.

The data regarding the socio-demographic information and associated risk factors were gathered through means of a structured questionnaire and responses were obtained at the time of sample collection. Earlier to sample collection, precise explanations were given about the aim of the study for all households who are willing to participate in the study. Age, marital status, education level, knowledge, attitude, and practice regarding the participant's usage of bed nets were among the sociodemographic characteristics whose data were gathered. The general content of the questionnaire was well structured close-ended questionnaire. The questionnaire was prepared for each participant and filled by a representative of the sample selected households under the direct supervision of the data collector.

The questionnaire deals with an individual's socio-demographic information, knowledge and use of preventive measures, use of anti-malarial drugs, and other malaria-related issues.

Sampling size and sampling techniques

In order to calculate the sample size, a confidence interval of 95% was used, along with the lowest acceptable result of 3% and a total of 384 houses. This number was then multiplied by 10% to account for non-response, yielding a total of 422 households. 16 of the 422 families that were

expected to participate in the study were not included because they did not respond. A response rate of 96.2% was consequently attained. Four hundred and six volunteers participated in the study.

Thus, the sample size is determined by the following formula [24].

$$n = Z^2 P (1-p) / d^2 \quad (1)$$

Where:

n = sample size for households

p = Average prevalence

d = Error value

z = Confidence interval z (95%) = 1.96

n = 384 members of households obtained

Contingency = 10% = 38 therefore, 384 + 38 = 422

The study participants were selected using simple random sampling. Based on proportional sampling to population sizes in town and surrounding countryside kebeles, the study sample size was determined. In each household one randomly selected respondent was taken as a study subject. In this study, 270, 67, 70, and 15 respondents were selected from *Dansha town, Yekatit, Alemgenet, and Selam kebele's*, respectively.

Inclusion and exclusion criteria and study variables

All household dwellers in Dansha town and its surrounding countryside were included in this study. Dwellers that are not present at the time of the study, those who were not volunteers, and dwellers of the town or surrounding countryside that did not live for at least one year were excluded.

The dependent variable of this study was the prevalence of malaria in the study. The independent variables were: socio-demographic characteristics, health information, type of living area, insecticide treated bed net possession in households, and distance from mosquito breeding site.

Statistical analysis

Data was entered and analyzed by the SPSS version 20 software package. Descriptive statistics were used in the analysis for frequencies, percentages and binary logistic regression analyses were conducted to describe

findings. P-value < 0.05 was considered statistically significant. The data were described and presented using tables and figures.

RESULTS AND DISCUSSION

Socio-demographic characteristics of respondents
 In this study, 60.2% and 39.8% of respondents were males and females, respectively. The respondents' ages ranged from 18 to 67 years. With regard to marital status, about 31.8% of respondents were single, 60.7% were married, 5.2% were divorced and 2.4% were widowed. The distribution of the study participants based on their educational status showed that 43.4% of the respondents could not read and write, 7.6% of the respondents could read and write, 18.2% were in their elementary school levels, 19.9% were in their secondary school levels and 10.9% were holders of university diploma and above (Table 1).

Table 1. Socio-demographic characteristics of respondents in Dansha town and surroundings

Characteristics		Frequency	Percentage
Sex	Male	254	60.2
	Female	168	39.8
Age	18-27	136	32.2
	28-37	112	26.5
	38-47	74	17.5
	48-57	57	13.5
	58-67	43	10.2
Marital status	Single	134	31.8
	Married	256	60.7
	Divorced	22	5.2
	Widowed	10	2.4
Religion	Orthodox	369	87.4
	Muslim	53	12.6
Educational status	Unable to read and write	183	43.4
	Read and write	32	7.6
	Elementary	77	18.2
	High school	84	19.9
	College and above	46	10.9

Knowledge of respondents on malarial disease
 Knowledge of Respondents about malaria is shown in Table 2. About 95% of them heard about malarial diseases from different sources, such as Radio, Television, Health professionals, friends, and others. Many responders demonstrated adequate understanding about malaria, including the accurate connection

between the disease and mosquito bites, its possible effects, and the appropriate treatment choices. About 75.8% of respondents said they thought malaria could be spread from an infected person to a healthy person through mosquito bites; 6.9% said it couldn't; the remaining 5.2% said they didn't know; and the rest said it couldn't be spread from person to person. With regard to conditions that are associated with malaria infection, 84.1% of respondents knew that stagnant water and swampy areas were the main sites of mosquito breeding. Mosquitoes often bite at night, as almost 92.9% of respondents correctly identified. More over 75.4% of research participants thought malaria could be prevented, and almost all 85.3%- thought it could be cured. The most often reported malaria preventive strategies were the usage of bed nets and drying stagnant water. The government's enhanced efforts to promote health and the study's timing may be to blame for this. Regarding symptoms of malaria, the respondents said that about 10.9% fever, 19.4% shivering, 20.9% headache, 14.9% vomiting, 11.1% chill, 38.5% loss of appetite, 6.4% back pain, 6.6% joints ache, and 1.2% other as important symptoms of malaria in the research region.

Table 2. Knowledge of respondents about malaria at Dansha Town and its surroundings

Knowledge	Response	Frequency	Percentage
Have information on malaria	Yes	401	95%
	No	21	5%
Is malaria a health problem in the study area	Yes	398	94.3%
	No	24	5.7%
Malaria is a transmittable disease	Yes	337	79.9%
	No	44	10.4%
	I don't know	41	9.7%
Malaria is transmittable by mosquito bite	Yes	320	75.8%
	No	29	6.9%
	I don't know	22	5.2%
When do mosquitoes bites mostly	Night	392	92.9%
	Day	30	7.1%
Know the breeding sites of mosquitoes	Yes	355	84.1%
	No	67	15.9%
Is malaria a curable disease?	Yes	360	85.3%
	No	11	2.6%

	I don't know	51	12.1%
Is there any method to prevent malaria?	Yes	318	75.4%
	No	104	24.6%
Symptoms of malaria -			
	Fever	46	10.9
	- Chill	47	11.1
	- Headache	88	20.9
	- Shivering	82	19.4
	- Vomiting	63	14.9
	- Back pain	27	6.4
	- Joints ache	28	6.6
	- Loss appetite	36	8.5
	- Other	5	1.2

Risk factors associated with malaria

Table 3 showed that about 14.2% of the respondents did not use mosquito nets, while (85.8%) of the respondents used mosquito nets to protect themselves from the bite of mosquitoes. About 87.7% of the respondents do not use body oil like buzz off to prevent mosquito bites. Out of 422 respondents, 291(69.0%) did not report malaria-associated problems to local health workers. In the study area, about 390(92.4%) of respondents lived near mosquito breeding sites. With regard to housing conditions, 34.8% of them lived in houses that had thatched roofs, 57.3% lived in houses that had corrugated iron sheets, and 7.8% lived in houses that had concrete roofs. With regard to wall opening, 87.7% of them lived in houses with openings on the walls due to the high temperature of the area which may enhance mosquito entrance into the living room of households (**Table 3**).

Table 3. Risk Factors associated with Malarial and the practice of the community to Prevent Malarial Disease in Dansha Town and its surroundings

Risk Factor Variables	Response	Frequency	Percent
Use mosquito net	Yes	362	85.8
	No	60	14.2
Use body oil (buzz off)	Yes	52	12.3
	No	370	87.7
Report problems to local health workers	Yes	131	31.0
	No	291	69.0
Mosquito breeding site near the living room	Yes	390	92.4
	No	32	7.6
Type of roof of living room	Thatched roof	147	34.8
	Corrugated iron sheet	242	57.3

	Concert roof	33	7.8
Presence of a wall opening	Yes	370	87.7
	No	52	12.3
Prevention Variables			
	Drain breeding sites	317	75.1
Community Campaign Activities to control the environment from mosquito	Spray chemicals	4	0.9
	Clear vegetation	13	3.1
	Participate in health education	88	20.9
Medical treatment	Take prescribed anti-malaria drugs	323	76.5
	Use traditional medicine	99	23.5
Method currently used	Impregnated mosquito net	349	82.7
	Fumigation	73	17.3
	Not available	14	3.3
Reason for not using an Impregnated mosquito net	Not affordable	9	2.1
	Not aware of its use	50	11.8

Practice of the community to prevent malaria

Some practices of respondents to prevent malaria are shown in **Table 3**. About, 75.1 % drained breeding sites of mosquitoes. Among the respondents, 76.5% and 24.5% had a practice of taking prescribed anti-malaria drugs and traditional medicine, respectively. Three hundred forty-nine (82.7%) of respondents used impregnated mosquito nets while the remaining used fumigation. Out of those who used fumigation to prevent malaria, 3.3%, 2.21%, and 11.8% do not use mosquito nets and justified them as not available, not affordable, and did not know about their use, respectively.

Prevalence of malaria in Dansha town and its' surroundings from 2012-2016

Trends of malaria infection among patients admitted at Dansha town health center

From 2012 to 2016, 87.4% of *Plasmodium falciparum* and 12.6% of *Plasmodium vivax* were identified in clinical records from the Dansha District Health Center for Malaria. The overall prevalence of malaria in the whole health record

was 28.4%. The prevalence of malaria was 28.7% in 2012, 25.7% in 2013, 25.9% in 2014, 28.5% in 2015, and 32.8% in 2016 (Table 4). The highest annual cumulative malaria prevalence, 32.8%, was detected in 2016. This was much greater than in 2012, 2013, 2014, and 2015. In the past five years of health records, malaria has been prevalent throughout the year; the peaks in prevalence were observed mostly from May to December. In 2016, high malaria prevalence was

widespread throughout the year, while the lowest annual cumulative prevalence of 25.7% was detected in 2013. The prevalence of *plasmodium falciparum* cases increased from 2012 to 2014 by 56.2%, but in 2015, it decreased by 19.03%, and in 2016, it increased by 24.8% whereas the prevalence of *plasmodium vivax* increased from 2012-2015, but it was decreased in the 2016 by 18.6%.

Table 4. Trends of malaria disease among patients admitted at Dansha town health center recorded from 2012 - 2016

Year	Parameters	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total
2012	No. examined	645	572	502	732	713	664	510	626	625	701	722	907	7919
	Positive	202	131	112	223	205	157	134	151	212	250	239	260	2276
	<i>P. falciparum</i>	155	110	89	167	143	127	98	119	215	202	205	195	1825
	<i>P. vivax</i>	47	21	23	56	62	30	36	32	17	48	34	45	451
2013	No. examined	578	430	566	608	898	835	728	674	634	726	766	764	8207
	Positive	178	122	137	207	222	212	108	110	125	212	235	241	2109
	<i>P. falciparum</i>	165	98	127	191	215	199	88	92	102	198	221	220	1916
	<i>P. vivax</i>	13	24	10	16	7	13	20	18	23	14	14	21	193
2014	No. examined	725	676	754	822	864	777	568	478	467	688	674	656	8199
	Positive	247	204	216	226	238	204	116	89	84	128	174	202	2128
	<i>P. falciparum</i>	222	192	201	212	218	185	95	74	71	114	154	190	1928
	<i>P. vivax</i>	25	12	15	14	20	19	21	15	13	14	20	12	200
2015	No. examined	436	494	761	846	772	698	747	790	470	557	747	648	7966
	Positive	158	189	218	253	247	218	112	252	116	167	226	111	2267
	<i>P. falciparum</i>	126	143	194	212	215	209	92	219	85	145	193	91	1920
	<i>P. vivax</i>	32	46	24	41	32	9	20	33	31	22	33	20	347
2016	No. examined	586	682	674	699	793	772	668	601	626	724	800	826	8451
	Positive	205	215	219	221	253	285	229	253	207	227	213	247	2774
	<i>P. falciparum</i>	180	197	195	191	208	259	197	242	189	215	193	236	2502
	<i>P. vivax</i>	25	18	24	30	45	26	32	11	18	12	20	11	272

Annual confirmed malaria cases recorded at Dansha town and its surroundings

Over the course of five years, the proportion of species composition in the research locations displayed various trends. Figure 1 shows that relative to *Plasmodium vivax*, *Plasmodium falciparum* was more prevalent in the Dansha town and its surrounding kebeles location. Starting from 2012 to 2014, the *Plasmodium falciparum* was highly increased by 80.2%, 90.8%, and 91.1%, respectively. However, it decreased in 2015 by 84.7% and once again increased rises in 2016 by 90.2%. During 2012,

the *Plasmodium vivax* was highly increased by 19.8%, but it was decreases starting from 2013 to 2014 by 9.2% and 8.9% respectively, and once again became increased in 2015 by 15.3% and also decline in 2016 by 9.8%. The prevalence rate of malaria decreased in 2013 by 25.7% and increased in 2014 by 28.8% and again decreased in 2015 by 28.4% and finally increased by 32.8% in 2016.




Figure 1. Annual confirmed malaria cases recorded at Dansha town and its surroundings from 2012-2016

Monthly health recorded data of malaria prevalence from Dansha distinct health center from 2012-2016

Different trends were seen in 2012 with regard to the species makeup of the plasmodium in relation to the monthly average number of cases. The major peak transmission of *Plasmodium falciparum* mostly occurred in September and November, While the minor transmission of *Plasmodium falciparum* occurred in March (89 and July. Regarding to the monthly distribution of *Plasmodium vivax*, the major peak transmission occurred in May, but the minor transmission of *Plasmodium vivax* occurred in September (**Table 4**).

Different trends were observed in 2013 with relation to the distribution of the plasmodium species relative to the monthly average of the number of cases. The major peak transmission of *Plasmodium falciparum* occurred in December and November; While minor transmission of *Plasmodium falciparum* occurred in August and July. Regarding the monthly distribution of *Plasmodium vivax*, the major peak transmission occurred mostly in February and July, but minor peak transmission of *Plasmodium vivax* in May (**Table 4**).

In 2014, there were varied trends in the plasmodium species composition based on the monthly average of the number of cases. The major peak transmission of *Plasmodium falciparum* mostly occurred in January and May; While the minor transmission of *Plasmodium falciparum* occurred in September and August. Regarding the monthly distribution of *Plasmodium vivax*, the major peak transmission occurred mostly in July, but minor peak

transmission of *Plasmodium vivax* in February (**Table 4**).

There were several trends in 2015 with regard to the distribution of the plasmodium species in relation to the monthly average of the number of cases. The major peak transmission of *Plasmodium falciparum* mostly occurred in June and March; While minor transmission of *Plasmodium falciparum* occurred in September and December. Concerning the monthly distribution of *Plasmodium vivax*, the major peak transmission occurred mostly in May, but minor peak transmission of *Plasmodium vivax* in December (**Table 4**).

In 2016, there were various trends in terms of the species that make up the plasmodium population in relation to the monthly average of the number of cases. The foremost peak transmission of *Plasmodium falciparum* mostly occurred in August; While the slight transmission of *Plasmodium falciparum* occurred in January. Regarding the monthly distribution of *Plasmodium vivax*, the foremost peak transmission occurred mostly in May, but minor peak transmission of *Plasmodium vivax* in August and December (**Table 4**).

Age, sex, and species breakdown of confirmed malaria cases in Dansha town and its surroundings

The prevalence of *Plasmodium falciparum* was 68.9% in children under the age of five. In contrast, this age group had a 31.1% *Plasmodium vivax* prevalence. The overall *Plasmodium falciparum* proportion for the age group of five to nine years was 90.5%, whereas the overall *Plasmodium vivax* proportion for the same age group was 9.5%. For the age group of ten to fourteen years, the overall *Plasmodium falciparum* proportion was 85.2%, whereas the overall *Plasmodium vivax* proportion was 14.8%. Similarly, the overall proportion of *Plasmodium falciparum* for the age group from fifteen and above years was 87.5 88.4%, while the overall proportion of *Plasmodium vivax* for the age group from fifteen and above years was 11.6% (**Table 5**).

Plasmodium falciparum was found in higher concentrations in the years 2013 (90.8%) and 2014 (90.6%), but it was found in lower concentrations in the years 2012 (80.2%) and 2015 (84.7%). A higher proportion of

Plasmodium vivax was registered in 2012 (19.8%) and 2015 (15.3%), whereas a lower proportion of *Plasmodium vivax* was seen in 2013 (9.2%). Males were more affected than females

by malaria parasites, but differ year to year. The infection rates among males were (66.3%) and females were (33.7%) (Table 5).

Table 5. Distribution of confirmed malaria cases by Age and Plasmodium species type at Dansha town and its surroundings from 2012 – 2016

Age	Species type	Years					Total
		2012	2013	2014	2015	2016	
< 5	<i>P. falciparum</i>	71.6	68.9	68.4	60.2	76.7	68.9
	<i>P. vivax</i>	28.4	31.1	31.6	39.8	23.3	31.1
5-9	<i>P. falciparum</i>	84.9	90.6	92.9	90.6	91.7	90.5
	<i>P. vivax</i>	15.1	9.4	7.1	9.4	8.3	9.5
10-14	<i>P. falciparum</i>	78.6	92.4	91.7	76.9	87.6	85.2
	<i>P. vivax</i>	21.4	7.6	8.3	23.1	12.4	14.8
≥ 15	<i>P. falciparum</i>	79.9	91.5	90.6	88.9	91.4	88.4
	<i>P. vivax</i>	20.1	8.5	9.4	11.1	8.6	11.6

Association of prevalence of malaria parasite infections with age and sex at Dansha town and surroundings from 2012 – 2016

The findings demonstrated that the prevalence of malaria parasite infection or species plasmodium was highly correlated with all age groups of malaria cases (P < 0.05), whereas sex was not significantly associated with malaria prevalence (Table 6). Although the prevalence of *Plasmodium falciparum* was 3.413 times [OR=3.413, 95%CI: 2.727 – 4.271] more likely higher than the prevalence of *Plasmodium vivax* among the age group under five years and the

prevalence of *Plasmodium falciparum* were 0.793 times [OR = 0.793, 95%CI: 0.680 – 0.923] less likely to the prevalence of *Plasmodium vivax* among the age group from five to nine years (Table 6).

The prevalence of *Plasmodium falciparum* was 1.315 times [OR = 1.315, 95%CI: 1.155 – 1.497] more likely to the prevalence of *Plasmodium vivax* among the age groups from ten to fourteen years. The prevalence of *Plasmodium falciparum* was 0.132 times more likely than the prevalence of *Plasmodium vivax* of malaria species among people from fifteen years (Table 6).

Table 6. Binary Logistic Regression Analysis of Malaria Prevalence by Age and sex in Dansha Town and Surroundings from 2012 – 2016

Age group	Species	Frequency	Percentage	Odds Ratio (95% CI)	P-value
< 5 years	<i>P. falciparum</i>	287	2.5%	3.413 (2.727 – 4.271)	0.000
	<i>P. vivax</i>	129	1.2%	1	
	Subtotal	416	3.7%		
5-9 years	<i>P. falciparum</i>	2510	21.7%	0.793(0.680 – 0.923)	0.003
	<i>P. vivax</i>	262	2.3%	1	
	Subtotal	2772	23.9%		
5-14 years	<i>P. falciparum</i>	2685	23.3%	1.315(1.155 – 1.497)	0.000
	<i>P. vivax</i>	465	4.1%	1	
	Subtotal	3150	27.3%		
≥ 15 Years	<i>P. falciparum</i>	4609	39.9%	0.132	0.000
	<i>P. vivax</i>	607	5.3%	1	
	Subtotal	5216	45.2%		
Sex					
Male	<i>P. falciparum</i>	6816	88.9%		0.996
	<i>P. vivax</i>	850	11.1%		1.000
	Total	7666			

	<i>P. falciparum</i>	3275	84.2%	0.996
Female	<i>P. vivax</i>	613	15.8%	-
	Total	3888		

In this study, a total of 11554 confirmed malaria cases were reported from 40742 suspected malaria examined with an overall prevalence of malaria of 28.4% from 2012-2016. This was significantly higher than the survey results of Ethiopia's national malaria prevalence from 2009 (0.3%) to 2013(0.2%), and a health record-based study conducted in the southern part of Ethiopia reported that 11.45% of 2521 febrile people who visited the health center for treatment tested positive for malaria [25]. It was much higher than the survey results national malaria prevalence in Ethiopia from 2015(0.3%) to 2011(1.3%) in areas below 2,000 m.a.s.l. 11.5% of 384 feverish people who attended the health center in northern Ethiopia, according to a research based on health facilities, tested positive for malaria¹. This might be due to the seasonal fluctuation of temperature [17], poor housing conditions of house hold dwellers [26], and inappropriate bed net usage [27]. Other reasons might be altitude variation and climate conditions that may contribute to the breeding site of the malaria parasite. Human behavior, the prevalence of malaria parasites, medical facilities, housing circumstances, jobs, community knowledge, attitudes, and behaviors about malaria transmission, prevention, treatment, and mosquito control are only a few of the variables that affect the spread of malaria [28].

However, an investigation of the incidence of malaria during a ten-year period (2002–2011) was undertaken in the northwest of Ethiopia showed that 39.6% confirmed malaria cases were reported out of 59, 208 cases [1].

The current study result was similar to a hospital-based survey conducted in the Arba-Minch town hospital where 34.7% of confirmed malaria cases were reported in all patients who had visited the outpatient department of the hospital and 60% of the total confirmed malaria cases were males [29].

Regarding *Plasmodium falciparum*, the prevalence rate in this study was 87.4%. This finding is much higher than a study done in Arsi Negelle health center, southern Ethiopia [25], Sire health center, East Wellaga zone, Western

Ethiopia [30] and Kola Diba, North Gondar, and Northwest Ethiopia [1] which were 19.8%, 66.1%, and 75%, respectively. The reason for the fluctuation of the prevalence of *P. falciparum* in different study areas might be due to seasonal fluctuation of ambient temperature [31], topography and humidity [32], availability of mosquito breeding sites, and absence of an integrated management system [33].

The prevalence of *P. vivax* in the current study was 12.6%. This is much lower than a study conducted in Arsi Negelle health center, southern Ethiopia, Sire health center, East Wellaga zone, Western Ethiopia [30] and Kola Diba, North Gondar, and Northwest Ethiopia [1]. which were 74%, 30.5%, and 25%, respectively.

In Ethiopia, insecticidal net coverage has increased to 53.3% [34]. Inpatient malaria cases have been reduced by 73% and child deaths by 62% since ITNs have been distributed nationwide, particularly long-lasting insecticide-treated nets (LLITNs) [35]. A According to research from rural Côte d'Ivoire [36] and other regions of Africa, high coverage of bed nets, especially those treated with an insecticide (ITNs and LLINs), resulted in a drop in malaria death and morbidity and reduces transmission [37, 38]. The findings of the present study revealed that 85.8% of respondents used ITNs. Several studies indicated that high stability in one place and the use of ITNs significantly reduce the proportion of malaria morbidity and mortality [38]. ITNs possession prevalence in this study was relatively higher than the study conducted in Iran that is 20% [39]. Without correct use, the mere existence of ITNs in the various houses of the respondents may not be sufficient to protect people against malaria morbidity. This may potentially follow from these observations. Inspections of ITNs have revealed difficulties in using them, including hanging them on the wall without using them and keeping them in packs. The results of this study also showed that respondents from the study region had a high level of general knowledge of malaria. Nearly majority of the responders (99.3%) have sufficient understanding about malaria. Seventy-five.8% of those surveyed connected malaria to

mosquito bites. This may be in contrast to a research by 30 in which 63.4% of participants connected malaria to mosquito bites. This outcome was comparable to those of other African nations like Kenya (77.3%) and Kampala (84%) [31].

Knowledge of a community is of paramount importance for the adoption of good practices for health improvement and malaria prevention. Deepen our understanding of the contextual factors that influence people's practice of malaria prevention and treatment to improve the effectiveness of malaria control interventions. For those residing in malaria-risk regions, health education is crucial to enhancing their understanding of the disease, their perceptions of it, and their behavior when seeking out health care [40]. Educated and uneducated people alike know malaria can be prevented. This might mean that malaria knowledge is not fully acquired through formal education. Knowledge of the disease might also come from non-formal sources, such as informal education.

However, the educational status and knowledge of most people living in malarious areas is limited. This might contribute to limited knowledge on how to use impregnated net, malarial treatment, and other preventive measures [41]. As a result, the current research region has a high prevalence of malaria, or 28.4%.

Regarding other aspects of knowledge of respondents on malaria, 75.8%, and 92.9% replied that malaria is caused by mosquito bites and biting at night, respectively. This result is consistent with research from Arba-Minch, where 98.2% of respondents said a mosquito bite causes malaria [42] and the Assosa zone, where 95% of respondents said insects bite at night [43]. However, this finding is higher than a study conducted in the Assosa zone by 34, which has shown that less than half (47.5%) of the study participants responded that malaria is caused by mosquito bites. Additionally, 79.9%, 75.4%, and 85.3% of participants in this research thought that malaria could be spread, avoided, and treated, respectively. About 76.5% of them replied that the use of modern medicine is an adequate treatment for malaria. This is almost in agreement with a study by [44].

Several factors determine malaria transmission, such as human behavior, the presence of malaria

parasites, and social and health facility factors. A community's knowledge, attitudes, and practices (KAPs) about malaria causation, transmission, treatment-seeking behavior, and mosquito control activities play a key role in determining malaria transmission rates [40]. Among the factors associated with malaria infections, not using mosquito nets and living around stagnant water seemed to have had an association with the prevalence of malaria in the study area.

With regard to housing conditions, 57.4% of respondents lived in houses that had corrugated iron sheets. The finding showed an association between the presence of malaria with the type of houses. Those who lived in houses having roofs with corrugated iron sheets were 15 times at higher risk, of getting malaria than those who lived under concrete roofs. The cause may be related to building characteristics that are linked to a high prevalence of malaria in the research region, such as an earth roof, tethering cattle within the home, the existence of windows, open eaves, the absence of a separate kitchen, and the presence of a single sleeping chamber. Poorly built homes have a higher risk of contracting malaria, which may be related to their appropriateness for an abundance of indoor vectors. Since homes are where malaria vectors mostly bite and rest, better housing may lower the likelihood of indoor malaria transmission in the research location. However, according on geography, socioeconomics, and specific household circumstances, housing conditions and their influence on the abundance of indoor vectors may differ [26].

CONCLUSION

The study revealed that different households in Dansha town and the surrounding countryside are highly exposed to malaria. In the research region, malaria was present generally at a rate of 28.4% greater than the prevalence of the country (4.1%) and much higher than the survey results of national malaria prevalence in Ethiopia from 2009 (0.3%) to 2013(0.2%). This might be an indication that different protective measures should be taken in Dansha town and the surrounding countryside by the community as well as the responsible government authorities. Public awareness formation could be important through strong health addition activities

incorporated with the community. Identification and elimination of causes of malaria including breeding sites should be strengthened. Reinforced health education should be implemented and it might be a promising as well as tangible measure to further reduce malarial infection. Dynamic participation of the community should be exploited in environmental organizations to control the transmission of malaria.

ACKNOWLEDGMENTS: We acknowledge Dansha Hospital for providing data and the community who participate in the study.

CONFLICT OF INTEREST: None

FINANCIAL SUPPORT: None

ETHICS STATEMENT: None

REFERENCES

1. Abebe Alemu DM. Ten year trend analysis of malaria prevalence in KolaDiba, Northwest Ethiopia. *Parasit Vectors*. 2012;5(1):173.
2. WHO. World malaria report. National Press Club Washington. 2013.
3. Irhan HB, Oran IB. Value Changes in National Currency in Foreign-Dependent Economies & Turkey Example in The Context of Crises. *J Organ Behav Res*. 2022;7(2):82-94.
4. Komov M, Panko J, Spector A, Stepanyan T, Tumanov E. The EAEU Potential and the Interests of Member-States in Cooperation to Ensure Sustainable Foreign Economic Strategies. *J Organ Behav Res*. 2021;6(1):166-72.
5. Taha MS, Elbasheir ME, Abakar MA, Abdallah EI, Elbashier MM, Omer AE, et al. The Impact of COVID-19 on Blood Coagulation Profile among Sudanese Hospitalized Adult Patients. *J Biochem Technol*. 2022;13(3):67-70.
6. Karpov VY, Medvedev IN, Komarov MN, Dorontsev AV, Kumantsova ES, Mikhailova OD. Possibilities of Students' Health Improvement through Physical Training in the Aquatic Environment. *J Biochem Technol*. 2021;12(4):67-71.
7. Union A. Five year review of the Abuja calls for accelerated action towards universal access to HIV/AIDS. Tuberculosis, and Malaria services. (Vol. 6). Geneva, Switzerland, Special session of the African Union Conference of Ministers of Health. Progress Report (2014 - 2010). 2010.
8. Abdaltif A, Abdallah MD, Yagowb MY, Mustafa MG, Alameen TA, Attar AO, et al. Transfusion Related Hepatitis C Virus Antibodies and Possible Risk Factors in Healthy Blood Donors. *Pharmacophore*. 2021;12(5):32-7.
9. Florina MG, Mariana G, Csaba N, Gratiela VL. The Interdependence Between Diet, Microbiome, and Human Body Health - A Systemic Review. *Pharmacophore*. 2022;13(2):1-6.
10. PMI. Malaria Operational Plan (MOP) Ethiopia FY 2011. Saving lives in Africa. 2011.
11. Muth Lakshmi K, Lakshmi K, Kannan A, Aniyam Y. Evaluation of novel MicroRNA profile-21 and 191 in oral leukoplakia and oral squamous cell carcinoma in comparison with healthy tissues – A cross-sectional study. *Clin Cancer Investig J*. 2021;10(6):275-82.
12. Dogan I, Khanmammadov N, Ahmed MA, Yıldız A, Saip P, Aydinler A, et al. Crizotinib in Metastatic ALK mutant Non-small Cell Lung Cancer Patients: A Single Centre Experience. *Clin Cancer Investig J*. 2022;11(3):25-9.
13. FMOH. Guideline for Malaria epidemic prevention and control in Ethiopia. (Second Ed.). Addis Ababa, Ethiopia: Federal Republic of Ethiopia, Ministry of Health. 2012.
14. Singh G, Goel N, Singh A, Gera R. Study of Factors Affecting the Time to Diagnosis and Treatment in Pediatric Acute Leukemia Patients- A study from India. *Clin Cancer Investig J*. 2022;11(3):35-40.
15. Ashwin SP, Sinduja P, Jayaraman S, Priyadarshini R. Assessment of 8-OHDG, SOD, and GPx Levels in OPMD and OSCC Patients. *Clin Cancer Investig J*. 2023;12(2):39-43.
16. FMOH. Ethiopia National Malaria Indicator Survey 2015. Technical summary Report Addis Ababa, Ethiopia. 2016.
17. Ayele TT. The risk factor indicators of malaria in Ethiopia. *Int J Med Med Sci*. 2013;5(7):335-47.

18. FMOH. Health and Health related indicators (Vol. 2). Expansion of Health extension program, FMOH, Addis Ababa, Ethiopia. 2009.
19. Gehebreyesus WK. Community based Malaria control in Tigray, Northern Ethiopia. *Parasitological*. 199;41(1-3):367-71.
20. WHO. Systems for the early detection of malaria epidemics in Africa: an analysis of current practices and future priorities, country experiences. Geneva, Switzerland: World Health Organization. 2006.
21. WHO. Global Malaria Program Roll Back Malaria Partnership. 2010.
22. Tsegedie. Tsegedie District, Finance and Economic Development Office, Dansha Town, Ethiopia. 2015.
23. CSA. Central Statistical Authority of Ethiopia. The Population and Housing Census of Ethiopia. 2010.
24. Naing WT. Practical issues in calculating the sample size for prevalence rate. *Arch Orofasc Sci*. 2006;1:9-14.
25. Hailemariam GS. Trend analysis of malaria prevalence in Arsi Negelle Health Center, Southern Ethiopia. *J Infect Dis Immun*. 2015;7(1):1-6.
26. Abebe MB. Impact of housing condition on indoor -biting and indoor -resting *Anopheles arabiensis* density in a high land area, Central Ethiopia. *Malar J*. 2013;12(1):393.
27. Killen ST. Preventing childhood malaria in Africa by protecting adults from mosquitoes with insecticide treated nets. *Public Lib Sci Med*. 2015;4(7):229.
28. Alemu A, Tsegaye W, Golassa L, Abebe G. Urban malaria and associated risk factors in Jimma town, south-west Ethiopia. *Malar J*. 2011;10(1):1-0.
29. Asnake. A five year (2014 -2010) Hospital based retrospective study of malaria in Gidole and its vicinity, Southern Ethiopia. Addis Ababa, Addis Ababa University, Addis Ababa, Ethiopia. 2010.
30. Gemechu T, Samuel A, Yewhalaw D. Ten years trend analysis of malaria prevalence and its correlation with climatic variables in Sibu Sire District, east Wollega zone, Oromia regional state, Western Ethiopia: a retrospective study. *Sci Technol Arts Res J*. 2015;4(4):99-105.
31. O'Meara BP. Effect of a fall in malaria transmission on morbidity and mortality in Kilifi. *Hancet Infect Dis*. 2016;372(9649):1555-62.
32. Atieli ZG. Topography as a modifier of breeding habitats and concurrent vulnerability to malaria risk in the western Kenya high lands. *Parasit Vectors*. 2011;4:241.
33. Woyessa DW. Prevalence of malaria infection in Butajira area, South-central Ethiopia. *Malar J*. 2012;11:84.
34. FMOH. Ethiopia National Malaria Indicator Survey 2007. Technical summary Report Addis Ababa, Ethiopia. 2008.
35. Otten AMB. Initial evidence of reduction of malaria cases and deaths in Rwanda and Ethiopia due to rapid scale- up of malaria prevention and Treatment. *Malar J*. 2009;8:1-8.
36. Henry ASY. Protective efficacy of lambda cyhalothrin treated nets in *Anopheles gambiae* pyrethroid resistant areas of cote d'Ivoire. *Am J Trop Med Hyg*. 2013;73(5):859-64.
37. Curtis MC. Insecticide treated bed- nets for malaria mosquito control. *J Am Mosq Control Assoc*. 2006;22(3):501-6.
38. Lengeler C. Insecticide treated bednets and curtains for preventing malaria. *Cochrane Database Syst Rev*. 2004;2.
39. Mohammad MA. Baseline results of the first malaria indicator survey in Iran at household level. *Malar J*. 2011;10(1):277.
40. Peterson BLS. Individual and household level factors associated with malaria incidence in a highland region of Ethiopia: a multilevel analysis. *Am J Trop Med Hyg*. 2009;80(1):103-11.
41. Kudom MB. The potential role of educational system in addressing the effect of inadequate knowledge of mosquitoes on the use of insecticide treated nets in Ghana. *Malar J*. 2010;9:256.
42. Astatkie. Knowledge and practice of malaria prevention methods among residents of Arba- Minch zuria district, Southern Ethiopia. *Ethiop J Health Sci*. 2010;20(3):185-93.

43. Legesse TA. Knowledge, attitude, and practice about malaria transmission and its preventive measure among households in urban areas of Assosa Zone, Western Ethiopia. *Ethiop J Health Dev.* 2007;21(2):157-65.
44. Paulander LH. Knowledge, attitudes, and practices about malaria in rural Tigray, Ethiopia. *Global Health Action.* 2009;2(1):1839.