

Fauna and Species Diversity of Ground Beetles (Coleoptera, Carabidae) in Meadows

Alexander Ruchin^{1*}, Sergei Alekseev², Anatoliy Khapugin¹, Mikhail Esin¹

¹Joint Directorate of the Mordovia State Nature Reserve and National Park «Smolny», Saransk, Republic of Mordovia, Russia.

²Ecological club «Stenus», Kaluga, Kaluga region, Russia.

ABSTRACT

The species diversity of ground beetles (Carabidae) was studied in the meadow ecosystems in the center of European Russia (Nizhniy Novgorod region and the Republic of Mordovia). Seventeen localities were studied. All meadow biotopes were divided into four types: dry meadows, dry meadows adjacent to forest shelter-belts, wet floodplain meadows, and floodplain meadows affected by livestock grazing. The highest species diversity was in dry meadows adjacent to forest shelter-belts (65 species) and wet floodplain meadows (62 species). The lowest number of species was in floodplain meadows affected by livestock grazing (24 species). Forty ground beetle species have been identified in dry meadows. Wet floodplain meadows had the highest Shannon's index, and the lowest Simpson index. The ground beetle fauna had high values of the Simpson and Berger-Parker indices in dry meadows. Only two species dominated in dry meadows, while four to seven species dominated in other habitats. According to the Jaccard similarity index, the most similar species composition of ground beetles was in dry meadows and dry meadows adjacent to forest shelter-belts. By reducing the number of species and specimens of ground beetles, trampling has a great effect on the fauna of floodplain meadows affected by livestock grazing.

Keywords: Abundance, Fauna, Ground beetle, Meadows, Species.

HOW TO CITE THIS ARTICLE: Ruchin A, Alekseev S, Khapugin A, Esin M. Fauna and Species Diversity of Ground Beetles (Coleoptera, Carabidae) in Meadows. Entomol Appl Sci Lett. 2021;8(3):28-39. <https://doi.org/10.51847/Nv94GSLSkN>

Corresponding author: Alexander Ruchin

E-mail ✉ ruchin.alexander@gmail.com

Received: 19/05/2021

Accepted: 27/08/2021

INTRODUCTION

Meadows are a type of vegetation characterized by the dominance of perennial herbaceous plants, mainly cereals, and sedges, under conditions of sufficient or excessive moisture. The presence of stands and turf is a common feature for all meadows. By location, there are three main meadow groups. Continental meadows are located on plains outside river floodplains. They are divided into dry and lowland meadows. Floodplain meadows are located in river valleys; they are flooded during floods. Mountain meadows are located above the upper border of the forest. In different worldwide regions, meadows are rich in perennial plant species [1] including threatened [2] and invasive [3] ones. They provide areas for

the inhabitation of birds [4], mammals [5, 6], reptiles [7], invertebrates [8-10], and others. The various meadow types have different abilities to recover after disturbance due to differences in carbon stock values [11, 12]. Since ancient times, meadows experienced exposure from human activities including unlimited grazing, plowing, afforestation, urbanization. It has resulted in a decline in areas covered by natural meadows [13].

Changes in open biocenoses (e.g. meadows, steppes, pastures) have been recently observed in many regions of the world [14-18]. The transformation of the vegetation cover has an ever-increasing impact on the ground beetle fauna in the biocenoses which are bioindicators of the ecosystem status [19-22]. These ground layer inhabitants of biogeocenoses are found in sufficient quantities in a wide variety of

landscapes (open and closed), biocenoses including territories of various destruction degrees. In recent decades, the anthropogenic transformation of meadow ecosystems (annual grass fires, unreasonable plowing of land, significant grazing, uncontrolled haying, afforestation of fields and meadows) has shown high rates [23-26]. The process of environmental impact leads to the disruption of the natural habitat, the emergence of secondary forest communities, and, consequently, changes in the natural ranges of some ground beetle species and population structure [16, 27-32]. This study is aimed to investigate the meadow ecosystems' ground beetle fauna in the center of European Russia.

MATERIALS AND METHODS

Description of biotopes

All biotopes were divided into four types. The main criteria for their distinguishing were the moisture degree, the presence of forest or afforestation near the biotope (within 100 m), anthropogenic impact in the form of grazing farm animals.

I – Dry meadows. These habitats are grazed meadows and abandoned fields located at sites on dry sandy soils. A certain degree of aridity is expressed in such biotopes. The following plant species have been found in these habitats: *Achillea millefolium* L., *Calamagrostis epigejos* (L.) Roth, *Bromus inermis* Leyss., *Trifolium arvense* L., *Trifolium pratense* L., *Artemisia vulgaris* L., *Lathyrus pratensis* L., *Leucanthemum vulgare* (Vaill.) Lam., *Matricaria matricarioides* (Less.) Porter, *Dactylis glomerata* L., *Cirsium arvense* (L.) Scop., *Agrimonia eupatoria* L., *Cichorium intybus* L., *Pimpinella saxifraga* L., *Astragalus danicus* Retz., *Fragaria viridis* Weston, *Carex spicata* Huds.

II – Dry meadows adjacent to forest shelterbelts. These are similar to the previous type but these differ by presence of *Betula*-formed or

mixed shelterbelts in 30–50 m apart of the meadows. The following plant species have been found in these habitats: *Cirsium arvense* (L.) Scop., *Consolida regalis* Gray, *Matricaria matricarioides*, *Achillea millefolium*, *Calamagrostis epigejos*, *Bromus inermis*, *Thlaspi arvense* L., *Brassica rapa* L., *Polygonum aviculare* L., *Cyanus segetum* Hill, *Leucanthemum vulgare* (Vaill.) Lam., *Viola arvensis* Murray, *Phleum pratense* L.

III – Wet floodplain meadows. They are located in floodplains of streams and small rivers on the relatively wet sandy and sandy-loam soils. The following plant species have been found in these habitats: *Rumex confertus* Willd., *Taraxacum officinale* (L.) Weber ex F.H.Wigg., *Carex spicata* Huds., *Carex vulpina* L., *Agrimonia eupatoria* L., *Bromus inermis*, *Tussilago farfara* L., *Dactylis glomerata* L., *Cichorium intybus* L., *Stellaria media* (L.) Vill., *Echium vulgare* L., *Scorzoneroidea autumnalis* (L.) Moench, *Trifolium hybridum* L., *Alchemilla* sp., *Scirpus sylvaticus* L., *Carex acuta* L.

IV – Floodplain meadows affected by livestock grazing. These habitats are similar to the previous type. But IV type differs from type III by the higher ground compaction due to grazing impact on the soil and vegetation cover.

Scientific names were used according to The PlantList database (<http://www.theplantlist.org/>).

Collection methods

We collected the material using pitfall traps from April to September in 2009, 2010, 2014, 2019. They were represented by 0.5-liter cups with 4% formalin solution. There were ten traps in each locality, they were installed in one line with a distance of two to three meters between them. In total, we studied 17 localities located in the Nizhny Novgorod region and the Republic of Mordovia (**Figure 1**). The study was conducted in each locality with only one line, consisting of ten traps.




Figure 1. Study Territory. Places, where Material Is Collected, Are Indicated by Red Dots.

Data analysis

The diversity analysis of ground beetles in the ecosystems was evaluated by the following diversity indexes: Shannon-Wiener (H'), which considers equal weight to the rare and abundant species, and Simpson's index ($1-D$), which is sensitive to changes in the most abundant species composition [33]. The uniformity among the coleopterans caught in the five sampling sites was calculated with the Berger and Parker index. Mathematical processing was carried out in Microsoft Excel. The tables show the average values.

In total, we have collected more than 3400 specimens during 4750 trap-days of the study.

The following scheme was adopted to characterize the numerical abundance of species: dominant species had numerical abundance exceeded 5%; subdominant species had numerical abundance from 2% to 5%; inconsiderable in number species had numerical abundance from 1% to 2%; rare species had numerical abundance less than 1%. The dynamic density of beetles was recognized as many beetle specimens caught per 100 traps per one day (exemplars / 100 trap-days, hereafter – ex./100 trap-days).

Species identification of ground beetles was carried out using the identification tables from the works [34, 35]. We used the Carabidae

system according to the website of the Zoological Institute of RAS [36] and based on a catalog of Kryzhanovskij *et al.* [37]. The nomenclature is given according to the catalog of the Palearctic beetles [38]. The asterisk "*" marks the species that were first registered in the Republic of Mordovia. The material is stored in the collection of the Mordovia State Nature Reserve (Pusha settlement, Russia).

RESULTS AND DISCUSSION

We caught 110 species of ground beetles belonging to 35 genera (**Table 1**) in all meadows. Four species were first registered in the Republic of Mordovia. The genera *Amara* (19 species), *Harpalus* (15 species), and *Pterostichus* (12 species) included the largest number of species. Wide holarctic and palearctic ranges are characteristic of many species. The most number of collected ground beetles were common and mass species in the forest-steppe zone in European Russia.

Table 1. The Fauna and Dynamic Density (ex./100 Trap-days) of Species Collected in Four Types of Meadow Biotopes

Species	Dry meadows	Dry meadows adjacent to forest shelter-belts	Wet floodplain meadows	Floodplain meadows affected by livestock grazing
Cicindelinae				
<i>Cylindera germanica</i> (Linnaeus, 1758)		7.58		
<i>Cicindela campestris</i> Linnaeus, 1758	0.07			0.61
Carabinae				
<i>Leistus ferrugineus</i> (Linnaeus, 1758)		0.08		
<i>Notiophilus germinyi</i> Fauvel, 1863		0.33		
<i>Loricera pilicornis</i> (Fabricius, 1775)		0.08	1.05	
<i>Clivina fossor</i> (Linnaeus, 1758)			0.07	
<i>Calosoma maderae</i> (Fabricius, 1775)		0.25		
<i>Carabus cancellatus</i> Illiger, 1798	0.07	0.33	0.46	
<i>Carabus clathratus</i> Linnaeus, 1761	0.07		0.13	
<i>Carabus glabratus</i> Paykull, 1790			0.13	
<i>Carabus granulatus</i> Linnaeus, 1758			1.96	
<i>Trechus secalis</i> (Paykull, 1790)	0.07	0.67	1.18	
<i>Trechus quadristriatus</i> (Schränk, 1781)		0.17		
<i>Bembidion biguttatum</i> (Fabricius, 1779)			1.70	
<i>Bembidion dentellum</i> (Thunberg, 1787)			1.24	
<i>Bembidion gilvipes</i> Sturm, 1825			0.07	
<i>Bembidion lampros</i> (Herbst, 1784)		0.25	0.13	
<i>Bembidion properans</i> (Stephens, 1828)		2.08	0.78	
* <i>Bembidion schuppelii</i> Dejean, 1831			0.72	
<i>Bembidion quadrimaculatum</i> (Linnaeus, 1760)		1.17		
<i>Patrobus atrorufus</i> (Ström, 1768)		0.08	0.07	
<i>Stomis puniceus</i> (Panzer, 1796)		0.08	0.26	
<i>Poecilus cupreus</i> (Linnaeus, 1758)	0.15	5.25	2.03	0.45
<i>Poecilus lepidus</i> (Leske, 1785)	0.22	0.83		
<i>Poecilus sericeus</i> Fischer von Waldheim, 1824	0.07			
<i>Poecilus versicolor</i> (Sturm, 1824)	6.76	1.58	4.44	1.06
<i>Pterostichus anthracinus</i> (Illiger, 1798)			13.46	
* <i>Pterostichus cursor</i> (Dejean, 1828)			0.13	
<i>Pterostichus gracilis</i> (Dejean, 1828)			4.90	0.15
<i>Pterostichus macer</i> (Marsham, 1802)		0.17		
<i>Pterostichus mannerheimii</i> (Dejean, 1831)			0.13	
<i>Pterostichus melanarius</i> (Illiger, 1798)	0.07	13.08	5.75	

<i>Pterostichus minor</i> (Gyllenhal, 1827)	0.07		0.39	
<i>Pterostichus niger</i> (Schaller, 1783)	0.07	0.17	1.44	
<i>Pterostichus nigrita</i> (Paykull, 1790)			3.20	
<i>Pterostichus oblongopunctatus</i> (Fabricius, 1787)	0.15	0.42		
<i>Pterostichus strenuus</i> (Panzer, 1796)		0.08	0.46	
<i>Pterostichus vernalis</i> (Panzer, 1796)			1.96	
<i>Calathus erratus</i> (C. Sahlberg, 1827)	1.25	13.25		3.79
<i>Calathus fuscipes</i> (Goeze, 1777)	2.94	3.83	3.01	0.61
<i>Calathus melanocephalus</i> (Linnaeus, 1758)	1.32	3.33	1.31	2.27
<i>Dolichus halensis</i> (Schaller, 1783)		0.08		
<i>Limodromus krynickii</i> (Sperk, 1835)		0.17		
<i>Agonum duftschmidii</i> J. Schmidt, 1994			3.07	
<i>Agonum fuliginosum</i> (Panzer, 1809)		1.92	0.13	
<i>Agonum viduum</i> (Panzer, 1796)			0.33	
<i>Oxytelus obscurus</i> (Herbst, 1784)		0.58	0.39	
<i>Synuchus vivalis</i> (Illiger, 1798)		0.25	0.20	
<i>Amara aenea</i> (De Geer, 1774)	0.81	2.92	0.33	1.36
<i>Amara apricaria</i> (Paykull, 1790)		0.25		
<i>Amara aulica</i> (Panzer, 1796)	0.22	0.08	0.72	
<i>Amara bifrons</i> (Gyllenhal, 1810)		2.42	0.07	
<i>Amara communis</i> (Panzer, 1797)	1.25		4.12	0.76
<i>Amara consularis</i> (Duftschmid, 1812)		0.67	0.07	
<i>Amara equestris</i> (Duftschmid, 1812)	8.24	2.08	0.65	3.03
<i>Amara eurynota</i> (Panzer, 1796)		0.08		
<i>Amara fulva</i> (Müller, 1776)				0.15
<i>Amara ingenua</i> (Duftschmid, 1812)	0.22			0.30
<i>Amara littorea</i> C.G. Thomson, 1857		0.08		
<i>Amara montivaga</i> Sturm, 1825		1.08		
<i>Amara nitida</i> Sturm, 1825	0.07	0.08		
<i>Amara ovata</i> (Fabricius, 1792)		20.58		0.15
<i>Amara plebeja</i> (Gyllenhal, 1810)	0.07		0.13	
<i>Amara praetermissa</i> (C.Sahlberg, 1827)			0.07	
<i>Amara similata</i> (Gyllenhal, 1810)		0.67		
<i>Amara sprete</i> Dejean, 1831			0.13	
<i>Amara tibialis</i> (Paykull, 1798)		0.08		
<i>Acupalpus meridianus</i> (Linnaeus, 1761)			0.07	
<i>Anisodactylus binotatus</i> (Fabricius, 1787)	0.07			
<i>Anisodactylus nemorivagus</i> (Duftschmid, 1812)		5.25		
<i>Anisodactylus signatus</i> (Panzer, 1796)		0.08		0.61
<i>Harpalus affinis</i> (Schränk, 1781)	0.59	0.92		1.06
<i>Harpalus calathoides</i> Motschulsky, 1844				1.36
<i>Harpalus calceatus</i> (Duftschmid, 1812)		0.17		
<i>Harpalus distinguendus</i> (Duftschmid, 1812)	0.66	0.67		
<i>Harpalus griseus</i> (Panzer, 1796)			0.07	
<i>Harpalus latus</i> (Linnaeus, 1758)	2.94	0.08	1.31	
<i>Harpalus luteicornis</i> (Duftschmid, 1812)		0.17		
<i>Harpalus progrediens</i> Schaubberger, 1922	0.37			0.91
<i>Harpalus pumilus</i> Sturm, 1818		0.08		

<i>Harpalus rubripes</i> (Duftschmid, 1812)	0.59	3.25	0.59	1.21
<i>Harpalus rufipes</i> (DeGeer, 1774)	0.74	1.75	2.42	1.52
<i>Harpalus smaragdinus</i> (Duftschmid, 1812)				0.45
<i>Harpalus tardus</i> (Panzer, 1796)				0.45
<i>Harpalus xanthopus winkleri</i> Schauburger, 1923	0.07	3.67	2.16	
<i>Harpalus zabroides</i> Dejean, 1829		0.17		
<i>Ophonus azureus</i> (Fabricius, 1775)		2.08		
<i>Ophonus cordatus</i> (Duftschmid, 1812)	0.07	4.08		
* <i>Ophonus diffinis</i> (Dejean, 1829)		0.33		
<i>Ophonus puncticeps</i> Stephens, 1828	0.07			
<i>Ophonus rufibarbis</i> (Fabricius, 1792)			0.20	0.15
<i>Ophonus stictus</i> Stephens, 1828	0.07			
* <i>Ophonus subquadratus</i> (Dejean, 1829)		11.83		
<i>Panagaeus bipustulatus</i> (Fabricius, 1775)	0.07	3.50	0.07	0.15
<i>Panagaeus cruxmajor</i> (Linnaeus, 1758)			0.13	
<i>Callistus lunatus</i> (Fabricius, 1775)	0.15			
<i>Chlaenius nigricornis</i> (Fabricius, 1787)			5.82	
<i>Oodes helopioides</i> (Fabricius, 1792)			0.13	
<i>Licinus depressus</i> (Paykull, 1790)	0.07	0.17	0.59	
<i>Badister bullatus</i> (Schränk, 1798)		0.58		
<i>Badister lacertosus</i> Sturm, 1815			0.72	
<i>Badister meridionalis</i> Puel, 1925		0.50		
<i>Badister peltatus</i> (Panzer, 1796).			0.26	
<i>Badister unipustulatus</i> Bonelli, 1813		0.17	3.86	
<i>Lebia chlorocephala</i> (Hoffmann, 1803)	0.07	0.08	0.07	
<i>Lebia cruxminor</i> (Linnaeus, 1758)	0.66	0.08	0.07	
<i>Philorhizus sigma</i> (Rossi, 1790)			0.07	
<i>Microlestes maurus</i> (Sturm, 1827)			0.13	
<i>Microlestes minutulus</i> (Goeze, 1777)	0.22			
<i>Cymindis angularis</i> Gyllenhal, 1810	0.07			0.15
Total number of exemplars	433	1547	1249	150
Shannon index	2.52	3.14	3.29	2.78
Simpson index (1-D)	0.14	0.07	0.06	0.08
Berger and Parker index	0.26	0.16	0.16	0.17
Number of species	40	65	62	24

Only nine ground beetle species were captured in all the studied meadow biocenoses (8.2% of the total fauna): *Poecilus cupreus*, *Poecilus versicolor* (meadow mesophile), *Amara aenea*, *Amara equestris*, *Harpalus rubripes*, *Harpalus rufipes*, *Calathus fuscipes*, *Calathus melanocephalus*, and *Panagaeus bipustulatus* (all of them are eurybionts). Fifty-four species (49.1%) were found in only one meadow type. Dominant species and their number differed in biocenoses. *Amara equestris* and *Poecilus versicolor* (25.9% and 21.2%, respectively) dominated with considerable numerical abundance in dry meadows. From these

calculations, it is seen that these two dominant species accounted for almost half of the studied specimens in dry meadows in terms of numerical abundance. This was reflected in the calculation results of the dominance indices. Four dominant species were found in dry meadows adjacent to forest shelter-belts: *Amara ovata* 16.0%, *Pterostichus melanarius* 10.1%, *Ophonus subquadratus* 9.2%, and *Calathus erratus* 10.3%. All these species are meadow mesoxerophiles and eurybionts. Five species dominated in wet floodplain meadows: *Pterostichus anthracinus* 16.5%, *Pterostichus melanarius* 7.0%, *Pterostichus gracilis* 6.0%,

Chlaenius nigricornis 7.1%, and *Poecilus versicolor* 5.4% (forest species were also present). Seven species dominated in floodplain meadows affected by livestock grazing: *Amara equestris* 13.3%, *Harpalus rufipes* 6.7%, *Harpalus rubripes* 5.3%, *Amara aenea* 6.0%, *Harpalus calathoides* 6.0%, *Calathus erratus* 16.7%, and *Calathus melanocephalus* 10.0%. Such a diverse composition of dominants and subdominants (we will not give their names) indicated a high level of species in abundance in three communities and is an indicator of a wide variety of ground beetle complexes.

In separate meadow biocenoses, the number of ground beetle species varied from 24 to 65 (**Table 1**). The Shannon index was the highest in wet floodplain meadows, and the Simpson index was minimal, i.e. maximum species diversity was recorded in this site with minimal dominance of species (**Table 1**). In addition, a high Shannon index was recorded in dry meadows adjacent to forest shelter-belts. The minimum value of this index was obtained in the dry meadow fauna.

It is known that an increase in the Simpson index and the Berger-Parker index means a decrease in biocenose diversity and an increase in the dominance of some species [39]. A

significant increase in these indices was obtained in dry meadows. This means that a significant decrease in biodiversity while increase in dominance degree of 1–2 species occurs in dry meadows as opposed to dry meadows adjacent to forest shelter-belts. As we indicated earlier, *Amara equestris* and *Poecilus versicolor* were such dominant species.

Cluster analysis based on Jaccard similarity index showed that the most similar species composition of ground beetles was in dry meadows and dry meadows adjacent to forest shelter-belts (**Figure 2**). Species diversity was 40 species in the first biotope while it was significantly higher and amounted to 65 species in the second one (**Table 1**). This similarity under significant differences in the fauna of these biotopes can be explained by the fact that a certain number of species that prefer to keep under the tree crowns and not migrate towards open meadow stations are present in dry meadows adjacent to forest shelter-belts. Floodplain meadows affected by livestock grazing differed most significantly from all other meadow biotopes. Intensive grazing has a serious effect on the ground beetle fauna in local natural conditions.




Figure 2. The Similarity of Four Meadow Biotope Types based on Jaccard Index: I – Dry Meadows; II – Dry Meadows Adjacent to Forest Shelter-belts; III – Wet Floodplain Meadows; IV – Floodplain Meadows Affected by Livestock Grazing

The species diversity of the genera *Amara*, *Harpalus*, and *Pterostichus* was considerable being represented by a total of 46 species. A

similar dependence was noted in meadow biotopes in the north and southwest of European Russia [40]. An analysis of the data

showed that when forming the population of ground beetles, the main importance belongs to the species groups of open habitats (meadows, meadow fields, and fields groups).

Dry meadows differ from other types of studied biotopes by the moisture regime. They are less humid and, therefore, herbaceous Poaceae species grow there much more frequently, as well as weed species. Typically, such meadows do not mow, and they do not serve as pastures. The ground beetle species composition is formed in such conditions by species of the meadow complex and eurybionts [40]. Similar results were obtained in our studies.

Forest edges and forest shelter-belts can also affect insect diversity by offering additional habitats such as wintering grounds, summer hibernation sites, mating sites, or feeding places. Such ecotones can reduce migration, change the daily and seasonal movements of insects. An increase in the species diversity of insects adjacent to the biotope forest is the most likely result [41-43]. This is exactly what was obtained in our conditions. The presence of forest shelter belts near dry meadows remarkably increases the species diversity of ground beetles that can use these belts for their life.

Wet floodplain meadows have a good herb layer and moderate humidity in our study sites. The species diversity of ground beetles was quite high in our research (62 species). A similar number of species was recorded in floodplain meadows in Belarus [44], Kirov region in Russia [45], Masovian Lowland in Poland [46]. On the other hand, a more considerable species diversity of ground beetles was found in floodplain meadows in the Ryazan region (center of European Russia), where authors attributed the obtained results to the heterogeneity of the relief and the strongly mosaic vegetation diversity [47]. The proximity of habitats with different soil and plant conditions ensures the migration and exchange of species, which also increases the species diversity of ground beetles both in general and in individual habitats [47]. In our conditions, wet floodplain meadows were quite similar to each other and did not differ in the diversity of vegetation.

In the center of European Russia, floodplain meadows affected by livestock grazing are summer pastures for these herbivores. They are

usually used during the growing season. Cattle grazing is carried out daily leading to a gradual degradation of the meadow system. It is known that under excessive soil pressure, floodplain meadows affected by livestock grazing can be severely crushed with mechanical damage to the soil cover and partial destruction of vegetation [48, 49]. An important role is played by a significant amount of manure, which may also not equally affect different insect groups, but not as obvious as trampling [50]. We think that in such biotopes the decrease in the species diversity of ground beetles is explained by extremely strong grazing.

CONCLUSION

Thus, 110 species of ground beetles belonging to 35 genera were recorded in meadow biocenoses in the Nizhniy Novgorod region and Republic of Mordovia. The genera *Amara*, *Harpalus*, and *Pterostichus* are represented by the highest species richness. The basis of meadow carabid fauna consisted of eurybionts and meadow species. The highest species diversity was noted in dry meadows adjacent to forest shelter-belts and wet floodplain meadows. Forest shelter-belts created heterogeneity in uniform meadow biotopes and therefore increased the number of species and abundance of ground beetles. Wet floodplain meadows are distinguished by well-developed herb layers and humidity, which also affects the species diversity of ground beetles. The lowest number of species was in floodplain meadows affected by livestock grazing. We attribute this to the significant trampling and the soil cover degradation in these meadows. In wet floodplain meadows, the Shannon index was the highest, while the Simpson index was the lowest. The ground beetle fauna had high values of the Simpson and Berger-Parker indices in dry meadows. Only two species dominated in dry meadows, while four to seven species dominated in other habitats. Such a diverse composition of dominants and subdominants indicated a high level of species abundance in three communities, and it is an indicator of a wide variety of ground beetle complexes. The highest similarity of species composition of ground beetles was noted between dry meadows and dry meadows adjacent to forest shelter-belts according to the Jaccard similarity

index. Floodplain meadows affected by livestock grazing differed significantly from all other types of meadow biotopes.

ACKNOWLEDGMENTS: None

CONFLICT OF INTEREST: None

FINANCIAL SUPPORT: None

ETHICS STATEMENT: None

REFERENCES

- Mseja GA, Furael BB, Lyakurwa GJ, Martin EH. Herbaceous vegetation communities around lake Manyara, Tanzania: response to environmental gradients. *Nat Conserv Res.* 2020;5(4):55-64. doi:10.24189/ncr.2020.056
- Suleymanova GF, Boldyrev VA, Savinov VA. Post-fire restoration of plant communities with *Paeonia tenuifolia* in the Khvalynsky National Park (Russia). *Nat Conserv Res.* 2019;4(Suppl.1):57-77. doi:10.24189/ncr.2019.048
- Dubovik DV, Skuratovich AN, Miller D, Spiridovich EV, Gorbunov YuN, Vinogradova YuK. The invasiveness of *Solidago canadensis* in the Sanctuary «Prilepsky» (Belarus). *Nat Conserv Res.* 2019;4(2):48-56. doi:10.24189/ncr.2019.013
- Assandri G, Bogliani G, Pedrini P, Brambilla M. Species-specific responses to habitat and livestock management call for carefully targeted conservation strategies for declining meadow birds. *J Nat Conserv.* 2019;52:125757. doi:10.1016/j.jnc.2019.125757
- Horncastle VJ, Chambers CL, Dickson BG. Grazing and wildfire effects on small mammals inhabiting montane meadows. *J Wildl Manage.* 2019;83(3):534-43. doi:10.1002/jwmg.21635
- Kerekes V, Ozogány K, Sándor I, Vegvari Z, Czető C, Nyírő B, et al. Analysis of habitat use, activity, and body condition scores of Przewalski's horses in Hortobagy National Park, Hungary. *Nat Conserv Res.* 2019;4(Suppl.2):31-40. doi:10.24189/ncr.2019.029
- Bakiev AG, Gorelov RA, Klenina AA. Post-fire abundance and age composition dynamics of *Lacerta agilis* (Reptilia, Lacertidae) in the Orenburg State Nature Reserve (Russia). *Nat Conserv Res.* 2019;4(Suppl.1):105-9. doi:10.24189/ncr.2019.047
- Hoyle H, Norton B, Dunnet N, Richards JP, Russell JM, Warren P. Plant species or flower colour diversity? Identifying the drivers of public and invertebrate response to designed and annual meadows. *Landsc Urban Plan.* 2018;180:103-13. doi:10.1016/j.landurbplan.2018.08.017
- Uligova TS, Gedgafova FV, Gorobtsova ON, Tsepkova NL, Rapoport IB, Tembotov RKh, et al. Meadow biogeocenoses in the subalpine belt of the Kabardino-Balkaria State High-Mountain Reserve (Central Caucasus). *Nat Conserv Res.* 2019;4(2):29-47. doi:10.24189/ncr.2019.012 [in Russian]
- Dedyukhin SV. Phytophagous beetles (Coleoptera: Chrysomelidae and Curculionoidea), protected and recommended for protection in the regions of the Middle Volga and the Urals. *Nat Conserv Res.* 2020;5(2):1-27. doi:10.24189/ncr.2020.013 [in Russian]
- Gebremedihin KM, Birhane E, Tadesse T, Gbrewahid H. Restoration of degraded drylands through exclosures enhancing woody species diversity and soil nutrients in the highlands of Tigray, Northern Ethiopia. *Nat Conserv Res.* 2018;3(1):1-20. doi:10.24189/ncr.2018.001
- Dusaeva GK, Kalmykova OG, Dusaeva NV. Fire influence on dynamics of above-ground phytomass in steppe plant communities in the Burtinskaya Steppe (Orenburg State Nature Reserve, Russia). *Nat Conserv Res.* 2019;4(Suppl. 1):78-92. doi:10.24189/ncr.2019.050
- Hurka H, Friesen N, Bernhardt KG, Neuffer B, Smirnov S, Shmakov A, et al. The Eurasian steppe belt: Status quo, origin and evolutionary history. *Turczaninowia.* 2019;22(3):5-71. doi:10.14258/turczaninowia.22.3.1
- Tchabovsky A, Savinetskaya L, Surkova E. Breeding versus survival: proximate causes of abrupt population decline under environmental change in a desert rodent, the midday gerbil (*Meriones meridianus*). *Integr Zool.* 2019;14(4):366-70.

15. Fedorov NI, Zharkikh TL, Mikhailenko OI, Bakirova RT, Martynenko VB. Forecast changes in the productivity of plant communities in the Pre-Urals Steppe site of Orenburg State Nature Reserve (Russia) in extreme drought conditions using NDVI. *Nat Conserv Res.* 2019;4(Suppl.2):104-10. doi:10.24189/ncr.2019.044
16. Cicort-Lucaciu AŞ. Road-killed ground beetles prove the presence of *Carabus hungaricus* (Coleoptera: Carabidae) in North-Western Romania. *Nat Conserv Res.* 2020;5(3):134-8. doi:10.24189/ncr.2020.035
17. Perazza G, Decarli M. Monitoring of *Cypripedium calceolus* (Orchidaceae) in the Adamello-Brenta Natural Park (Italy). *Nat Conserv Res.* 2020;5(Suppl.1):178-84. doi:10.24189/ncr.2020.045
18. Bakka SV, Kiseleva NY, Shestakova AA. Current Status, Problems and Prospects of Conservation of Meadow Steppes in the Nizhny Novgorod Region. *IOP Conf Ser Earth Environ Sci.* 2020;543(1):012014. doi:10.1088/1755-1315/543/1/012014
19. Rainio J, Niemelä J. Ground beetles (Coleoptera: Carabidae) as bioindicators. *Biodivers Conserv.* 2003;12(3):487-506. doi:10.1023/A:1022412617568
20. Venn SJ, Kotze DJ, Lassila T, Niemelä JK. Urban dry meadows provide valuable habitat for granivorous and xerophilic carabid beetles. *J Insect Conserv.* 2013;17(4):747-64. doi:10.1007/s10841-013-9558-8
21. Ruchin AB, Alekseev SK, Khapugin AA. Post-fire fauna of carabid beetles (Coleoptera, Carabidae) in forests of the Mordovia State Nature Reserve (Russia). *Nat Conserv Res.* 2019;4(Suppl.1):11-20. doi:10.24189/ncr.2019.009
22. Bondarenko AS, Zamotajlov AS, Belyi AI, Khomitskiy EE. Fauna and ecological characteristics of ground beetles (Coleoptera, Carabidae) of the Nature Sanctuaries «Prichernomorskiy» and «Tuapsinskiy» (Russia). *Nat Conserv Res.* 2020;5(3):66-85. doi:10.24189/ncr.2020.032
23. Dymitryszyn I. The effect of the construction and renovation of a highway bypass in Central Poland on the carabid beetle fauna (Coleoptera: Carabidae). *Eur J Entomol.* 2014;111(5):655-62. doi:10.14411/eje.2014.081
24. Kataev GD. The impact of industrial emissions of copper-nickel smelter complex on the status of populations and communities of small mammals in the Kola Peninsula. *Nat Conserv Res.* 2017;2(Suppl.2):19-27. doi:10.24189/ncr.2017.033 [in Russian]
25. Rozhnov VV, Lavrinenko IA, Razzhivin VYu, Makarova OL, Lavrinenko OV, Anufriev VV, et al. Biodiversity revision of a large arctic region as a basis for its monitoring and protection under conditions of active economic development (Nenetsky Autonomous Okrug, Russia). *Nat Conserv Res.* 2019;4(2):1-28. doi:10.24189/ncr.2019.015 [in Russian]
26. Ruchin AB, Alekseev SK, Semishin GB. Seasonal activity dynamics of imago *Carabus coriaceus* Linnaeus, 1758 (Coleoptera, Carabidae) in mixed forests. *Proc Mordovia State Nat Reserve.* 2019;23:239-44. [in Russian]
27. Desender K, Dekoninck W, Dufrêne M, Maes D. Changes in the distribution of carabid beetles in Belgium revisited: have we halted the diversity loss? *Biol Conserv.* 2010;143(6):1549-57. doi:10.1016/j.biocon.2010.03.039
28. Ruchin AB, Egorov LV. Overview of insect species included in the Red Data Book of Russian Federation in the Mordovia State Nature Reserve. *Nat Conserv Res.* 2017;2(Suppl.1):2-9. doi:10.24189/ncr.2017.016 [in Russian].
29. Ruchin AB, Khapugin AA. Red data book invertebrates in a protected area of European Russia. *Acta Zool Acad Sci Hung.* 2019;65(4):349-70. doi:10.17109/AZH.65.4.349.2019
30. Zamotajlov AS, Serdyuk VYu, Khomitskiy EE, Belyi AI. New data on distribution and biology of some rare ground beetles (Coleoptera, Carabidae) in South Russia. *Nat Conserv Res.* 2019;4(4):81-90. doi:10.24189/ncr.2019.066 [in Russian]
31. Egorov LV, Ruchin AB, Semenov VB, Semionenkov OI, Semishin GB. Checklist of the Coleoptera of Mordovia State Nature

- Reserve, Russia. ZooKeys. 2020;962:13-122. doi:10.3897/zookeys.962.54477
32. Nora AHN, Mohamed T, Tahar M, Abderrahmane L. Importance of low perennial plant formations to the sustainability of forest ecosystems in the Saida Mountains (Algeria). *World J Environ Biosci.* 2020;9(2):71-6.
 33. Peet RK. The measurement of species diversity. *Annu Rev Ecol Syst.* 1974;5:285-307. doi:10.1146/annurev.es.05.110174.001441
 34. Kryzhanovskij OL. Family Carabidae – Ground beetles. Determinant of the insects of the European part of the USSR. 1965;2:29-77. [in Russian].
 35. Kryzhanovskij OL. Coleoptera. Beetles of the Suborder Adephaga: The Trachypachidae, Carabidae Families. *Fauna of the USSR*, 1 (2), Leningrad. 1983.
 36. Makarov KV, Kryzhanovskij OL, Belousov IA, Zamotajlov AS, Kabak II, Kataev BM, et al. Systematic list of carabid beetles (Carabidae) of Russia. 2018. Available from: http://www.zin.ru/Animalia/Coleoptera/rus/car_rus.htm [Retrieved on 08.12.2019]. [in Russian]
 37. Kryzhanovskij OL, Belousov IA, Kabak II, Kataev BM, Makarov KV, Shilenkov VG. A checklist of the ground-beetles of Russia and adjacent lands (Insecta, Coleoptera, Carabidae). Sofia – Moscow: PENSOFT Publishers; 1995. 271 p.
 38. Löbl I, Löbl D. (Eds.). *Catalogue of Palaearctic Coleoptera. Revised and updated revision. Vol. 1. Archostemata – Myxophaga – Adephaga.* Lieden-Boston: Brill; 2017. 1443 p. doi:10.1163/9789004330290
 39. Chifundera KZ. Using diversity indices for identifying the priority sites for herpetofauna conservation in the Democratic Republic of the Congo. *Nat Conserv Res.* 2019;4(3):13-33. doi:10.24189/ncr.2019.035
 40. Bulokhova NA. Species composition and domination structure of ground beetles (Coleoptera, Carabidae) in meadow ecosystems of southwestern Russia (Bryansk Region). *Entomolog Oboz.* 1995;74(4):758-63. [in Russian]
 41. Burgio G, Sommaggio D, Marini M, Puppi G, Chiarucci A, Landi S, et al. The influence of vegetation and landscape structural connectivity on butterflies (Lepidoptera: Papilionoidea and Hesperioidea), carabids (Coleoptera: Carabidae), syrphids (Diptera: Syrphidae), and sawflies (Hymenoptera: Symphyta) in Northern Italy Farmland. *Environ Entomol.* 2015;44(5):1299-307. doi:10.1093/ee/nvv105
 42. Jahnová Z, Knapp M, Boháč J, Tulachová M. The role of various meadow margin types in shaping carabid and staphylinid beetle assemblages (Coleoptera: Carabidae, Staphylinidae) in meadow dominated landscapes. *J Insect Conserv.* 2016;20(1):59-69.
 43. Allema B, Hemerik L, Rossing WAH, Groot JCJ, van Lenteren JC, van der Werf W. Dispersal of a carabid beetle in farmland is driven by habitat-specific motility and preference at habitat interfaces. *Entomol Exp Appl.* 2019;167(8):741-54. doi:10.1111/eea.12804
 44. Solodovnikov IA. The ground beetles (Coleoptera, Carabidae) of Belarus Poozeriya with the catalogue of Carabidae of Belarus and adjacent countries. Vitebsk: Vitebsk State University Publisher; 2008. 325 p. [in Russian]
 45. Tselishcheva LG. The spatial structure of populations of ground beetles (Coleoptera, Carabidae) in flood-plain communities of the Nurgush Nature Reserve. In: Potemkin NA. (Ed.): Long-term processes in natural complexes of nature reserves of Russia. Velik Luki: Cent Forest State Nat Res. 2012. P. 239-44. [in Russian]
 46. Chechlowski W. Carabid beetles (Coleoptera, Carabidae) of moist meadow on the Mazovian Lowland. *Memor Zool.* 1989;43:141-67.
 47. Trushitsyna OS, Matalin AV, Makarov KV. Long-term dynamics and spatial distribution of stable and labile components in ground beetle communities (Coleoptera: Carabidae) in a mosaic of flood-plain meadows. *Period Biol.* 2016;118(3):255-72. doi:10.18054/pb.2016.118.3.3928
 48. Twardowski JP, Pastuszko K, Hurej M, Gruss I. Effect of different management practices on ground beetle (Coleoptera: Carabidae) assemblages of uphill grasslands. *Pol J Ecol.* 2017;65(3):400-9.

49. Fan F, Liang C, Tang Y, Harker-Schuch I, Porter JR. Effects and relationships of grazing intensity on multiple ecosystem services in the Inner Mongolian steppe. *Sci Total Environ.* 2019;675:642-50.
50. Lenoir L, Lennartsson T. Effects of timing of grazing on arthropod communities in semi-natural grasslands. *J Insect Sci.* 2010;10(1):60. doi:10.1673/031.010.6001