

The Rate of Crop Residue Decomposition as a Function of the Chemical Composition of Field Crops

Boris Alexandrovich Sotnikov^{1*}, Vladimir Alexandrovich Kravchenko¹, Roman Viktorovich Shchuchka¹

¹ *Bunin Yelets State University, Kommunarov str., 28.1, Yelets, 399770, Russia.*

ABSTRACT

The data on the degree of post-harvest residue decomposition at any given time is needed to determine the reserves of nutrients released from the organic fertilizer applied to the soil and to determine the humification factor of the crop residues being cultivated. Among other factors, this requires an assessment of post-harvest residues in terms of organic matter inputs as well as their chemical composition. To this end, post-harvest residues in the crop rotation were studied in laboratory experiments. Authors have laid a model (laboratory) experience, where duration of the decomposition of crop residues were studied. The data obtained suggest that it is essential to alternate crops with high and slow rates of decomposition because there is a short period after harvesting of predecessors (especially nonfallow) and before sowing winter crops so that often the process of rapid decomposition of post-harvest residues coincides with the growth and development of winter crops.

Keywords: Crop residues, Decomposition, Humification, Crop rotation, Biomass, Controlled conditions.

HOW TO CITE THIS ARTICLE: Sotnikov B A, Kravchenko V A, Shchuchka R V. The Rate of Crop Residue Decomposition as a Function of the Chemical Composition of Field Crops. *Entomol. Appl. Sci. Lett.* 2021;8(2):16-9. <https://doi.org/10.51847/MuZoszyPy0>

Corresponding author: Boris Alexandrovich Sotnikov

E-mail ✉ boris.a.sotnikov@mail.ru

Received: 20/03/2021

Accepted: 22/06/2021

INTRODUCTION

To assess the post-harvest residues of a crop as a potential source of humus matter formation, it is necessary to know the time of their complete decomposition [1, 2]. Such information for some crops of crop rotations, in the conditions of the Central Black Earth Region, is practically absent.

MATERIALS AND METHODS

Therefore, we laid a model (laboratory) experience and studied the duration of crop residues decomposition. Such experiments are the ones where we can get more or less unambiguous answers to certain questions related to post-harvest residue decomposition, as most of the factors influencing residue decomposition were maintained in optimal parameters. These parameters, according to M.M. Kononova [3-5], should be the following – the temperature of about 30°C and soil moisture of 60-80% of the total moisture capacity. A simultaneous increase

in temperature and humidity or their simultaneous reduction decreases the rate of decomposition of crop residues. When one of the factors increases, the intensity of decomposition of organic residues obeys the law of minimum and maximum. If anaerobiosis is constant, the decomposition processes slow down sharply and gradually subside.

RESULTS AND DISCUSSION

A study of residue decomposition under controlled conditions showed (**Table 1**) that the rate of decomposition of non-humified crop residues from the rotation (washed directly in the stationary experiment) was higher than under field conditions. They included residues from previous years.

In the first year, the most rapid reduction in biomass was observed in the winter vetch-oat mixture and maize for silage, both in the occupied and in the green manure fallow. On these variants, half of the biomass applied to soils is

decomposed. This is explained by the fact that post-harvest residues of these crops contained large quantities of easily degradable substances and they (under favorable conditions) decom-

posed quickly. This process was slower in grain crops – winter wheat and barley, which decomposition rate was 1.5-2.0 times lower.

Table 1. The Rate of Decomposition of Crop Residues in the Crop Rotation (Laboratory Experiment in Vessels)

Crop	Mass of residues before setting up the experiment		Decomposed from the beginning			
			1 year		2 year	
	g	%	g	%	g	%
Winter vetch-oat mixture	15	100	<u>7.6</u>	<u>50.7</u>	<u>8.53</u>	<u>56.9</u>
			7.5	49.8	8.35	55.5
Winter wheat	15	100	<u>3.5</u>	<u>23.3</u>	<u>4.07</u>	<u>27.1</u>
			4.1	27.3	4.63	30.8
Maize for silage	15	100	<u>6.3</u>	<u>42.0</u>	<u>9.66</u>	<u>64.4</u>
			7.0	46.7	10.58	70.6
Barley	15	100	<u>5.3</u>	<u>35.3</u>	<u>8.94</u>	<u>59.6</u>
			5.8	38.7	8.92	59.5

Note: above the line — in a crop rotation with occupied fallows, below the line — in a crop rotation with green manure fallows.

A study of crop biomass decomposition only (root and above-ground parts) under controlled conditions showed (**Table 2**) that their decomposition rate was higher than under field conditions.

In the first year, the most rapid reduction in biomass was observed in the winter vetch-oat mixture and maize for silage grown both in the rotation with occupied fallow and in the rotation with green manure fallow. More than half of the

ploughed biomass was decomposed on these variants. This is explained by the fact that post-harvest residues of these crops contained a large amount of easily degradable substances and they (under favorable conditions) decomposed quickly. This process was slower in grain crops – winter wheat and barley – with decomposition rates of 40.1-40.8% and 60.8-61.4%, respectively, which is 1.1-1.5 times higher than mixed residues.

Table 2. The Rate of Decomposition of the Biomass of the Crops in the Crop Rotation (Laboratory Experiment in Vessels)

Crop	Mass of residues before setting up the experiment		Decomposed from the beginning			
			1 year		2 year	
	g	%	g	%	g	%
Winter vetch-oat mixture	15	100	<u>8.3</u>	<u>55.3</u>	<u>10.5</u>	<u>69.9</u>
			8.6	57.3	10.4	69.5
Winter wheat	15	100	<u>5.1</u>	<u>34.0</u>	<u>6.00</u>	<u>40.1</u>
			5.3	35.3	6.10	40.8
Maize for silage	15	100	<u>8.0</u>	<u>53.3</u>	<u>11.2</u>	<u>74.4</u>
			8.3	55.3	11.6	77.6
Barley	15	100	<u>5.9</u>	<u>39.3</u>	<u>9.10</u>	<u>60.8</u>
			6.4	42.7	9.20	61.4

Note: above the line — in a crop rotation with occupied fallows, below the line — in a crop rotation with green manure fallows.

This is one of the important points to consider when planning crop rotations. Crops with high and slow decomposition rates should be alternated, as a short period elapses after the harvesting of the crop's predecessors (especially nonfallow crops) before sowing winter crops so that the rapid decomposition of their post-harvest residues often coincides with the growth and development of winter crops. If the amount of fertilizers applied to winter crops

(especially nitrogen fertilizers) is insufficient, nitrogen starvation of crops may occur, because decomposition of nitrogen-poor post-harvest residues of barley, winter crops, and maize for silage occurs with immobilization of soil mineral nitrogen. Therefore, it is necessary that in the period from harvesting of predecessors to the sowing of winter crops, most of the post-harvest residues have time to decompose and release as many nutrients as possible.

In general, according to the rate of decomposition in the first year in the laboratory experiment, the studied crops can be placed in the following descending order: winter vetch-oat mixture – maize for silage – barley – winter wheat. In the second year, the biomass of the winter vetch-oat mixture and winter wheat was almost completely decomposed – the percentage of decomposition was between 3.5 and 6.7. The decomposition rate of maize and barley residues decreased, leaving 20 and 23%, respectively, undecomposed.

In terms of the rate of biomass decomposition in the second year of the laboratory experiment, the studied crops can be placed in the following descending order: winter vetch-oat mixture – winter wheat – maize for silage – barley.

On average, the biomass of maize for silage decomposed more in two years, slightly less in barley, winter vetch-oat mixture, and winter wheat. The decomposition rate of these crops was 64.4, 59.6, 56.9, and 27.1% respectively against occupied fallow and 70.5, 59.5, 55.7, and 30.9% against green manure fallow.

As to the influence of biomass chemical composition on decomposition rate, crop residues are characterized by a wide variety of chemical composition, which, under other equal conditions, strongly affects the intensity of mineralization-humification processes of organic matter [5-8]. The higher the content of easily mobilized

compounds (water-soluble, simple sugars, starch, and protein) in the decomposing material, the higher the decomposition rate and, conversely, the higher the content of lignin, wax, and resins in the decomposing substrate, the lower the decomposition rate.

In natural phytocenoses, crop residues contain about 2% nitrogen and 3-5% ash elements [9]. The content of nitrogen and ash elements in crops depends on the cultivated crops and fertilizers applied. Perennial leguminous grasses have a similar chemical composition to natural phytocenoses. Poorer in nitrogen and ash elements are post-harvest crop residues of annual legumes and row crops (from 1 to 2%). Less than 1% of nitrogen is contained in crop residues of grain crops.

To identify changes in the chemical composition during the decomposition of crop residues, we compared their nutrient content in different periods. Our research showed (**Table 3**) that the crop residues of the studied crops differed from each other in terms of nutrient content.

The residues of alfalfa, clover, vetch-oat and maize for silage contained the most nitrogen. Grain biomass contained less than 1% of nitrogen.

Crop residues (except winter wheat, where the content of this element was as high as 0.12%) did not differ much in phosphorus content. Its amount varied from 0.24 to 0.28%.

Table 3. The Chemical Composition of Crop Residues

Crop	Nutrient content of crop residues, %				Ratio C: N
	C	N	P	K	
1. Clover	35	1.46	0.28	1.35	24.3
2. Alfalfa	42	1.68	0.24	1.32	25.0
3. Vetch-oat	46	1.23	0.24	2.28	37.0
4. Barley	48	0.58	0.27	1.10	83.0
5. Maize for sowing	45	1.08	0.24	1.15	42.0
6. Winter wheat	48	0.56	0.12	0.91	86.0
7. Winter rye	48	0.50	0.20	0.84	96.0
LSD ₀₅	0.3	0.2	0.05	0.06	-

In terms of potassium content, crop residues can be arranged in the following descending order: vetch-oat, clover, alfalfa, maize for silage, barley, winter wheat, and winter rye.

A widely used indicator for estimating the rate of decomposition of crop residues is their carbon to nitrogen ratio (C: N). It was found that

the rate of decomposition of crop residues reaches its maximum at an initial C: N ratio of 25.

The studies showed that the crop residues of the studied crops differed from each other in terms of carbon and nitrogen content.

It was already higher in alfalfa and clover residues (24.3 and 25.0). Due to this narrow ratio, these residues are capable of rapid mineralization. Immediately after they are ploughed in, a vigorous microbial process begins, resulting in most of the biomass decomposing, which creates favorable conditions for the growth and development of subsequent crops in the crop rotation.

This ratio was wider on the other variants of the experiment (from 37.0 to 96.0). This is because these residues, according to scientists [10, 11], are enriched with lignin, waxes, and resins and contain few protein compounds. Therefore, additional nitrogen mineral fertilizers (10-15 kg a.d.m. per 1 ton) are needed to accelerate the rate of residue decomposition in these variants, while this is not required for clover and alfalfa residue decomposition.

CONCLUSION

Thus, in terms of C: N, and, consequently, higher decomposition rate, crop residues can be placed in the following descending order: clover, alfalfa, vetch-oat, maize for silage, winter rye, winter wheat, and barley. Besides, under the laboratory experiment conditions (controlled conditions), the rate (potential capacities) of decomposition was higher than under field conditions. However, the presence (addition) of residues from previous years reduced the rate of decomposition.

ACKNOWLEDGMENTS: None

CONFLICT OF INTEREST: None

FINANCIAL SUPPORT: None

ETHICS STATEMENT: Authors are aware of, and comply with, best practice in publication ethics specifically with regard to authorship and compliance with policies on research ethics.

REFERENCES

1. Benzineb E, Kambouche N, Hamiani A, Belahouel S, Zitouni H, Toumi H. Phenolics Compounds and Biological Activity of Leaves of *Anabasis Articulata*, an Algerian Medicinal Plant. *Int J Pharm Res Allied Sci.* 2019;8(4):1-5.

2. Kanjekar AP. On Anti-Diabetic Potential of Phyto-nanoparticles Comparison with Hormonal Therapy and Medicinal Plants. *Int J Pharm Phytopharmacol Res.* 2019;9(1):103-11.
3. Kononova MM. Issues of soil humus and modern tasks of its study. Moscow: Izd-vo AN SSSR; 1951. 391 p.
4. Kononova MM. Humus of the main types of soils of the USSR, its nature and ways of formation. *Soil Sci.* 1956;(3):18-30.
5. Kononova MM. Organic matter, Moscow: Izd-vo AN SSSR; 1963. 313 p.
6. Mi W, Sun Y, Gao Q, Liu M, Wu L. Changes in humus carbon fractions in paddy soil given different organic amendments and mineral fertilizers. *Soil Tillage Res.* 2019 Dec 1;195:104421. DOI: 10.1016/j.still.2019.104421
7. Wu M, Wei S, Liu J, Liu M, Jiang C, Li Z. Long-term mineral fertilization in paddy soil alters the chemical structures and decreases the fungistatic activities of humic acids. *Eur J Soil Sci.* 2019;70(4):776-85.
8. Zhang T, He X, Li M, Liu Y, Xi B, Zhang H, et al. Spectral characteristics of humus in chernozem in Qinghai-Tibet Plateau under reclamation and long-term fertilization. *Acta Pedologica Sinica.* 2019;56(2):398-407.
9. Hafez M, Rashad M, Popov AI. The biological correction of agro-photosynthesis of soil plant productivity. *J Plant Nutr.* 2020;43(19):2929-80.
10. Dedov AV, Korzhov SI. The transformation of various mixtures of organic substances. Proceedings of scientific-practical International Conference. Voronezh: VGU; 1995. 40 p.
11. Mishustin EN. The use of hay as a fertilizer. *Agrochemistry.* 1977;8:49-54.