

## Study of Butterfly Diversity in Chilkigarh, West Bengal (India)

Sourav Karan<sup>1</sup>, Rakesh Acharya<sup>1</sup>, Koushik Sen<sup>1</sup>, Sanjib Kumar Das<sup>1\*</sup>

<sup>1</sup>Department of Zoology, Jhargram Raj College, Jhargram, W.B-721507, India.

### ABSTRACT

An attempt has been made to understand the butterfly community of Chilkigarh, a village with heritage site and one of the famous tourist destinations in Jhargram subdivision, West Bengal, India. These winged jewels were studied adopting conventional sampling techniques followed by measurement of different diversity indices. A total of 59 species belonging to 6 families and 14 subfamilies have been documented between December 2021 and November 2022, with a good number of species to genus ratio (1.31:1). Among 59, 11 are protected under different schedules of the Indian Wildlife Protection Act, 1972. Nymphalids with the highest percentage (42.3%) secured the dominant status among the families. Relative abundance study reveals 10 species with subdominant status but no one with dominant status. Analysis of different diversity indices indicates that Chilkigarh carries rich butterfly fauna. Information from this preliminary study may provide a direction for future investigations, such as the identification of new species, host plants, nectar plants, and seasonal fluctuations over time.

**Keywords:** Butterfly diversity, Lepidoptera, Pollinator, Diversity indices, Chilkigarh, Jhargram.

**HOW TO CITE THIS ARTICLE:** Karan S, Acharya R, Sen K, Das SK. Study of Butterfly Diversity in Chilkigarh, West Bengal (India). Entomol Appl Sci Lett. 2023;10(4):26-34. <https://doi.org/10.51847/DczmEKgc9h>

**Corresponding author:** Sanjib Kumar Das

**E-mail** ✉ [sanjib.biology2012@gmail.com](mailto:sanjib.biology2012@gmail.com)

**Received:** 18/09/2023

**Accepted:** 07/12/2023

### INTRODUCTION

Faunal components of forest ecosystems play a crucial role in the maintenance and sustainability of that ecosystem, and ecological indicator species are used worldwide for assessing biodiversity. The effect of forest management on the structure and function of a forest ecosystem can be illustrated by studying bioindicators [1, 2]. Butterflies, among the insect groups, due to their short life history traits, host plant preferences, easily identifiable features (unique wing color patterns), high diversity, sensitivity to microclimate as well as environmental changes, themselves achieve such a status to be accepted as bioindicator [3]. Butterflies are primary consumers in the forest ecosystem and play an important role as herbivores in the stability of food webs [4, 5]. To maintain and improve community structure, they serve as pollinators [5-7], surrogate species for floral and faunal diversity [8], host of parasitoids [5, 9], and prey of predators [4, 5, 10]. Chilkigarh, a rural and


tribal area situated on the bank of Dulung River, mostly surrounded by Sal Forest, has become a famous tourist destination for the presence of Chilkigarh Raj Palace and Kanak Durga Sacred Grove. In 2018, it received the status of Chilkigarh Kanak Durga Biodiversity Heritage Site from the Environment Department, Govt. of West Bengal, India. Altogether, 388 plant species, including 105 with medicinal values and 26 species of megafauna (vertebrates), have been reported [11]. Recently, 37 species of birds have been documented from different sites in Chilkigarh [12]. With the aim to elucidate the butterfly community of this area, a study has been done to gather knowledge about their diversity, dominant family group, species-to-genus ratio, legally protected species, etc., that will surely provide a route for future investigations regarding their nectar plants, host plants, seasonal fluctuations and ultimately planning for conservation.

### MATERIALS AND METHODS

**Study area**

Chilkigarh (**Figure 1**) is a tribal area (latitudes 22° 27' 20" N and 22° 56' 50" N and longitudes 86° 52' 20" E and 86° 53' 10" E), having an average elevation of 60-85 m above the sea level [13], situated in Jamboni CD block, under Jhargram Subdivision of District Jhargram, West Bengal. The western boundary falls under the lower ranges of the Chhotanagpur Plateau, the northwest area is uninhabitable, and most of the areas are non-productive due to the nature of the lateritic soil. Dulung, a monsoon-nourished river, passes down across the village. Chilkigarh forest

is situated on the eastern side of the bank of this river, comprising heterogeneous vegetation of semideciduous, deciduous, and evergreen trees [11]. This area can be classified under the category of "Tropical Moist Deciduous Forest" as huge Sal plants predominate. Different types of shrubs, herbs, climbers, and grasses provide resources to small to large creatures. Dulung river bank, Chilkigarh Kanak Durga Sacred Grove, Open grasslands, Chilkigarh Raj Palace, Sonajhuri garden, Sal forests, and Agricultural lands were selected for study.



**Figure 1.** Location map of Chilkigarh in West Bengal, India [12]

**Data collection**

The study was carried out from 7:00 am to 10:00 am and 3:00 pm to 6:00 pm for 12 months (December 2021 to November 2022), adopting sampling techniques such as Pollard walk method [14], Direct searching method [15] & Time Constrained method [16].

**Identification & documentation of butterfly species**

Photographic documentation was done by visiting different sites once or twice per month to capture photos from the best possible angles using a Canon IXUS 190 Digital camera and mobile phone camera – Redmi 6 Pro & Redmi Note 8. Species were identified using the following references [17-

19] and further consulting the website of Butterflies of India [20].

#### Community analysis

To understand the structure of butterfly community  $\alpha$ -diversity, i.e., the diversity of species within the community has been measured using the following diversity indices. All data were calculated using MS Excel 2019 software, and results were further verified using statistical software PAST version 4.03 [21].

#### Species richness

The Shannon-Wiener index, commonly known as the Shannon index of diversity [22], sometimes erroneously called the Shannon-Weaver index, was derived independently by Shannon and Wiener, which apply information theory to measure species diversity. Rare species with very few individuals can contribute some value to this index [23]. It is calculated considering Eq. 1 as follows:

$$H' = - \sum pi \ln pi \quad (1)$$

Where  $H'$  is the value of the Shannon index and  $pi$  is the proportion of individuals of  $i$ th species in the community. The value usually ranges between 1.5 to 3.5 and rarely exceeds 4.5. The value of  $H'$  is related to species richness but is also influenced by underlying species abundance distribution. Margalef's index [24] is used to calculate species richness considering Eq. 2 as follows:

$$I_{Mg} = S - 1 / \ln N \quad (2)$$

Where  $S$  is the total number of species and  $N$  is the total number of individuals in  $S$  species.

#### Species abundance

Simpson's index [25] is the measure of the probability that two organisms picked at random from a community will belong to the same species. This index relates the contribution made by each species to the total number of individuals present. It can be calculated considering Eq. 3 as follows:

$$D = \sum_{i=1}^S (pi)^2 \quad (3)$$

The value of  $D$  ranges between 0 to 1 and is inversely proportional to the wealth of species. As the value of Simpson's index increases, the species diversity decreases. Therefore, the more the index value is inclined to 0, the more abundance will be in the community.

Simpson's index of diversity = 1 - probability of picking two organisms that are the same species and calculated considering following Eq. 4 as follows:

$$D = 1 - \sum_{i=1}^S (pi)^2 \quad (4)$$

Where  $D$  is the value of Simpson's index of diversity and  $pi$  is the proportion of individuals of the species in the community. Simpson's index of diversity gives relatively little weightage to rare species and more weightage to common species. It ranges from 0 (low diversity) to a maximum of  $(1-1/S)$ , where  $S$  is the total number of species.

**Table 1.** Family-wise checklist with common and scientific names along with relative abundance, dominant status, and WPA status of each butterfly species encountered at Chilkigarh

| Sl. No.   | Common Name   | Scientific name                          | Abundance | Relative abundance (%) | Dominant status * | WPA Schedule status |
|---|---------------|--|-----------|------------------------|-------------------|---------------------|
| <b>Family: Nymphalidae (Brush-footed Butterflies)</b>                             |               |  |           |                        |                   |                     |
| <b>Subfamily: Biblidinae (Castors &amp; Jokers)</b>                               |               |  |           |                        |                   |                     |
| 1   | Angled Castor | <i>Ariadne ariadne</i> (Linnaeus,1763)   | 79        | 4.990                  | SD                |                     |
| 2   | Common Castor | <i>Ariadne merione</i> (Cramer,1777)     | 16        | 1.010                  | SR                |                     |
| <b>Subfamily: Danainae (Milkweed Butterflies)</b>                                 |               |  |           |                        |                   |                     |
| 3   | Blue Tiger    | <i>Tirumala limniace</i> (Cramer,1775)   | 28        | 1.768                  | R                 |                     |
| 4   | Common Crow   | <i>Euploea core</i> (Cramer,1780)        | 93        | 5.874                  | SD                | IV                  |
| 5   | Plain Tiger   | <i>Danaus chrysippus</i> (Linnaeus,1758) | 23        | 1.452                  | R                 |                     |
| 6   | Striped Tiger | <i>Danaus genutia</i> (Cramer,1779)      | 14        | 0.884                  | SR                | I                   |
| <b>Subfamily: Heliconiinae (Costers, Lacewings, Fritillaries &amp; Relatives)</b> |               |  |           |                        |                   |                     |

|  |                          |  |    |       |    |    |
|--|--------------------------|--|----|-------|----|----|
| 7  | Common Leopard           | <i>Phalanta phalantha</i> (Drury,1773)             | 20 | 1.263 | R  |    |
| 8  | Tawny Coster             | <i>Acraea violae</i> (Fabricius, 1775)             | 58 | 3.663 | SD |    |
| <b>Subfamily: Limenitinae (Barons, Sailors, Sergeants &amp; Relatives)</b> |                          |  |    |       |    |    |
| 9  | Baronet                  | <i>Euthalia nais</i> (Forster,1771)                | 21 | 1.326 | R  |    |
| 10   | Commander                | <i>Moduza procris</i> (Cramer,1777)                | 15 | 0.947 | SR |    |
| 11   | Common Baron             | <i>Euthalia aconthea</i> (Cramer,1777)             | 18 | 1.137 | R  | II |
| 12   | Grey Count               | <i>Tanaecia lepidea</i> (Butler,1868)              | 2  | 0.126 | SR | II |
| 13   | Chestnut-streaked Sailer | <i>Neptis jumbah</i> (Moore, 1857)                 | 34 | 2.147 | R  |    |
| <b>Subfamily: Nymphalinae (Pansies, Eggflies &amp; Relatives)</b>          |                          |  |    |       |    |    |
| 14   | Great Eggfly             | <i>Hypolimnas bolina</i> (Linnaeus,1758)           | 44 | 2.779 | R  |    |
| 15   | Blue Pansy               | <i>Junonia orithya</i> (Linnaeus,1758)             | 37 | 2.337 | R  |    |
| 16   | Chocolate Pansy          | <i>Junonia iphita</i> (Cramer,1779)                | 46 | 2.905 | R  |    |
| 17   | Grey Pansy               | <i>Junonia atlites</i> (Linnaeus,1763)             | 49 | 3.095 | R  |    |
| 18   | Lemon Pansy              | <i>Junonia lemonias</i> (Linnaeus,1758)            | 25 | 1.579 | R  |    |
| 19   | Peacock Pansy            | <i>Junonia almana</i> (Linnaeus,1758)              | 42 | 2.653 | R  |    |
| 20   | Yellow Pansy             | <i>Junonia hierta</i> (Fabricius,1798)             | 17 | 1.073 | SR |    |
| <b>Subfamily: Satyrinae (Browns)</b>                                       |                          |  |    |       |    |    |
| 21   | Bamboo Treebrown         | <i>Lethe europa</i> (Fabricius,1775)               | 4  | 0.252 | SR |    |
| 22   | Common Evening Brown     | <i>Melanitis leda</i> (Linnaeus,1758)              | 56 | 3.537 | SD |    |
| 23   | Common Four-ring         | <i>Ypthima huebneri</i> (Kirby,1871)               | 18 | 1.137 | R  |    |
| 24   | Common Palmfly           | <i>Elymnias hypermnestra</i> (Linnaeus,1763)       | 23 | 1.452 | R  |    |
| 25   | Common Bushbrown         | <i>Mycalesis perseus</i> (Fabricius,1775)          | 19 | 1.200 | R  |    |
| <b>Family: Lycaenidae (Blues &amp; Hairstreaks)</b>                        |                          |  |    |       |    |    |
| <b>Subfamily: Polymmatinae (Weak Blues)</b>                                |                          |  |    |       |    |    |
| 26   | Common Pierrot           | <i>Castalius rosimon</i> (Fabricius,1775)          | 52 | 3.284 | SD | I  |
| 27   | Dark Grass Blue          | <i>Zizeeria karsandra</i> (Moore,1865)             | 10 | 0.631 | SR |    |
| 28   | Forget-me-not            | <i>Catochrysops strabo</i> (Fabricius,1793)        | 12 | 0.758 | SR |    |
| 29   | Lesser Grass Blue        | <i>Zizina otis</i> (Fabricius,1787)                | 21 | 1.326 | R  |    |
| 30   | Lime Blue                | <i>Chilades lajus</i> (Stoll,1780)                 | 8  | 0.505 | SR | I  |
| 31   | Pointed Ciliate Blue     | <i>Anthene lycaenina</i> (C & R. Felder,1868)      | 22 | 1.389 | R  | II |
| 32   | Quaker                   | <i>Neopithecops zalmora</i> (Butler,1870)          | 4  | 0.252 | SR |    |
| <b>Subfamily: Theclinae (Strong Blues, Hairstreaks)</b>                    |                          |  |    |       |    |    |
| 33   | Common Guava Blue        | <i>Deudorix isocrates</i> (Fabricius,1793)         | 7  | 0.442 | SR | II |
| 34   | Common Silverline        | <i>Spindasis vulcanus</i> (Fabricius,1775)         | 5  | 0.315 | SR |    |
| 35   | Peacock Royal            | <i>Tajuria cippus</i> (Fabricius,1798)             | 4  | 0.252 | SR | II |
| 36   | Bengal Slate Flash       | <i>Rapala manea</i> (Hewitson,1863)                | 4  | 0.252 | SR |    |
| 37   | Indian Oakblue           | <i>Arhopala atrax</i> (Hewitson,1862)              | 2  | 0.126 | SR |    |
| <b>Family: Papilionidae (Swallowtails)</b>                                 |                          |  |    |       |    |    |
| <b>Subfamily: Papilioninae</b>   |                          |  |    |       |    |    |
| 38   | Lime Swallowtail         | <i>Papilio demoleus</i> (Linnaeus,1758)            | 52 | 3.284 | SD |    |
| 39   | Common Mormon            | <i>Papilio polytes</i> (Linnaeus,1758)             | 79 | 4.990 | SD |    |
| 40   | Common Banded Peacock    | <i>Papilio crino</i> (Fabricius,1793)              | 3  | 0.189 | SR |    |
| 41   | Blue Mormon              | <i>Papilio polymnestor</i> (Cramer,1775)           | 6  | 0.379 | SR |    |
| 42   | Common Mime              | <i>Chilasa clytia</i> (Linnaeus,1758)              | 15 | 0.947 | SR | I  |
| 43   | Common Jay               | <i>Graphium doson</i> (C&R Felder,1864)            | 13 | 0.821 | SR |    |
| 44   | Tailed Jay               | <i>Graphium agamemon</i> (Linnaeus,1758)           | 49 | 3.095 | R  |    |
| 45   | Spot Swordtail           | <i>Graphium nomius</i> (Esper,1799)                | 1  | 0.063 | SR |    |
| 46   | Common Rose              | <i>Atrophaneura aristolochiae</i> (Fabricius,1775) | 45 | 2.842 | R  |    |
| <b>Family: Pieridae (Whites &amp; Yellows)</b>                             |                          |  |    |       |    |    |

| Subfamily: Pierinae (Whites)                                       |                           |  |    |       |    |    |
|--|---------------------------|--|----|-------|----|----|
| 47   | Common Gull               | <i>Cepora nerissa</i> (Fabricius,1775)           | 15 | 0.947 | SR | II |
| 48   | Common Jezebel            | <i>Delias eucharis</i> (Drury,1773)              | 19 | 1.200 | R  |    |
| 49   | Indian Common Wanderer    | <i>Pareronia hippia</i> (Fabricius,1787)         | 62 | 3.916 | SD |    |
| 50   | Eastern Striped Albatross | <i>Appias olferna</i> (Swinhoe,1890)             | 2  | 0.126 | SR |    |
| 51   | Psyche                    | <i>Leptosia nina</i> (Fabricius,1793)            | 83 | 5.243 | SD |    |
| Subfamily: Coliadinae (Yellows)                                    |                           |  |    |       |    |    |
| 52   | Mottled Emigrant          | <i>Catopsilia pyranthe</i> (Linnaeus,1758)       | 54 | 3.411 | SD |    |
| 53   | Oriental Lemon Emigrant   | <i>Catopsilia pomona</i> (Fabricius,1775)        | 6  | 0.379 | SR |    |
| 54   | Three-spot Grass Yellow   | <i>Eurema blanda</i> (Boisduval,1836)            | 23 | 1.452 | R  |    |
| Family: Hesperidae (Skippers)                                      |                           |  |    |       |    |    |
| Subfamily: Pyrginae (Flats & Angles)                               |                           |  |    |       |    |    |
| 55   | Common Snow Flat          | <i>Tagiades japedus</i> (Stoll,1781)             | 5  | 0.315 | SR |    |
| Subfamily: Hesperinae (Bobs, Hoppers, Redeyes, Swifts & Relatives) |                           |  |    |       |    |    |
| 56   | Dark Palm Dart            | <i>Telicota ancilla</i> (Herrich-Schaffer, 1869) | 39 | 2.463 | R  |    |
| 57   | Rice Swift                | <i>Borbo cinnara</i> (Wallace,1866)              | 35 | 2.210 | R  |    |
| 58   | Common Red Eye            | <i>Matapa aria</i> (Moore,1865)                  | 18 | 1.137 | R  |    |
| Family: Riodinidae (Metalmarks)                                    |                           |  |    |       |    |    |
| Subfamily: Riodininae  |                           |  |    |       |    |    |
| 59   | Double-banded             | <i>Abisara bifasciata</i> (Moore,                | 7  | 0.442 | SR |    |

\*RA<1=Subprecedent (SR); 1.1-3.1=Recedent (R); 3.2-10=Subdominant (SD); 10.1-31.6=Dominant (D) and >31.7%=Eudominant

### Species evenness

Pielou's index [26] was used to measure species evenness. It was calculated considering Eq. 5 as follows:

$$E = H' / \ln S \quad (5)$$

Where  $H'$  is the Shannon index and  $S$  is the total number of species. Value of  $e$  ranges from 0 to 1. More the index value inclined towards 1, the more will be the evenness in the community. Dominance status of each species was enumerated on the basis of relative abundance following Engelmann's scale [27]. Rank-abundance curve (Whittaker plot) is prepared, taking abundance rank on the X axis against relative abundance on the Y axis to graphically represent the relative species abundance [28].

## RESULTS AND DISCUSSION

In our study, overall, 59 species of butterflies were recorded with a total count of 1583 individuals belonging to 45 genera under 6 families from Chilkiharh (Table 1; Figures 3 and 4). The family Nymphalidae appeared to be the most dominant family (42.3% with 25 species), followed by Lycaenidae (20.3% with 12 species), Papilionidae (15.2% with 9 species), Pieridae

(13.5% with 8 species), Hesperidae (6.7% with 4 species) and Riodinidae (1.6% with 1 species) (Figure 2). Previous reports also support our findings that Nymphalidae is the dominant family in the neighboring districts: Purulia [29], Haldia [30], Midnapore [23, 30], and Howrah [31].





Figure 2. Percentage composition of Butterfly families.

The ratio of species to genus is 1.31: 1. The proportion of butterflies under six families from genera to species is represented in Figure 5.

11 species were found legally protected under different Schedules of the Wildlife (Protection) Act, 1972 [32], but none were found globally threatened as per the IUCN Red List (Ver 3.1) [33]. Of these legally protected species Striped Tiger (*D. genutia*), Common Pierrot (*C. rosimon*),

Lime Blue (*C. lajus*), Common Mime (*C. Aclytia*) are protected under Schedule I. Common Baron (*E. aconthea*), Grey Count (*T. lepidea*), Pointed Ciliate Blue (*A. lycaenina*), Common Guava Blue (*V. isocrates*), Peacock Royal (*T. cippus*) and Common Gull (*C. nerissa*) are protected under Schedule II and Common Crow (*E. core*) is protected under Schedule IV. In the family, Nymphalidae *Euploea core* was found to be the most abundant species, while *Tanaecia lepidea* was the least one. Under the family Lycaenidae, *Castalius rosimon* was more common, while *Arhopala atrax* was the least common. Similarly, in papilionidae, *Papilio polytes* was well


encountered compared to only a single species of *Graphium nomius*. In Pieridae, *Leptosia nina* was counted more than *Appias olferna*. *Abisara bifasciata* is the only species recorded under the family Riodinidae. An analysis of relative abundance following Engelmann's scale [26] reveals the absence of dominant species in Chilkigarh but 10 species viz. *Ariadne ariadne*, *Euploea core*, *Acraea terpsicore*, *Melanitis leda*, *Castalius rosimon*, *Papilio demoleus*, *Papilio polytes*, *Pareronia hippia*, *Leptosia nina* and *Catopsilia pyranthe* were subdominant in nature (Table 1).



**Figure 3.** 1. Angled Castor 2. Common Castor 3. Common Crow 4. Blue Tiger 5. Plain Tiger 6. Striped Tiger 7. Common Leopard 8. Tawny Coster 9. Baronet 10. Commander 11. Common Baron 12. Grey Count 13. Chestnut-streaked Sailer 14. Great Eggfly (male) 15. Great Eggfly (female) 16. Gray Pansy 17. Yellow Pansy (male) 18. Blue Pansy (male) 19. Lemon Pansy 20. Peacock Pansy 21. Chocolate Pansy 22. Bamboo Tree Brown 23. Common Evening Brown 24. Common Four-ring 25. Common Palmfly (male) 26. Common Bush Brown 27. Dark Grass Blue 28. Lesser Grass Blue 29. Lime Blue 30. & 31. Common Pierrot 32. Forget-me-not 33. Pointed Ciliate Blue 34. Quaker 35. Common Guava Blue.




**Figure 4.** 36. Common Silverline 37. Peacock Royal 38. Indian Oak Blue 39. Bengal Slate Flash 40. Common Jay 41. Tailed Jay 42. Spot Swordtail 43. Blue Mormon 44. Common Mormon (male) 45. Common Mormon (female) 46. Common Rose 47. & 48. Lime Butterfly 49. Common Mime (male) 50. Common Mime (female) 51. Common Banded Peacock 52. Indian Common Wanderer 53. Common Gull 54. Eastern Striped Albatross 55. Common Jezebel 56. Psyche 57. Three Spot Grass Yellow 58. Mottled Emigrant 59. Oriental Lemon Emigrant 60. Double Banded Judy (male) 61. Double Banded Judy (female) 62. Common Red Eye 63. Dark Palm Dart 64. Rice Swift 65. Common Snow Flat.



**Figure 5.** Genus to species proportion of butterflies under six families.

The calculated values of Shannon index (1) and Margalef's index (2) are 3.73 and 7.87, respectively, indicating that the butterfly community of Chilkgarh has high species richness which is consistent with the other findings [23, 30, 34]. The calculated value of Simpson's index (3) is 0.029. As the value is more inclined towards 0, it suggests a high proportion of species abundance. The value of Simpson's index of diversity (4) is 0.9708, suggesting the studied butterfly community is a diversified one. The species evenness (5) for the studied community is  $E=0.9148$ , which indicates high evenness, as it is more inclined to 1.

Given that the abundances of the high-ranking and low-ranking species are very different, the rank abundance curve for the community exhibits strong evenness with a comparatively low steep inclination in the Whittaker plot. High evenness between the various species is conditioned by a modest gradient (**Figure 6**).



**Figure 6.** Whittaker plot of rank-abundance of butterfly community of Chilkiagarh.

### CONCLUSION

This preliminary investigation suggests that the Chilkiagarh area has rich butterfly diversity. Identification of different host plants, nectaring plants, studying seasonal variation, searching for new species, measuring different environmental parameters that affect their life cycle, and correlating all these together in the future will surely help us to predict the complete picture of butterfly community in this area. Surveys at regular intervals will make us aware of any anthropogenic impact due to tourism. If any, accordingly, conservation strategies can be planned to restore these beautiful creatures.

**ACKNOWLEDGMENTS:** The authors are grateful to Dr. Rahul Kumar Datta (Associate Professor & Head, Department of Zoology, Jhargram Raj College) for his valuable suggestions during this work.

**CONFLICT OF INTEREST:** None

**FINANCIAL SUPPORT:** None

**ETHICS STATEMENT:** In this study none of the butterfly species were captured or harmed by any means. Images in the figure are the result of live photography in their natural habitat.

### REFERENCES

- Pearce JL, Venier LA. The use of ground beetles (Coleoptera: Carabidae) and spiders (Araneae) as bioindicators of sustainable forest management: a review. *Ecol Indic.* 2006;6(4):780-93.
- Maleque MA, Maeto K, Ishii HT. Arthropods as bioindicators of sustainable forest management, with a focus on plantation forests. *Appl Entomol Zool.* 2009;44(1):1-1.
- Lee CM, Kim SS, Kwon TS. Butterfly fauna in Mount Gariwang-san, Korea. *J Asia-Pac Biodivers.* 2016;9(2):198-204. doi:10.1016/j.japb.2016.02.005
- Rusman R, Atmowidi T, Peggie D. Butterflies (Lepidoptera: Papilionoidea) of Mount Sago, West Sumatra: Diversity and Flower Preference. *Hayati J Biosci.* 2016;23(3):132-7. doi:10.1016/j.hjb.2016.12.001
- Chowdhury S, Chowdhury SK. A study on butterfly diversity and related host plants in Joychandi Hill of Purulia district, West Bengal, India. *IOSR J Environ Sci Toxicol Food Technol.* 2020;14(10):55-60. doi:10.1093/aesa/91.3.323
- Atmowidi T, Buchori D, Manuwoto S, Suryobroto B, Hidayat P. Diversity of pollinator insects in relation to seed set of mustard (*Brassica rapa* L.: Cruciferae). *HAYATI J Biosci.* 2007;14(4):155-61. doi:10.4308/hjb.14.4.155
- Mukherjee S, Banerjee S, Saha GK, Basu P, Aditya G. Butterfly diversity in Kolkata, India: An appraisal for conservation management. *J Asia-Pac Biodivers.* 2015;8(3):210-21. doi:10.1016/j.japb.2015.08.001
- Ehrlich PR, Raven PH. Butterflies and plants: a study in coevolution. *Evolution.* 1964;586-608.
- Van Nouhuys S, Hanski I. Colonization rates and distances of a host butterfly and two specific parasitoids in a fragmented landscape. *J Anim Ecol.* 2002;71:639-50.
- Hammond PC, Miller JC. Comparison of the biodiversity of Lepidoptera within three forested ecosystems. *Ann Entomol Soc Am.* 1998;91(3):323-8.
- Bhakat RK. Biodiversity Conservation Through A Sacred Grove. *Indian J Biol Sci.* 2015;21:59-62.



12. Das SK, Karan S, Sen K. Biodiversity of Avifauna in Chilkigarh, Jhargram, West Bengal, India. *World J Environ Biosci.* 2022;11(3):8-13. doi:10.51847/jNtkP7dkxS
13. Saadi SM, Mondal I, Sarkar S, Mondal AK. Medicinal plants diversity modelling using remote sensing & GIS technology of Chilkigarh, West Bengal, India. *Trop Plant Res.* 2020;7(2):440-51.
14. Pollard E. A method for assessing changes in the abundance of butterflies. *Biol Conserv.* 1977;12(2):115-34. doi:10.1016/0006-3207(77)90065-9
15. Sutherland WJ, editor. *Ecological census techniques: a handbook.* Cambridge university press; 2006.
16. Suman A, Ravikanthachari N, Kunte K. A comparison between time-constrained counts and line transects as methods to estimate butterfly diversity in tropical habitats. *BioRxiv.* 2021:2021-09. doi:10.1101/2021.09.04.458959
17. Evans WH. *The Identification of Indian Butterflies.* 2nd ed. Bombay Natural History Society. 1932.
18. Wynter-Blyth MA. *Butterflies of the Indian region.* Bombay Natural History Society. 1957.
19. Kehimkar I. *The Book of Indian Butterflies.* Bombay Natural History Society. Oxford University Press. Walton Street, Oxford, New York; 2008.
20. Kunte K, Sondhi S, Roy P. (Chief Editors). *Butterflies of India, v. 3.03.* Indian Foundation for Butterflies and National Centre for Biological Sciences. Available from: <https://www.ifoundbutterflies.org/>
21. Hammer Ø, Harper DA. Past: paleontological statistics software package for education and data analysis. *Palaeontol Electron.* 2001;4(1):1-9.
22. Shannon CE, Weaver W. *The Mathematical Theory of Communication.* The University of Illinois Press, Urbana and Chicago; 1963. 117pp.
23. Biswas SJ, Patra D, Roy S, Giri SK, Pal S, Hossain A. Butterfly diversity throughout Midnapore urban area in West Bengal, India. *J Threat Taxa.* 2019;11(14):14816-26. doi:10.11609/jott.4587.11.14.14816-14826
24. Margalef R. Temporal succession and spatial heterogeneity in phytoplankton. In: *Perspectives in marine biology, Buzzati-Traverso (ed.), Univ. Calif. Press, Berkeley; 1958. pp.323-47.*
25. Simpson GG. Species density of North American recent mammals. *Syst Zool.* 1964;13(2):57-73.
26. Pielou EC. *An Introduction to Mathematical Ecology,* John Wiley, New York, NY, USA; 1969. viii+286pp.
27. Engelmann HD. Zur Dominanzklassifizierung von Bodenarthropoden. *Pedobiologia.* 1978;18:378-80.
28. Whittaker RH. Dominance and Diversity in Land Plant Communities: Numerical relations of species express the importance of competition in community function and evolution. *Science.* 1965;147(3655):250-60. doi:10.1126/science.147.3655.250
29. Samanta S, Das D, Mandal S. Butterfly fauna of Baghmundi, Purulia, West Bengal, India: a preliminary checklist. *J Threat Taxa.* 2017;9(5):10198-207. doi:10.11609/jott.2841.9.5.10198-10207
30. Pahari PR, Mishra NP, Sahoo A, Bhattacharya T. A study on the butterfly diversity of Haldia industrial belt and adjacent rural area in Purba Medinipur district, West Bengal, India. *World Sci News.* 2018(97):207-24.
31. Dwari S, Mondal AK, Chowdhury S. Diversity of butterflies (Lepidoptera: Rhopalocera) of Howrah district, West Bengal, India. *J Entomol Zool Stud.* 2017;5(6):815-28.
32. The Wildlife (Protection) Act, 1972, with the Wildlife (Protection) Amendment Act, 2002. *The Gazette of India,* 148pp. Available from: [https://web.archive.org/web/20201130145631/http://legislative.gov.in/sites/default/files/A1972-53\\_0.pdf](https://web.archive.org/web/20201130145631/http://legislative.gov.in/sites/default/files/A1972-53_0.pdf)
33. IUCN. *The IUCN Red List of Threatened Species.* Version 2022-1. 2022. Available from: <https://www.iucnredlist.org>
34. Mahata A, Mishra NP, Palita SK. Butterflies (Lepidoptera: Rhopalocera) of the undivided Midnapore district, West Bengal, India: a preliminary report. *J Threat Taxa.* 2020;12(17):17347-60. doi:10.11609/jott.5142.12.17.17347-17360