



Influence of Non-Hormonal Drugs on Hormonal Background and Biochemical Parameters of Blood

Diana Alievna Ashibokova¹, Khady Aslanovna Kostoeva², Galina Vladimirovna Osipchuk³, Sergey Nikolaevich Povetkin^{4*}, Alexey Viktorovich Kozlikin⁵, Alexander Nikolaevich Simonov⁶

¹Department of Therapy, Faculty of Medicine of Stavropol State Medical University, Stavropol, Russia.

²Department of Therapy, Faculty of Medicine of Yaroslavl State Medical University, Yaroslavl, Russia.

³Laboratory of Biotechnologies in Reproduction and Embryo Transplantation, Scientific-Practical Institute of Biotechnologies in Animal Science and Veterinary Medicine, Maksimovka Village, Moldova.

⁴Laboratory of Food and Industrial Biotechnology, Faculty of Food Engineering and Biotechnology, North Caucasus Federal University, Stavropol, Russia.

⁵Department of Food Technologies, Faculty of Biotechnology, Don State Agrarian University, Persianovsky, Russia.

⁶Basic Department of Epizootiology and Microbiology, Faculty of Veterinary Medicine of Stavropol State Agrarian University, Stavropol, Russia.

ABSTRACT

Any impact on the body and the physiological processes occurring in the body affect, to varying degrees, the composition of the blood. When making a diagnosis and selecting therapeutic agents, it is especially important to timely and correct conduct biochemical and hematological studies of blood parameters. When prescribing various drugs, including protein-mineral complexes and micro- and macroelements, it is especially important to understand what effect this or that drug will have on the body as a whole. In this scientific article, the influence of a non-hormonal complex, designed to correct the hormonal background of the body, was studied using the example of sheep. Animals of the experimental group daily, for 50 days, were given 5 ml of a preparation containing iodine with amylopectin and were injected with 1.5 ml of E-selenium and 0.5 ml of a tissue preparation mixed with 1 ml of a 0.5% novocaine solution. At the end of the experiment, the dynamics of protein and carbohydrate-lipid metabolism in laboratory animals were studied. It has been established that the non-hormonal agents used do not hurt the body and contribute to a more optimal recovery of the body.

Keywords: Non-hormonal agents, Hormonal correction, Protein metabolism indicators, Carbohydrate-lipid metabolism indicators.

HOW TO CITE THIS ARTICLE: Ashibokova DA, Kostoeva KhA, Osipchuk GV, Povetkin SN, Kozlikin AV, Simonov AN. Influence of Non-Hormonal Drugs on Hormonal Background and Biochemical Parameters of Blood. Entomol Appl Sci Lett. 2023;10(1):119-24. <https://doi.org/10.51847/rX63RpnEhu>

Corresponding author: Sergey Nikolaevich Povetkin

E-mail ✉ ruslankalmykov777@yandex.ru

Received: 10/12/2022

Accepted: 04/03/2023

INTRODUCTION

Blood, or rather its components, is the main indicator of metabolism in the body [1-3]. Moving through a network of vessels and capillaries, blood contacts the cells of all tissues and organs,

thus providing the possibility of their nutrition and respiration, supplying them with the necessary enzymes, hormones, and other substances, without which the normal functioning of the body is impossible [4, 5]. Any impact on the body and the physiological

processes occurring in the body affect, to varying degrees, the composition of the blood. It is for this reason that, both in scientific research and in making a diagnosis, doctors often perform biochemical and hematological blood tests [6]. Such studies are especially important when making a diagnosis, selecting means of therapy and preventing pathologies, and studying the effect of various drugs on the body [7-9]. The information obtained during the performance of such studies allows us to more quickly and efficiently solve many problems and issues that arise in medicine.

In therapy aimed at correcting the hormonal background, many specialists intensively use various means, including non-hormonal ones: vitamin-mineral complexes, vitamins, micro-macroelements, etc. [10-13]. This stabilizes the state of the body, improves the quality and quantity of produced sperm in men, and improves the quality of eggs in women [14, 15]. However, few specialists seriously think about how exactly a drug or a complex of drugs affects the metabolic processes in the body, what happens to carbohydrate-lipid or protein-mineral metabolism, how quickly metabolic processes are activated, whether the means used have a hidden negative effect on organism [16-18].

In this scientific work, a study is made of the effect of taking similar non-hormonal drugs on biochemical blood parameters using the example of laboratory animals.

MATERIALS AND METHODS

The study of the effect of non-hormonal correction of the hormonal background on the biochemical parameters of the blood was carried out on laboratory animals.

The studies were carried out in accordance with the "Rules of Laboratory Practice in the Russian Federation" (Order of the Ministry of Health of the Russian Federation No. 708n dated August 23, 2010). Animal experiments were carried out in accordance with the rules adopted by the European Convention for the Protection of Vertebrate Animals Used for Experimental and Other Scientific Purposes [19].

Two similar groups of 25 young rams were selected for the study. The Rams were clinically healthy, of average weight and size, and aged 1.5 to 2.5 years. Animals of both groups were kept under the same conditions and on the same diet. Animals of the experimental group daily, 50 days, with feed were given 5 ml of a preparation containing iodine with amylopectin and, intramuscularly, were injected with 1.5 ml of E-selenium and a tissue preparation of 0.5 ml mixed with 1 ml of a 0.5% solution of novocaine. Animals of the control group were not prescribed or given any drugs. Blood samples were taken before the start of the study and at the end of the experiment.

To study physiological and biochemical changes in the body of laboratory animals, blood was taken from the vessels, in which the indicators of protein metabolism and carbohydrate-lipid metabolism were determined according to the standard method [20, 21].

To characterize the clinical state of the animals, body temperature was measured, pulse and respiration rates were determined, and the state of health and behavior were monitored.

RESULTS AND DISCUSSION

The results obtained were statistically processed and summarized in **Tables 1 and 2**.

Table 1. Indicators of protein metabolism in rams

Indicator	Standard	Groups	Numerical value	
			At the beginning of the experience	At the beginning of the experience
Total protein, g/l	42-97	Experience	20,37±0,67	43,59±5,80
		Control	28,11±2,30	54,18±13,75
Albumins, g/l	22,6-40,4	Experience	17,02±1,20	20,13±2,03
		Control	24,77±3,19	23,23±0,22
Globulins, g/l	35-49	Experience	3,35±1,26	23,46±7,84
		Control	3,34±0,89	30,96±13,53
Creatinine, mmol/l	53-174	Experience	154,41±57,22	169,27±22,57

		Control	181,33±58,14	194,04±49,45
Urea, mmol/l	3,3-9,3	Experience	4,02±0,91	5,103±1,66
		Control	2,75±0,23	3,24±0,65

In the course of the research, it was found that at the beginning of the experiment, hypoproteinemia, hypoalbuminemia, and hypoglobulinemia were observed in animals of both groups. At the same time, the level of creatinine and urea at the beginning of the experiment was almost normal. This is a sign that at the beginning of the experiment, the diet of the animals was not sufficiently balanced for some time, therefore, first of all, the diet was corrected and only then the experiment was carried out [22].

By the end of the experiment, the level of protein, albumin, and globulin in both groups changed significantly. So in both groups, the level of total protein was normal, but in the experimental group, the protein by the end of the experiment increased by 21.24% more than in the control. At the same time, the total protein concentration in the experimental group by the end of the experiment was 43.59 g/l, and in the control group, it was 54.18 g/l. Such dynamics can be explained only by the fact that the level of creatinine and urea also increased [23]. Consequently, in the rams of the experimental group during the period of the experiment, there was a more rapid recovery of the body after the correction of the diet.

Albumin in the experimental group increased by 17%, while in the control group, it decreased by 5.8%. At the same time, by the end of the experiment in both groups, the share of finely dispersed proteins - albumins accounted for about 50% of the total protein, which indicates good tissue hydrophilicity. At the same time, the level of globulins increased, which can be explained by an increase in the ability of the organism to a specific humoral response [24]. However, in the experimental group, the level of globulins increased by 31% less than in the control group. Consequently, in the experimental

group there was less need for the formation of immune globulins, which can only be explained by the fact that in the process of experimenting, the body received the necessary trace elements that improve immunity (E-selenium and iodine). Creatinine is a breakdown product of creatinine phosphate in muscles in the cycle of providing the body with energy for muscle contraction [25]. Given that the amount of produced creatinine is inextricably linked with the total body weight, in particular with muscle mass, we believe that the dynamics of creatinine concentration in our study showed the rate of recovery of muscle mass after the winter keeping of rams in barns. In the rams of the experimental group, the level of creatinine increased by 8.77%, and in the control group by 6.55%. Consequently, the restoration of muscle mass occurred more intensively in the animals of the experimental group.

Urea is one of the main indicators of the process of protein hydrolysis in the body [26]. The results of biochemical studies show that the concentration of urea in the experimental group at the beginning of the experiment is higher than in the control group, where the urea level is slightly below the minimum value of the norm. The high concentration of urea in the blood serum of rams of the experimental group indicates a high disintegration of feed protein [27]. Thus, it can be seen that in the rumen of rams, good hydrolysis of vegetable protein to amino acids occurs with further deamination of the protein to ammonia. By the end of the experiment in both groups, the level of urea increased, as there is an improvement in the cicatricial hydrolysis of feed proteins, which indicates an improvement in feed conversion. At the same time, feed conversion was better in the experimental group, where the urea level increased by 26%, while in the control by 17.8%.

Table 2. Indicators of carbohydrate-lipid metabolism in rams

Indicator	Standard	Groups	Numerical value	
			At the beginning of the experience	At the end of the experience
Glucose, mmol/l	2,5-3,3	Experience	3,13±0,43	4,52±1,12
		Control	3,6±0,69	4,715±0,55

Triglycerides, mmol/l	0,66-0,88	Experience	0,29±0,16	1,59±0,93
		Control	0,087±0,0017	0,77±0,16
Cholesterol, mmol/l	1,1-2,3	Experience	0,78±0,113	1,91±0,07
		Control	0,28±0,103	2,05±0,27

Carbohydrate metabolism was studied by the dynamics of glucose in blood serum. At the beginning of the experiment, the glucose level was within normal limits. Although in the control group, it was slightly above the upper limit of normal. During the experiment, the glucose content increased in both groups, but in the experimental group, this figure increased by 13% more than in the control group. As for the level of triglycerides, as one of the indicators of lipid metabolism, it can be seen that at the beginning of the experiment, the level of triglycerides was below the lower limit of normal in both groups [28]. This can be explained by the fact that the animals received an unbalanced diet in the winter. After the correction of the diet during the experiment, the level of triglycerides was restored to physiological norms. At the same time, in the experimental group, the level of triglycerides after recovery was 1.6 times lower compared to the level of recovery in the control group. Triglycerides enter the body not only with feed but are also synthesized in the liver from carbohydrates. In the experimental group, the glucose level is higher than in the control, therefore, such indicators of triglycerides and glucose in the experimental group can also be explained by the fact that in the experimental group, there was a lower utilization of glucose for the formation of triglycerides. Such dynamics can be explained by the fact that the organism of animals in the experimental group recovered more quickly by the end of the experiment. The same fact confirms the level of cholesterol, a source of lipids for lipid membranes during the formation of cells, in particular, the lipid membrane of spermatozoa [29]. So, in the experimental group, the cholesterol level at the beginning of the experiment was low in both groups, during the experiment it increased several times and by the end of the experiment was about 2 mmol/l in both groups [30]. It can be seen that in the experimental group, by the end of the experiment, the degree of cholesterol recovery was normal but 2.98 times lower than in the control. Such dynamics can be explained precisely by the fact that for the synthesis of cell

walls, lipids were primarily utilized by the body of rams of the experimental group, while triglycerides were used for other metabolic processes. It should be noted that in the experimental group, all processes in the body of rams during their preparation for the breeding period proceeded more quickly and with less stress for the immune system.

CONCLUSION

In this scientific article, the influence of a non-hormonal complex, designed to correct the hormonal background of the body, was studied using the example of sheep. Animals of the experimental group daily, for 50 days, were given 5 ml of a preparation containing iodine with amyloextrin and were injected with 1.5 ml of E-selenium and 0.5 ml of a tissue preparation mixed with 1 ml of a 0.5% novocaine solution. At the end of the experiment, the dynamics of protein and carbohydrate-lipid metabolism in laboratory animals were studied. It has been established that the non-hormonal agents used do not hurt the body and contribute to a more optimal recovery of the body.

ACKNOWLEDGMENTS: None

CONFLICT OF INTEREST: None

FINANCIAL SUPPORT: The study was funded by a grant from the Ministry of Science and Higher Education of the Russian Federation "Study of the mechanisms of interaction of lactic acid microorganisms, lactose-fermenting yeast and biologically active substances during microcapsulation of various fractions of microbiota" by Decree of the Government of the Russian Federation No. 220 in the form of a subsidy from the federal budget for state support of scientific research conducted under the leadership of leading scientists of Russian educational institutions of higher education, scientific institutions and state scientific centers of the Russian Federation (IX stage), Agreement No. 075-15-2022-1129 01.07.2022.

ETHICS STATEMENT: The protocol for experiments with animals complied with the requirements of the European Convention for the Protection of Vertebrate Animals used for Experimental and other Scientific Purposes.

REFERENCES

- Nobandegani AS, Motamedifar M. Antibiotic Sensitivity Profile of the Bacterial Isolates from the Blood Samples of the Patients in Different Wards of a Major Referral Hospital, Shiraz, Iran 2015-2016. *Pharmacophore*. 2019;10(2):30-6.
- Van Ta T, Nguyen HB, Tran HT. Evaluating the Association of Red Blood Cell Parameters and Glycemic Control in Type 2 Diabetic Patients at Tien Giang General Hospital. *Arch Pharma Pract*. 2019;10(4):153-9.
- Yusransyah, Halimah E, Suwantika AA. Effect of pharmacist counseling on adherence and blood pressure of hypertensive Prolanis patients in sixteen primary healthcare centers. *J Adv Pharm Edu Res*. 2020;10(4):8-14.
- Murrant CL, Fletcher NM. Capillary communication: the role of capillaries in sensing the tissue environment, coordinating the microvascular, and controlling blood flow. *Am J Physiol Heart Circ Physiol*. 2022;323(5):H1019-36. doi:10.1152/ajpheart.00088.2022
- Mračková J, Růžičková T, Ševčík P, Karlíková M, Mraček J, Polívka J. Role of blood biomarkers in spontaneous intracerebral hemorrhage - a prospectively studied cohort of patients. *Cas Lek Cesk*. 2020;159(5):185-91.
- Jørgensen HL, Lind BS. Blood tests - too much of a good thing. *Scand J Prim Health Care*. 2022;40(2):165-6. doi:10.1080/02813432.2022.2104436
- Racaru-Honciuc V, Betea D, Scheen AJ. Hormonal deficiencies in the elderly: is there a role for replacement therapy?. *Rev Med Suisse*. 2014;10(439):1555-6, 1558-61. [In French].
- Orsaeva AT, Tamrieva LA, Mischvelov AE, Osadchiy SS, Osipchuk GV, Povetkin SN, et al. Digital clinic "smart ward. *Pharmacophore*. 2020;11(1):142-6.
- Raevskaya AI, Belyalova AA, Shevchenko PP, Karpov SM, Mishvelov AE, Simonov AN, et al. Cognitive Impairments in A Range of Somatic Diseases Diagnostics, Modern Approach to Therapy. *Pharmacophore*. 2020;11(1):136-41.
- Osipchuk GV, Povetkin SN, Simonov AN, Verevkina MN, Karatunov VA, Yakovets MG. On the Issue of Non-Hormonal Stimulation of the Reproductive Function of Rams, *Pharmacophore*. 2020;11(2):73-6.
- Ziruk IV, Egunova AV, Kopchekchi ME, Frolov VV, Babina KI, Povetkin SN, et al. Morphometric of Pig Livers under Different Doses of Minerals in Feed Allowance. *Int Trans J Eng, Manag, Appl Sci Technol*. 2020;11(14):11A14L, 1-10. doi:10.14456/ITJEMAST.2020.277
- Maslova AY, Tskaeva AA, Ashurova ZA, Abazova A, Ismailov MM, Ismailova MM, et al. Study of the effect of Baricitinib on the Course of COVID-19. *J Pharm Res Int*. 2021;33(35A):204-13.
- Bertolla RP. Sperm biology and male reproductive health. *Sci Rep*. 2020;10(1):21879. doi:10.1038/s41598-020-78861-7
- Yoshimura Y, Barua A. Female Reproductive System, and Immunology. *Adv Exp Med Biol*. 2017;1001:33-57. doi:10.1007/978-981-10-3975-1_3
- Nazar FN, Magnoli AP, Dalcero AM, Marin RH. Effect of feed contamination with aflatoxin B1 and administration of exogenous corticosterone on Japanese quail biochemical and immunological parameters. *Poult Sci*. 2012;91(1):47-54. doi:10.3382/ps.2011-01658
- Brinkworth GD, Noakes M, Keogh JB, Luscombe ND, Wittert GA, Clifton PM. Long-term effects of a high-protein, low-carbohydrate diet on weight control and cardiovascular risk markers in obese hyperinsulinemic subjects. *Int J Obes Relat Metab Disord*. 2004;28(5):661-70. doi:10.1038/sj.ijo.0802617
- Maslova AY, Mishvelov AE, Dudusheva MJ, Blejyants GA, Minaev SV, Shchetinin EV, et al. Using a Decision Tree with a Feedback Function to Select Therapeutic Tactics for Viral Infection of the Respiratory Tract in the Medical Expert System. *Int Trans J Eng*

- Manag Appl Sci Technol. 2022;13(8):13A8F, 1-10.
18. Galabueva AI, Biragova AK, Kotsoyeva GA, Borukayeva ZK, Yesiev RK, Dzgoeva ZG. Optimization of Modern Methods of Treating Chronic Generalized Periodontitis of Mild Severity. *Pharmacophore*. 2020;11(1):47-51.
 19. Blinov AV, Nagdalian AA, Povetkin SN, Gvozdenco AA, Verevkin MN, Rzhepakovsky IV, et al. Surface-Oxidized Polymer-Stabilized Silver Nanoparticles as a Covering Component of Suture Materials. *Micromachines*. 2022;13(7):1105. doi:10.3390/mi13071105
 20. Cheprasova AA, Popov SS, Pashkov AN, Verevkin AN, Krylâ ED, Mittova VO. Oxidative status, carbohydrate, and lipid metabolism indicators in saliva and blood serum of type 1 diabetes mellitus patients. *Biomed Res Ther*. 2022;9(8):5233-40. doi:10.15419/bmrat.v9i8.761
 21. Chen W, Chen G. The Roles of Vitamin A in the Regulation of Carbohydrate, Lipid, and Protein Metabolism. *J Clin Med*. 2014;3(2):453-79. doi:10.3390/jcm3020453
 22. Bachinina KN, Povetkin SN, Simonov AN, Pushkin SV, Blinova AA, Sukhanova ED, et al. Effects of Selenium Preparation on Morphological and Biochemical Parameters of Quail Meat. *Int Trans J Eng Manag Appl Sci Technol*. 2021;12(13):1-7.
 23. Ilyasov KK, Demchenkov EL, Chernyshkov AS, Rodin IA, Pushkin SV, Povetkin SN, et al. Features of the phytopharmacological preparations in the metaphylaxis of urolithiasis. *Pharmacophore*. 2020;11(5):66-71.
 24. Bartsch YC, Fischinger S, Siddiqui SM, Chen Z, Yu J, Gebre M, et al. Discrete SARS-CoV-2 antibody titers track with functional humoral stability. *Nat Commun*. 2021;12(1):1018. doi:10.1038/s41467-021-21336-8
 25. Pandhi P, Streng KW, Anker SD, Cleland JG, Damman K, Dickstein K, et al. The value of spot urinary creatinine as a marker of muscle wasting in patients with new-onset or worsening heart failure. *J Cachexia Sarcopenia Muscle*. 2021;12(3):555-67. doi:10.1002/jcsm.12690
 26. Vitolo M. Notes on urea hydrolysis by urease. *World J Pharm Pharm Sci*. 2022;11:96-135. doi:10.20959/wjpps20223-21380
 27. Liu M, Li M, Liu J, Wang H, Zhong D, Zhou H, et al. Elevated urinary urea by high-protein diet could be one of the inducements of bladder disorders. *J Transl Med*. 2016;14:1-7. doi:10.1186/s12967-016-0809-9
 28. Dzhabrailova US, Vagabov VM, Akhaeva ZN, Kasimova ZZ, Kolesnikov SP, Bondarenko NG. Characterization of Physico-Chemical Parameters and Toxicological Properties of Neocytin. *Pharmacophore*. 2022;13(5):44-50. doi:10.51847/igW0BabCma
 29. Bernabò N, Machado-Simoes J, Valbonetti L, Ramal-Sanchez M, Capacchietti G, Fontana A, et al. Graphene Oxide increases mammalian spermatozoa fertilizing ability by extracting cholesterol from their membranes and promoting capacitation. *Sci Rep*. 2019;9(1):8155. doi:10.1038/s41598-019-44702-5
 30. Tanigawa K, Luo Y, Kawashima A, Kiriya M, Nakamura Y, Karasawa K, et al. Essential Roles of PPARs in Lipid Metabolism during Mycobacterial Infection. *Int J Mol Sci*. 2021;22(14):7597. doi:10.3390/ijms22147597