



The Efficiency of Ethanolic Extract of *Ocimum basilicum* Leaves and Flowers against Mosquito Larvae

Hassan Ahmed Rudayni¹, Nosiba Suliman Basher¹, Lamyah Ahmed AL-keridis^{2*},
Nasir Adam. Ibrahim³, Elnour Abdelmageed⁴

¹Department of Biology, Faculty of Science, Imam Mohammed Ibn Saud Islamic University, Riyadh, Saudi Arabia.

²Department of Biology Faculty of Science, Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia.

³Biochemistry and Physiology Department, Faculty of Veterinary Medicine, Al-butana University, Rufaa, Sudan.

⁴Department of Biology and Chemistry, Faculty of Education, University of Al Qadarif, Al Qadarif, Sudan.

ABSTRACT

Mosquito control remains a powerful method to control mosquito-transmitted diseases. Focusing on early stages is very important for many reasons because they are easy to handle, significantly affected by ambient environmental influences, natural, and chemically toxic. The current study was aimed to examine the chemical components and larvicidal effect of Ethanolic extract of *O. basilicum* leaves and flowers on mosquito larvae. Fresh *Ocimum basilicum* was collected, dried in the shade for one week, and extracted with ethanol. The larvae were placed in plastic containers and bioassayed according to the protocol approved by WHO. After 24 hours of the recovery period, mortality percentage was recorded. Dead larvae were examined under a microscope to observe any morphological changes. The statistical analysis using SPSS program version 19 was done to determine Probit data. The phytochemical investigation of *O. basilicum* leaves presented many components such as flavonoids, glycosides, tannins, and steroids, whereas alkaloids, saponins, and terpenoids were absent.

Moreover, *O. basilicum* flowers showed the presence of only tannins, terpenoids, steroids, and flavonoids, but not Saponins, glycosides alkaloids. The action of leaves and flower extracts were given LC₅₀ values of 17.78 ppm, 16.98 ppm, 15.48 ppm, and 15.84 and LC₉₅ values were 56.23 ppm, 64.56 ppm, 66.06 ppm, and 50.11 ppm against larvae of *Anopheles. Arabiensis* and *Culex. quinquefasciatus*, respectively. Moreover, when treated with the ethanolic extract, mosquito larvae showed some morphological changes, such as decoloration and alimentary canal deformity.

Keywords: Larvicidal, Ethanol extract, *Ocimum basilicum*, *Anopheles arabiensis*, *Culex quinquefasciatus*, phytochemical.

HOW TO CITE THIS ARTICLE: Rudayni HA, Basher NS, AL-keridis LA, Ibrahim NA, Abdelmageed E. The Efficiency of Ethanolic Extract of *Ocimum basilicum* Leaves and Flowers against Mosquito Larvae. Entomol Appl Sci Lett. 2021;8(3):46-53. <https://doi.org/10.51847/5wPMV7xyXL>

Corresponding author: Lamyah Ahmed AL-keridis

E-mail ✉ laalkeridis@pnu.edu.sa

Received: 17/04/2021

Accepted: 28/08/2021

INTRODUCTION

Mosquitoes, including *Culex pipiens*, *Anopheles*, and *Aedes* which belong to the family of Diptera: Culicidae, are responsible for transmitting many diseases. In addition to their higher cost, chemical insecticides remain one of the main factors that cause damage to the environment.

Most mosquitoes are becoming resistant to insecticides [1, 2]. Research in the last years has concentrated on the discovery of new active compounds from plants as natural products [3]. Plants are significant resources for discovering new natural products to replace chemical ones [4]. Many mosquito insecticides derived from plants proved to be inhibitors [5].

Ocimum basilicum L - belongs to the family Lamiaceae - is used globally in folk medicine [6]. The *Aedes aegypti*, showed high sensitivity to *Ocimum basilicum* L leaf extract with concentrations of 0.3%, 0.6%, 0.9%, 1.2%, 1.5% [7]. Basal leaf powder, when extracted by ethanol, becomes more toxic to insects. Therefore, it is recommended to use for controlling American cockroaches [8]. Basil essential oils were more effective against the third instar of *A. aegypti* larvae. It gives the best LC₅₀ and LC₉₀ values when applied in the laboratory [9, 10]. Oil constituents gave 100% mortality when tested against mites [11]. The basil leaves gave good results when used against agriculture pests in rice and showed a high mortality rate [12]. The current study was aimed to examine the chemical components and larvicidal effect of Ethanolic extract of *O. basilicum* leaves and flowers on mosquito larvae. The study results would likely be aimed to contribute to the search for natural products in mosquito control programs.

MATERIALS AND METHODS

Plant material

Fresh samples of *Ocimum basilicum* were collected randomly from the garden of the College of Science, University of Imam Mohammed Iben Saud Islamic, where plant parts were processed and identified by a department member. Leaves and flowers samples were dried in the shade for one week, then were powdered and kept in a sealed plastic container until using for phytochemical investigation and bioassay experiments.

Phytochemical screening of extracts

The selected parts of *O. basilicum* were screening preliminary phytochemical to identify the chemical constituents (main classes). The methods described by Mohamed Nour (2009) [13]. were used to screen alkaloids, saponins,

tannins, flavonoids, glycosides, steroids, and terpenoids.

Preparation of ethanolic extracts

The plant parts (leaves and flowers) were extracted by ethanol, according to Nosiba S *et al.* [14].

Larvae of mosquitoes

Under laboratory conditions, mosquito larvae were collected and reared in the Biology Department laboratory, Faculty of Science. The larvae were placed in plastic containers and provided with all nutrients required.

Larval bioassay

The larvicidal application was performed according to the protocol approved by WHO [15]. The test included 20 larvae (third and early fourth instar stages) of mosquito *A. arabiensis*, and *C. quinquefasciatus* put 250 ml of tap water into container cups. Sequences of concentrations were used. Each attention was of three replicates of extract, in addition to control groups without section. After 24 hours, the mortality percentage was recorded. Dead larvae were counted and examined under a microscope to observe morphological changes.

Data and statistical analysis

Data were analyzed using SPSS program version 19 to determine Probit parameters that included: percentage mortalities of larvae *A. arabiensis* and *C. quinquefasciatus*, LC₅₀ and LC₉₀ values at specific P-value, slope, regression (R²), and x-coefficient.

RESULTS AND DISCUSSION

The phytochemical constituents of *O. basilicum* leaves and flowers

O. basilicum leaves showed the variant amount of phytochemical components (**Table 1**).

Table 1. The phytochemical constituents of *O. basilicum* leaves and flowers

Plant part	Alkaloids	Saponins	Tannins	Flavonoids	Glycosides	Steroids	Terpenoids
<i>O. basilicum</i> leaves	-	-	+	+	+	+	-
<i>O. basilicum</i> flowers	-	+	-	+	++	-	+

- means the absence of the main class

+ indicates the presence of the main class

++ implies the presence of the main class in a relatively higher concentration

The action of ethanol extract of leaves and flower of basil plants on mosquito larvae (24hrs)

The results obtained by the current study showed that leave ethanol extracts of basil plant

tested at different concentrations of the lethal concentration (LC₅₀), which was 17.78 ppm for *Anopheles arabiensis* and 16.98 ppm for *Culex quinquefasciatus* larvae. Meanwhile, flowers ethanol extract at different concentrations showed lethal concentration (LC₅₀) that was 15.48 ppm for *Anopheles arabiensis* and 15,84

ppm for *Culex quinquefasciatus* larvae (Tables 2 and 3; Figures 1 and 2).

According to LC₅₀ value, the flower presented good action against *Anopheles* larvae compared to *Culex* larvae. Additionally, the flowers extract indicated more potency and biological activity compared to the extract obtained from leaves.

Table 2. The action of ethanol extract of leaves on mosquito larvae (24hrs)

Conc. (ppm)	Log- Conc	<i>A. arabiensis</i>		<i>C. quinquefasciatus</i>	
		Mortality %	Probit	Mortality %	Probit
59,3	1.773	97	6.88	92	6.41
47,44	1.676	82	5.92	87	6.13
30,08	1.551	75	5.07	80	5.84
23,72	1.375	57	5.18	52	5.05
11,86	1.074	47	4.92	32	4.53
R2		0.64		0.97	
slope		2.36		2.79	
x-coefficient		2.07		1.43	
LC ₅₀		17.78 ppm		16.98 ppm	
LC ₉₅		56.23 ppm		64.56 ppm	




Figure 1. Log- Probit curve of action of ethanol extract of leaves on mosquito larvae (24hrs)

Table 3. The action of ethanol extract of flower on mosquito larvae (24hrs)

Conc. (ppm)	Log- Conc	<i>A. arabiensis</i>		<i>C. quinquefasciatus</i>	
		Mortality %	Probit	Mortality %	Probit
58.56	1.76	90	6.28	97	6.88
46.86	1.67	85	6.04	87	6.13
35.15	1.54	72	5.58	77	5.74
23.43	1.36	57	5.18	62	5.31
5.86	0.76	35	4.61	30	4.48
R2		0.912		0.88	
slope		1.60		2.12	
x-coefficient		3.25		2.69	
LC ₅₀		15.48ppm		15.84ppm	
LC ₉₅		66.06ppm		50.11ppm	




Figure 2. Log- Probit curve of action of ethanol extract of flower on mosquito larvae. (24hrs)

The damage on mosquito larvae caused by basil plant extracts

Mosquitoes larvae treated with plant parts section of basil revealed larval malformation. Discoloration (bright color), disconnected

alimentary canal, and swollen alimentary canal that was not attached to the head were the main changes observed (**Figure 3**).



Control larva



Bright color larva



Larvae swollen alimentary canal that was not attached to head



Larva disconnected the alimentary canal

Figure 3. The Damage on mosquito larvae caused by basil plant extracts

The mosquitoes are a series of vectors transmitted diseases, and control plans remained reflected as a very difficult triangle, both in health and entomological sciences [16]. Recently, the use of plant products received significant attention and proved to be

alternative sources as parasitic control agents since - compared to synthetic products - they constitute a rich source of eco-friendly bioactive compounds. The plant extracts are suitable and safe alternatives due to their low toxicity to mammals and easy biodegradability [17, 18].

Identifying novel effective mosquitocidal compounds are essential to overcome the increasing resistance rates of synthetic insecticide concern for the environment and food safety, the unacceptability of many chemicals insecticide [19]. The *Ocimum basil* proved to have richness in phytochemical components such as terpenoids, alkaloids, phenolics, flavonoids, tannins, saponin, reducing sugars, cardiac glycosides, steroids, and glycosides that cause pharmacological advantages [20]. The current study showed that the phytochemical constituents of *O. basilicum* in leaves were tannins, glycosides, steroids, and flavonoids, while saponins, alkaloids, and terpenoids were absent. On the other hand, *O. basilicum* flowers showed the presence of tannins, terpenoids, steroids, and flavonoids. While, Saponins, glycosides alkaloids were not detected.

The study suggested that the basil content phytochemicals act more toxic to mosquito leave and other stages. In the present study, the larvicidal activity of *O. basilicum* seemed to be supported by Azhari *et al.* [9]. Further, it has the larvicidal potential and repellent against the dengue vector *Ae. aegypti*. Also, it showed toxicity in experimental animals. The study of the insecticidal potentiality of basil was tested against *Culex quinquefasciatus* (Say) under laboratory conditions. Five concentrations of aqueous plant extracts against the 4th instar larvae for 24 and 48h were used after exposure. The LC₅₀ values demonstrated (5.32%) [21]. The previous results support our results and suggest that bioactive basil content can be more effective with water extract and other solvents. *Ocimum basilicum* L leaves extracted by ethanol and hexane and applied against the larvae of *Anopheles arabiensis* Patton. Results showed that basil leaves hexane extracts at the rate of 10% were the best to repel the adult insect for up to 2h [22]. This result is not similar to the result obtained by the current study because the application was in adult mosquitoes. Moreover, when the ethanol extract efficacy was considered, it supports our findings. In the dose-response test of *Ocimum basilicum* (*O. basilicum*), a high repellency against *Ae. Aegypti* is exhibited and a strong, effective dose against *Cx. quinquefasciatus* is obtained. In addition, the two essential oils exhibited

moderate repellency against *Ae. aegypti*, *An. dirus* and *Cx. quinquefasciatus*, at 60, 90, and 78 min with *C. nardus*, and 54, 96, and 72 min with *S. aromaticum*, respectively [23]. These results were consistent with results obtained by the current study. The leaves powder (LP) of *Ocimum basilicum* is extracted by ethanolic and used as a larvicide of *Anopheles arabiensis* 3rd instar larvae. The study accomplished malaria control findings by using basil as a bio-agent (Basil leaves extracts LC₅₀ of 58mg/l and LC₉₀ of 143mg/l) [24]. The previous studies were similar to the results obtained by the current study. Many plants may affect the mosquito that given lethal effect against *Anopheles* and *Culex larvae*, it can be recommended to be used in mosquito control programs [15, 24, 25]. Many researcher suggest that essential oils of *Ocimum basilicum* could be used as natural repellent [26, 27] *Anopheles larvae* is more susceptible than *Culex* [28]. Kumar *et al.* recommended the efficient use of basil leaf essential oil as an efficient repellent and as a moderate larvicide agent against *Ae. aegypti* [29, 30]. The previous studies indicate that the plant has the ability to inhibit mosquito larvae and control it without using chemical pesticides due to the sensitivity of mosquito stages. Some plants such as *L. Camara A. indica* leaves, and flower leaves give changes in morphology [14, 31]. This finding is similar to the results of the current study. Changes in *O. basilicum* leaves, and flower ethanol extracts in morphological (Bright color, swelled disconnected alimentary canal and head) of *A. arabiensis* and *C. quinquefasciatus* were noted.

CONCLUSION

The study concluded that according to LC's values obtained, the plant extract displays more potentiality against *A. arabiensis* larvae than *C. quinquefasciatus* larvae, and the flowers were better than leaves in mortality rate. Therefore, the study recommended that *O. basilicum* ethanol extract is the best natural larvical and it is eco-friendly.

ACKNOWLEDGMENTS: This research was funded by the Deanship of Scientific research at Princess Nourah bint Abdulrahman University through the fast-track Research Funding Program 2021.

CONFLICT OF INTEREST: None

FINANCIAL SUPPORT: This work has received funds from The Deanship of Scientific research, Princess Nourah bint Abdulrahman University through the fast-track Research Funding Program, 2021.

ETHICS STATEMENT: This work was approved by Ethics Committee of Imam Mohammed Ibn Saud Islamic University. The data obtained from this work were analyzed according guidelines of ethical standards of the Declaration of Helsinki.

REFERENCES

1. El Ouali Lalami A, El-Akhal F, Ez Zoubi Y, Taghzouti K. Study of Phytochemical Screening and Larvicidal Efficacy of Ethanolic Extract of *Salvia officinalis* (Lamiaceae) from North Center of Morocco against *Culex pipiens* (Diptera: Culicidae) Vector of Serious Human Diseases. *Int J Pharmacog Phytochem Res.* 2016;8(10):1663-8. Available online on www.ijppr.com
2. Krishnappa K, Pandiyan J, Elumalai K, Baranitharan M, Jayakumar S, Gokulakrishnan J. GC-MS analysis and mosquitocidal properties of *Loranthes pentandrus* Linn.(Loranthaceae) against human vector mosquitoes (Diptera: Culicidae). *Acad J Med Plants.* 2019;7(12):261-8. doi:10.15413/ajmp.2019.0154
3. Lupi E, Hatz C, Schlagenhaut P. The efficacy of repellents against *Aedes*, *Anopheles*, *Culex* spp. A literature review. *Trav Med Infect Dis.* 2013;11(6):374-411.
4. Asadollahi A, Khoobdel M, Zahraei-Ramazani A, Azarmi S, Mosawi SH. Effectiveness of plant-based repellents against different *Anopheles* species: a systematic review. *Malar J.* 2019;18(1):1-20. doi:10.1186/s12936-019-3064-8.
5. Sukumar K, Perich MJ, Boobar LR. Botanical derivatives in mosquito control: a review. *J Am Mosq Control Assoc.* 1991;7(2):210-37.
6. Miele M, Dondero R, Ciarallo G, Mazzei M. Methyleugenol in *Ocimum basilicum* L. Cv. *genovese gigante*. *J Agric Food Chem.* 2001;49(1):517-21.
7. Husna I, Setyaningrum E, Handayani TT, Kurnia Y, Palupi EK, Umam R, et al. Utilization of Basil Leaf Extract as Anti-Mosquito Repellent: A Case Study of Total Mosquito Mortality (*Aedes aegypti* 3rd Instar). *J Phys: Conf Ser.* 2020;1467(1):012014.
8. Ukoroije BR, Island W, Island W, Island W. The efficacy of *Ocimum gratissimum* leaf powder and ethanol extract on adult *Periplaneta americana* under laboratory condition. *Open Access Libr J.* 2018;5(04):1. doi:10.4236/oalib.1104455
9. Azhari H, Abdurahman H, Mashitah M, DO SJ. Bioactive compounds from Basil (*Ocimum basilicum*) essential oils with larvicidal activity against *Aedes aegypti* larvae. 3rd Int. Conf Bio, Env Chem, Singapore; 2012;46:21-4.
10. Amer A, Mehlhorn H. Larvicidal effects of various essential oils against *Aedes*, *Anopheles*, and *Culex* larvae (Diptera, Culicidae). *Parasitol Res.* 2006;99(4):466-72.
11. Perumalsamy H, Kim JY, Kim JR, Hwang KN, Ahn YJ. Toxicity of basil oil constituents and related compounds and the efficacy of spray formulations to *Dermatophagoides farinae* (Acari: Pyroglyphidae). *J Med Entomol.* 2014;51(3):650-7.
12. López MD, Jordán MJ, Pascual-Villalobos MJ. Toxic compounds in essential oils of coriander, caraway and basil active against stored rice pests. *J Stored Prod Res* 2008;44(3):273-8.
13. Mohamed Nour AA. Chemical component of the essential oil of basil (*Ocimum basilicum* L.) active against *Salmonella typhi* [Doctoral dissertation, M. Sc. Thesis]. Sudan: University of Gezira; 2009.
14. Hamid NS, Kehail MA, Ibrahim NA, Abdel-Rahman EH. Larvicidal activity of ethanol extracts of *Azadirachta indica* (neem) against *Anopheles Arabiensis* and *Culex Quinquefasciatus* larvae, Gezira state, Sudan. *Int J Entomol Res.* 2021;6(1):138-41.
15. World Health Organization. Pesticides and their application: for the control of vectors and pests of public health importance. World Health Organization; 2006. WHO/CDS/WHOPES/GCDPP/2006.1.

16. Yousif HY. Susceptibility of *Anopheles arabiensis* Patton and *Culex quinquefasciatus* Say Larvae to *Ixora* (*Ixora coccinea* L) Leaves and Flowers Extracts [dissertation on the internet]. Sudan: University of Gezira; 2013. [cited 2013 May 16]. Available from: <http://repo.uofg.edu.sd/handle/123456789/2165>
17. Aouati A, Roubhi AH, Boudjahem I, Berchi S. Study of toxicological activity of the essential oil of *Ocimum basilicum* L. against *Culicida* larvae in Algeria. AIP Conf Proc. 2019;2190(1):020073. doi:10.1063/1.5138559
18. Madhumitha G, Rajakumar G, Roopan SM, Rahuman AA, Priya KM, Saral AM, et al. Acaricidal, insecticidal, and larvicidal efficacy of fruit peel aqueous extract of *Annona squamosa* and its compounds against blood-feeding parasites. Parasitol Res. 2012;111(5):2189-99. doi:10.1007/s00436-011-2671-2
19. Shaalan EA, Canyon D, Younes MW, Abdel-Wahab H, Mansour AH. A review of botanical phytochemicals with mosquitocidal potential. Environ Int. 2005;31(8):1149-66. doi:10.1016/j.envint.2005.03.003.
20. Al-Snafi AE. Chemical constituents and pharmacological effects of *Ocimum basilicum*-A review. Int J Pharm Res. 2021;13(2):2997-3013.
21. Iqbal J, Ishtiaq F, Alqarni AS, Owayss AA. Evaluation of larvicidal efficacy of indigenous plant extracts against *Culex quinquefasciatus* (Say) under laboratory conditions. Turk J Agric For. 2018;42(3):207-15. doi:10.3906/tar-1711-69
22. Elsiddig FEI, Khei SEM. Effect of Neem, Cafure, and Rehan Organic Extracts Compared to Other Chemical and Biological Standard Insecticides on *Anopheles arabiensis* Patton Mosquito. [dissertation on the internet]. Agricultural Studies College of Agricultural Studies: Sudan University of Science and Technology; 2007. [cited 2007 April 1]. Available from: <http://repository.sustech.edu/handle/123456789/3504>
23. Phasomkusolsil S, Soonwera M. Comparative mosquito repellency of essential oils against *Aedes aegypti* (Linn.), *Anopheles dirus* (Peyton and Harrison) and *Culex quinquefasciatus* (Say). Asian Pac J Trop Biomed. 2011;1(1):S113-8. doi:10.1016/S2221-1691(11)60136-6
24. Ali, HE. Effect of Basil (*Ocimum basilicum* L.) Leave Powder and Ethanolic Extract on the 3rd Larval Instar of *Anopheles arabiensis* (Patton,1905) *Culicidae: Diptera*. [dissertation on the internet]. Sudan: University of Gezira; 2013. [cited 2013 January 31]. Available from: <http://repo.uofg.edu.sd/handle/123456789/2171>
25. Panneerselvam C, Murugan K, Kovendan K, Kumar PM, Subramaniam J. Mosquito larvicidal and pupicidal activity of *Euphorbia hirta* Linn.(Family: Euphorbiaceae) and *Bacillus sphaericus* against *Anopheles stephensi* Liston.(Diptera: Culicidae). Asian Pac J Trop Med. 2013;6(2):102-9. doi:10.1016/S1995-7645(13)60003-6
26. Ileke KD, Adesina JM. Toxicity of *Ocimum basilicum* and *Ocimum gratissimum* extracts against main malaria vector, *Anopheles gambiae* (Diptera: Culicidae) in Nigeria. J Arthropod Borne Dis. 2019;13(4):362.
27. Adam AA, Ahmed SA, Mohamed TA, Azrag RA, Mustfa SE, Hamdi AA. Evaluation of repellent activities of the essential oil of *Ocimum basilicum* against *Anopheles* mosquito and formulation of mosquitoes repellent cream. Biomed Res Clin Prac. 2019;4:2-5. doi:10.15761/BRCP.1000184
28. Abdalla AI, Kehail MA, Abdelrahim YM, Ibrahim NA. Phytochemical screening of *Calotropis procera* ait flower parts and their larvicidal potentialities against *Anopheles* and *Culex* Larvae, Gezira State, Sudan. Int J Biol Res. 2017;2(2):88-92.
29. Govindarajan M, Rajeswary M. Ovicidal and adulticidal potential of leaf and seed extract of *Albizia lebbek* (L.) Benth. (Family: Fabaceae) against *Culex quinquefasciatus*, *Aedes aegypti*, and *Anopheles stephensi* (Diptera: Culicidae). Parasitol Res. 2015;114(5):1949-61. doi:10.1007/s00436-015-4384-4

30. Kumar S, Warikoo R, Mishra M, Samal RR. Impact of *Ocimum basilicum* Leaf Essential Oil on The Survival and Behaviour of an Indian Strain of Dengue Vector, *Aedes aegypti* (L.). *Vector Biol J.* 2017;2:2. doi:10.4172/2473-4810.1000122
31. Alghamdi AA, Basher NS. Efficacy of leaves and flowers ethanol extracts of the invasive species *Lantana camara* Linn as a Mosquito larvicidal. *Int J Mosq Res.* 2020;7(5):43-7.