



## Antimicrobial Bio-Components from Red Algae Species: a Review of Application and Health Benefits

Sivak Elena<sup>1\*</sup>, Bugaev Sergey<sup>2</sup>, Sokolov Mikhail<sup>3</sup>, Glinushkin Alexey<sup>4</sup>

<sup>1</sup>Associate Professor, Doctor of Agricultural Sciences, Department of Standardization and Equipment of Processing Industries, Kursk State Agricultural Academy named after I.I.

Ivanov, Kursk, Russia

<sup>2</sup>Associate Professor, Candidate of Agricultural Sciences, Department of general zootechnics, Kursk State Agricultural Academy named after I.I. Ivanov, Kursk, Russia

<sup>3</sup>Academician of the Russian Academy of Sciences, Professor, Department of Herbology, All Russian Research Institute of Phytopathology, Moscow Region, Russia

<sup>4</sup>Professor of the Russian Academy of Sciences, Doctor of Agricultural Sciences, Department of resistance studies, All Russian Research Institute of Phytopathology, Moscow Region, Russia

### ABSTRACT

Primarily, Seaweeds contain beneficial metabolites for human body. Algae, a group of seaweeds, have potential of being used to produce food products, pharmaceutical products, biodiesel, etc. There are several groups of Algae throughout the world like, red, green algae etc. Some natural metabolites of red algae are steroid, terpenoid, acetogenin. Regarding growing awareness of consumers on application of natural antimicrobial products, it could be noted that Marine seaweed and algae could be considered as proper sources. The aim of this review is to present a brief description of antimicrobial properties of red algae.

**Keywords:** Red algae, antimicrobial compounds, seaweeds

**HOW TO CITE THIS ARTICLE:** Sivak Elena\*, Bugaev Sergey, Sokolov Mikhail, Glinushkin Alexey, Antimicrobial bio-components from red algae species: a review of application and health benefits, Entomol Appl Sci Lett, 2018, 5 (3):85-90.

**Corresponding author:** Sivak Elena

**E-mail** ✉ [kursknich@gmail.com](mailto:kursknich@gmail.com)

**Received:** 18/03/2018

**Accepted:** 29/07/2018

### INTRODUCTION

To date, some bioactive compounds have been identified with potentials of treating toward HIV, Ebola, Hepatitis C, and H5N1 virus. Moreover, sulfated polysaccharide found in red algae might attenuate retroviral replications. In general, seaweeds are considered as one of the superb components of marine organisms and currently are used as vegetables in diets (Nisizawa). People from coastal beach area consume both fresh and dry forms of seaweeds [1]. Algae are in a wide range of utilizations such as food, medicine, etc. Seaweeds have 13000 years' history of using in different applications [2, 3]. Phytochemical Substances derived from red algae are frequently used in industries like food, pharmacy, textile, etc. Seaweeds are rich in enzymes, antioxidants, functional components.

They also have medicinal applications like antibiotics Swaminathan and anticoagulant [4]. Recently, seaweeds (macro-algae) are being used for many applications such as food [5]. With respect to parameters like pigment and morphological characteristics, seaweeds are classified into three groups: red algae, brown algae and green algae, scientifically known as Rhodophyceae, Phaeophyceae and Chlorophyceae, respectively [6]. Several technological applications for seaweeds could be defined like using as thickening or gelling agent, and as vegetables. Additionally, seaweeds contain some groups of gum including carrageenan, agar, and alginat [7]. Incorporating seaweeds into pasta makes it possess biologically active compounds such as fucoxanthin and fucosterol [8]. Furthermore, seaweeds biochemical profile contains different sources of protein, fat, carbohydrate, etc., and nutritionally would be considered as functional ingredients [9]. Tables 1 and 2 exhibit some bioactive components of red algae as well as some of its biological properties.



Figure 1 . Red algae [10]

Table 1 . Bioactive compounds of several species of red algae

Bioactive molecules	Red algae	Virus targets (within the replication cycle)	Reference
Bromophenols	Polysiphonia morrowii	Fish pathogenic viruses, infectious hematopoietic necrosis virus and infectious pancreatic necrosis virus	[11]
carrageenan	Gigartina	Herpes simplex virus types 1 (HSV-1) and 2 (HSV-2)	[11]
Griffithsin	Griffithsia sp	GRFT Cellular intrusion of the HIV-1, Ebola, SARS, hepatitis C and H5N1 virus	[10, 12]
Sulphated galactans Galactose, xylose	Grateloupia filicina, Grateloupia longifolia	HIV-1 inhibitory activity	[13]
Sulfated glucuronogalactan Galactose	Schizymenia dubyi	HIV-1 inhibitory activity	[14]

Table 2 . Some of the biological activities of red algae

Biological properties	Reference
Antiulcer products and suspending agents in radiological preparation	[15]
terpenoids, phlorotannins, and steroids	[16, 17]
polysaccharides, vitamins, minerals, and fibers	[18, 19]
antibacterial	[20]
antiviral	[21]
antifungal	[22]
anticoagulant	[23]
antitumor	[24]
anti-inflammatory activities	[25]

The presence of antioxidants could be a possible reason for prevention of oxidative damage while the seaweeds are exposed to sunlight [26]. If we consider seaweeds as antioxidant sources, and simultaneously, protecting the human body from disrupting reactions, therefore, the hypothesis of potential defense against oxidation would be possible [9].

In fact, antioxidants prevent DNA damage or mutation with reduction of reactive oxygen and obstruction of free radicals [27]. Some methods to detect antioxidant activity include beta-carotene bleaching (BCB) assay, ferric reducing antioxidant power (FRAP) assay, DPPH free radical scavenging assay [28-30].

#### Antimicrobial effects

Antimicrobials refer to any kind of substances with ability of killing or inhibiting the growth of microorganisms like bacteria, yeasts, molds with

the least damage to the host cells. Since seaweeds are living in aquatic environment where microbiota is rich, it could be noted that they have accustomed to have some antimicrobial components [31]. Repeatedly, to avoid increasing resistance of pathogens to the conventional antimicrobials, we can consider them as proper alternatives [32]. A bulk of published documents have been assigned to pronounce the micro- and macro-algae as potential antimicrobial sources against pathogens. To date, many studies have focused on isolated bioactive natural compounds, and for developing and designing new drugs to control pathogens.

Both food industry and consumers achieve to awareness of having more natural antioxidants, and healthier foods [33]. In literature, several studies have revealed efficacy of algae derived bioactive compounds against gram (+) but not gram (-) bacteria. New antimicrobial substances derived from algae are promising to act against pathogens [34].

**Table 3.** Some antimicrobial components derived from red algae

Anti-HIV compounds	red seaweeds	[34]
polybromocatechol compounds effects on a HeLa cell	Neorhodomela aculeata	[35]
HSV-1 and HSV-2	Sargassum vulgare	[36]
antiviral activity	Stoechospermum marginatum	[37]
anti-HSV-1	Laminaria angustata	[38]

Recently, several studies on seaweeds led to extraction of two new prenylated para-xylenes from *Caulerpa racemos*, which acts against some pathogens like *Candida glabrata*, *Trichophyton rubrum*, and *Cryptococcus neoformans* [39, 40].

#### Antibacterial Compounds From Red Algae

Several species of red algae have been reported, and this review deals with some of them such as *Laurencia* spp., *Gracillaria* spp., *Acanthophora* spp and other rare species.

**Table 4.** Antimicrobial compounds derived from *Laurencia* (Rhodomelaceae, Ceramiales)

C15 acetogenin en-yne	antistaphylococcal	8-256 mg/mL	[41]
halogenated sesquiterpene alcohol	antiherbivore activity		[42]
5-acetoxypalisadin B	Antimicrobial activities		[43]
palisadin A	Antimicrobial activities		[43]
palisadin B	Antimicrobial activities		[43]
pannosanol	Antimicrobial activities	60 µg/disk against <i>Proteus mirabilis</i> and 100 µg/disk against <i>Chromobacterium violaceum</i> and <i>Vibrio cholera</i>	[43]
pannosane	Antimicrobial activities	60 µg/disk against <i>Chromobacterium violaceum</i>	[43]
3(Z) Chlorofucin	Antimicrobial activities	100 µg/disk against <i>Chromobacterium violaceum</i>	[43]
10-acetoxyangasiol	Antimicrobial activities	10-acetoxyangasiol against <i>Vibrio cholerae</i> at 100 µg/mL	[44]
1-methyl-2,3,5-tribromoindole	Antimicrobial activities	<i>Staphylococcus</i> sp. at MIC 300 µg/mL	[44]

#### *Acanthophora* spp.

Some of the bioactive compounds have derived from them are  $\beta$ -sitosterol, saringosterol,  $5\alpha$ -cholestane-3,6-dione, and cholest-4-ene-3-one [45, 46]. It has been reported that some of the sterols have cytotoxic and antimicrobial properties.

#### *Gracillaria* spp.

This genus contains more than 300 species. The main reason that make this group especial is a commercial product derived from them called

agar which is extensively used in food and cosmetic industries [47-50].

#### *Asparagopsis* spp.

Detected chemical components of this genus by chromatographic methods are octadec-9-enoic

acid 2,3-dihydroxypropyl ester, 9-octadecanoic acid, octadecanoic acid, and chlorobenzene.

Members of this genus have been revealed to be effective against pathogens like *Escherichia coli*, and *Pseudomonas aeruginosa*. GC-MS method

have been used to identify the antimicrobial compounds (bromoform, and dibromoacetic acid) of this genus [51].

### CONCLUSION

In conclusion, we could mention that red algae are an appropriate source of promising antimicrobial agents, which could even play a role in synthesis of novel substances. Technical and chemical methods could employ some templates for designing new antimicrobial products. Additionally, other science branches like, biology, chemical engineering, along with biomedical techniques are needed to align synthetic products as natural antimicrobial products.

### FUNDING

This work was partly supported by the Russian Foundation Basic Research (RFBR) under the project 15-29-05893.

### REFERENCES

- Cox, S., Abu-Ghannam, N. and Gupta, S. An assessment of the antioxidant and antimicrobial activity of six species of edible Irish seaweeds. *International Food Research Journal* 2010; 17: 205-220.
- Seenivasan R; Rekha M; Geetha S. Antibacterial Activity and Phytochemical Analysis of Selected Seaweeds from Mandapam Coast, India. *Journal of Applied Pharmaceutical Science*. 2012; 2(10): 159-169.
- Radhika D; Veerabahu C; Priya R. Antibacterial Activity Of Some Selected Seaweeds From The Gulf of Mannar Coast, South India *Asian J. Pharm. Clin. Res.*, 2012; 5(4), 89-90.
- Koyanagi S, Tanigawa N, Nakagawa H, Soeda S and Shimeno H. Oversulfation of fucoidan enhances its anti-angiogenic and antitumor activities. *Biochem Pharmacol* 2003; 65: 173-179.
- Dhargalkar, V. K. and Verlecar, X. N. 2009. Southern Ocean seaweeds: A resource for exploration in food and drugs. *Aquaculture* 287: 229-242.
- Manivannan, K., Thirumaran, G., Devi, G. K., Anantharaman, P. and Balasubramanian, T. 2009. Proximate Composition of Different Group of Seaweeds from Vedalai Coastal Waters (Gulf of Mannar): Southeast Coast of India. *Middle-East Journal of Scientific Research* 4(2): 72-77.
- McHugh, D. J. 2003. A Guide to Seaweed Industry. FAO Fisheries Technical Paper No. 441.
- Prabhasankar, P., Ganesan, P., Bhaskar, N., Hirose, A., Stephen, N., Gowda, L. R., Hosokawa, M. and Miyashita, K. 2009. Edible Japanese seaweed, wakame (*Undaria pinnatifida*) as an ingredient in pasta: Chemical, functional and structural evaluation. *Food Chemistry* 115(2): 501-508.
- Plaza, M., Cifuentes, A. and Ibáñez, E. 2008. In the search of new functional food ingredients from algae. *Trends in Food Science & Technology* 19(1): 31-39.
- Emau P, Tian B, O'keefe BR. et al. "Griffithsin, a potent HIV entry inhibitor, is an excellent candidate for anti-HIV microbicide". *J. Med. Primatol.* 2007; 36: 244-53.
- Kim SY, Kim SR, Oh MJ, Jung SJ, Kang SY. In vitro antiviral activity of red alga, *Polysiphonia morrowii* extract and its bromophenols against fish pathogenic infectious hematopoietic necrosis virus and infectious pancreatic necrosis virus. *Journal of Microbiology*. 2011; 49: 102-106.
- Philip M, Anna A, Sandrine B et al. Griffithsin Has Antiviral Activity against Hepatitis C Virus. *Antimicrob Agents Chemother.* 2011; 55: 5159-5167.
- Wang HB, Mo QH, Yang Z. (2015) HIV vaccine research: the challenge and the way forward. *J Immunol Res.*, 2015:503978.
- Bourgougnon N, Lahaye M, Quemener B, Chermann JC et al. Annual variation in composition and in vitro anti-HIV-1 activity of the sulfated glucuronogalactan from *Schizymenia dubyi* (Rhodophyta, Gigartinales). *Journal of Applied Phycology*. 1996; 8: 155-161.
- Rajasulochana P; Dharmotharan R; P Krishnamoorthy; S Murugesan. Antibacterial Activity of the Extracts of Marine Red and Brown Algae. *Journal of American Science*, 2009; 5(3): 20-25.
- Mtolera, M.S.P. and Semesi, AK. Antimicrobial activity of extracts from six green algae from Tanzania. In: *Current Trends in Marine Botanical Research in the East African Region* (eds M. Bjérk, A.K. Semesi, M. Pedersen and B. Bergman). Gotab AB, Uppsala, Sweden, 1996; pp. 211-217.
- Taskin, E., Ozturk, M., Taskin, E. and Kurt, O. Antibacterial Activities of Some Marine Algae from the Aegean Sea (Turkey). *African*

- Journal of Biotechnology, 2007; 6: 2746-2751.
18. Lahaye M. Marine algae as sources of fibres: Determination of soluble and insoluble dietary fibre contents in some 'sea vegetables'. *J. Sci. Food Agric.*, 1999; 54, 587-594.
  19. Darcy-Vrillon B. Nutritional aspects of the developing use of marine macroalgae for the human food industry. *Int. J. Food Sci. Nutr.* 1993; 44: 523-535.
  20. Burkholder PR., Burkholder LM., Almodovar LR. Antibiotic activity of some marine algae of Puerto Rico. *Bot. Mar.* 1960; 2: 149- 156.
  21. Richards JT; Kern ER; Glasgow LA; Overall JC; Deign EF; Melvin MT. Antiviral Activity of Extracts from Marine Algae. *Antimicrobial Agents Chemother.* 1978; 14, 24-30.
  22. Oumaskour K, Boujaber N, Etahiri S; Assobhei O. Screening of antibacterial and antifungal activities in green and brown algae from the coast of Sidi Bouzid (El Jadida, Morocco). *African Journal of Biotechnology*, 2012; 11(104): 16831-16837.
  23. Yasantha A; Ki-Wan L; Se-Kwon K; You-Jin J. Anticoagulant activity of marine green and brown algae collected from Jeju Island in Korea. *Bioresour. Technol.*, 2007; 98: 1711-1716.
  24. Abirami RG; Kowsalya S. Anticancer Activity of Methanolic and Aqueous Extract of *Ulva Fasciata* In Albino Mice. *International Journal of Pharmacy and Pharmaceutical Science*, 2012; 4(2): 681-684.
  25. Boonchum W; Peerapornpisal Y; Kanjanapathi D; Pekkoh J; Amornlerdpison D; Pumas C; Sangpaiboon P; Vacharapiyasophon P. Antimicrobial and anti-inflammatory properties of various seaweeds from the gulf of Thailand. *Int. J. Agric. Biol.*, 2011; 13: 100-104.
  26. Matsukawa, R., Dubinsky, Z., Kishimoto, E., Masaki, K., Masuda, Y., Takeuchi, T., Chihara, M., Yamamoto, Y., Niki, E. and Karube, I. 1997. A comparison of screening methods for antioxidant activity in seaweeds. *Journal of Applied Phycology* 9: 29-35.
  27. Khan, N., Afaq, F. and Mukhtar, H. 2008. Cancer chemoprevention through dietary antioxidants progress and promise. *Antioxidants & Redox Signaling* 10(3): 475-510.
  28. Chew, Y. L., Lim, Y. Y., Omar, M. and Khoo, K. S. 2008. Antioxidant activity of three edible seaweeds from two areas in South East Asia. *LWT* 41: 1067-1072.
  29. Manivannan, K., Thirumaran, G., Devi, G. K., Anantharaman, P. and Balasubramanian, T. 2009. Proximate Composition of Different Group of Seaweeds from Vedalai Coastal Waters (Gulf of Mannar): Southeast Coast of India. *Middle-East Journal of Scientific Research* 4(2): 72-77.
  30. Kumar, K. S., Ganesan, K. and Subba Rao, P. V. 2008. Antioxidant potential of solvent extracts of *Kappaphycus alvarezii* (Doty) Doty - An edible seaweed. *Food Chemistry* 107: 289-295.
  31. Shannon E, Abu-Ghannam N. Antibacterial derivatives of marine algae: An overview of pharmacological mechanisms and applications. *Marine Drugs*. 2016; 14(4):81.
  32. Falaise C, François C, Travers MA, Morga B, Haure J, Tremblay R, Leignel V. Antimicrobial Compounds from Eukaryotic Microalgae against Human Pathogens and Diseases in Aquaculture. *Marine drugs*. 2016; 14(9):159.
  33. Gyawali R, Ibrahim SA. Natural products as antimicrobial agents. *Food control*. 2014; 46:412-429.
  34. Bourgougnon N. Anti-HIV compounds from red seaweeds. *Biomaterials and bioprocessing*, Vol. 9. Recent advances in marine biotechnology. Science Publishers, Enfield. 2003; 165-206.
  35. Park SH, Song JH, Kim T, Shin WS, Park GM, Lee S, Kwon DH. Anti-human rhinoviral activity of polybromocatechol compounds isolated from the Rhodophyta, *Neorhodomela aculeata*. *Marine drugs*. 2012; 10(10):2222-2233.
  36. Plouguerné E, Ioannou E, Georgantea P, Vagias C, Roussis V, Hellio C, et al. Antimicrofouling activity of lipidic metabolites from the invasive brown alga *Sargassum muticum* (Yendo) Fensholt. *Marine Biotechnology*. 2010; 12(1):52-61.
  37. Adhikari U, Mateu CG, Chattopadhyay K, Pujol CA, Damonte EB, Ray B. Structure and antiviral activity of sulfated fucans from *Stoechospermum marginatum*. *Phytochemistry*. 2006; 67:2474-2482.
  38. Saha S, Navid MH, Bandyopadhyay SS, Schnitzler P, Ray B. Sulfated polysaccharides from *Laminaria angustata*: Structural features and in vitro antiviral activities. *Carbohydrate Polymers*. 2012; 87:123-130.

39. Kaaria PK, Matiru VN, Wakibia JG, Ndung'u M W, Bii CC. Evaluation of antibacterial and antifungal activity of macro algae associated microbes from the Northern and Southern coast of Kenya. *Journal of Microbiology and Biotechnology Research*. 2017; 5(5):7-13.
40. Liu AH, Liu DQ, Liang TJ, Yu XQ, Feng MT, Yao LG, et al. Caulerprenylols A and B, two rare antifungal prenylated para-xylenes from the green alga *Caulerpa racemosa*. *Bioorganic & Medicinal Chemistry Letters*. 2013; 23(9):2491-2494.
41. Kladi, M., Vagias, C., Stavri, M., M. Rahman, M.M., Gibbons, S., and Roussisa, V., 2008, *Phytochem. Lett.*, 1(1), 31-36.
42. Dos Santos A.O., Veiga-Santos P., Ueda-Nakamura, T., Filho, B.P., Sudatti, D.B., Bianco, E.M., Pereira, R.C., and Nakamura, C.V., 2010, *Mar. Drugs*, 8(11), 2733-2743.
43. Suzuki, M., Daitoh, M., Vairappan, C.S., Abe, T., and Masuda, M., 2001, *J. Nat. Prod.*, 64(5), 597-602.
44. Vairappan, C.S., Ishii, T., Lee, T.K., Suzuki, M., and Zhaoqi, Z., 2010, *Mar. Drugs*, 8(6), 1743-1749.
45. Wahidulla, S., D'Souza, L., and Govenker, M., 1998, *Phytochemistry*, 48(7), 1203-1206.
46. Lang, K.L., Palermo, J.A., Falkenberg, M., and Schenkel, E.P., 2007, *Biochem. Syst. Ecol.*, 35,805-808.
47. Prakash, O., Roy, R., and Bhakuni, D.S., 1989, *J.Nat. Prod.*, 52(4), 686-692.
48. Marinho-Sariano, E., 2001, *J. Biotechnol.*, 89(1),81-84.
49. Francavilla, M., Franchi, M., Monteleone, M., and Caroppo, C., 2013, *Mar. Drugs*, 11(10), 3754-3776.
50. Andriamanatoanina, H., Chambat, G., and Rinaudo, M., 2007, *Carbohydr. Polym.*, 68, 77-88.
51. Paul, N.A., de Nys, R., and Steinberg, P.D., 2006, *Mar. Ecol. Prog. Ser.*, 306, 87-101.