



Overview of the Helminth Fauna of Slow Worms of Genus *Anguis* (Reptilia, Anguidae) in the Western Palearctic

Alexander A. Kirillov^{1*}, Nadezhda Yu. Kirillova¹, Alexander B. Ruchin²

¹ Institute of Ecology of Volga basin of RAS, Togliatti, Russia.

² Joint Directorate of the Mordovia State Nature Reserve and National Park «Smolny», Saransk, Russia.

ABSTRACT

The overview summarizes data on helminths of slowworms of genus *Anguis* inhabiting the Western Palearctic. The data was collected using all available literature for 180-year period and original data of the authors. The data contain information on 13 countries where slowworms have been studied. 21 parasite species are listed (2 trematodes, 1 cestoda, 17 nematodes and 1 acanthocephalan), each with the location in the host, geographical localities, countries, prevalence, range of infection intensity and literature sources. Most of helminths are «random» slowworms' parasites. There are only 5 common parasites: *Entomelas entomelas*, *Oswaldocruzia filiformis*, *Oswaldocruzia lisnykiensis*, *Oxysomatium brevicaudatum* and *Neoxysomatium caucasicum*, all of them nematodes and with direct life cycle. *Entomelas entomelas*, *Oswaldocruzia filiformis* and *Oxysomatium brevicaudatum* are found in most examined legless lizards.

Keywords: Helminths, Slowworm, *Anguis*, Western Palearctic.

HOW TO CITE THIS ARTICLE: Alexander A. Kirillov, Nadezhda Yu. Kirillova, Alexander B. Ruchin; Overview of the Helminth Fauna of Slow Worms of Genus *Anguis* (Reptilia, Anguidae) in the Western Palearctic, Entomol Appl Sci Lett, 2019, 6 (1): 51-65.

Corresponding author: Alexander A. Kirillov

E-mail ✉ parasitolog@yandex.ru

Received: 02/11/2016

Accepted: 1/03/2018

INTRODUCTION

The latest molecular research [1-3] shows that genus *Anguis* includes 5 genera of legless lizards, inhabiting Palearctic: *A. fragilis* (LINNAEUS 1758) (Western and Central Europe), *A. colchica* (NORDMANN 1840) (Eastern Europe and Western Asia), *A. graeca* (BEDRIAGA 1881) (Southern Balkans), *A. cephalonica* (WERNER 1894) (The Peloponnese) and *A. veronensis* (POLLINI 1818) (Italy and Southeastern France). In some parts of species of this genus a range of hybridization is observed [4-6]. The northern limit of the spread of *Anguis* almost reaches the Arctic Circle; the eastern limit passes through Tobol River valley in Western Siberia [7-9]. *Anguis* species prefer to live in forest habitats. They use forest floor, rotten stumps, tree trunks, stones and animal burrows as their shelters. Their ratio consists of earthworms, slugs and other soil invertebrates [10-12]. The purpose of this work is to summa-

rize the existing data on helminths of the genus *Anguis* lizards inhabiting Western Palearctic.

MATERIALS AND METHODS

Our overview is based on all available literature sources and on our original data. A number of researchers have been studying genus *Anguis* (more than 220 specimens) in 12 European countries since 1845 till 2018 (Fig. 1). Our study is based on the analysis of helminth fauna of *Anguis* genus collected from several regions of Russia in 2000–2018. In our research, we examined only road-killed slowworms. Animal dissection, fixation and treatment of parasitological material were performed according to standard techniques [13, 14]. Our collected material consisted exclusively of nematodes. Collected parasites were killed by heating in water. Clarification of objects was carried out in lactic acid. The whole mount of nematodes was made by enclosing helminths in Glycerin-Jelly. Helminths were defined according to V.P. Sharpilo (1976) [15].

If the literature sources lacked quantitative indicators, presence of parasite in the region was marked with a "+" sign. Parasite taxonomy is provided according to the Fauna Europaea website (<http://www.fauna-eu.org/>). For each para-

site are given: the number of examined lizards, its locations in the host, localities of detection, country, parasite prevalence (P, % – proportion of infected reptiles), range of intensity of infection (I, specimens) and authors of record.




Fig. 1. A map of helminth research sites of slow worms in the Western Palearctic. Legend: stars on the map mark the catching places of lizards. 1 – Dujardin (1845); 2 – Jones *et al.* (2011); 3 – Baylis (1928); 4 – Markowski (1933); 5 – Moravec (1963); 6 – Borkovcova, Kopriva (2005); 7 – Bertman, Okulewicz (1984, 1987) and Lewin (1990); 8 – Muhling (1898); 9 – Markov (1952); 10 – Shimalov *et al.* (2000), Shimalov (2010); 11 – Svitin (2017); 12 – Shevchenko (1963); 13 – Mihalca *et al.* (2007); 14 – Schad *et al.* (1960) and Dusen *et al.* (2010); 15 – Sumer *et al.* (2019); 16 – Sharpilo (2003); 17 – Sharpilo (1962); 18 – Sharpilo (1974); 19 – Kirillov *et al.* (2015); 20 – Kirillov *et al.* (2018); 21 – Kirillov (2000). [16-43]

RESULTS AND DISCUSSION

The information on the parasitic worms of European slowworms is short and fractional. As a rule, data on *A. fragilis* parasites were obtained during the study of the overall helminth fauna of reptiles or all vertebrates in general. First information about helminth fauna of *Anguis* genus dates back to the mid-19th century, when F. DUJARDIN (1845) described a new species of nematodes – *Entomelas entomelas* [16]. P. MUHLING (1898) found the nematodes *Oswaldocruzia filiformis* (GOEZE 1782) and *Cosmocerca ornata* in the lizards of the Kaliningrad region of Russia (formerly, East Prussia) [17]. *Entomelas entomelas* and *Oxysomatium brevicaudatum* (ZEDER 1800) were noted in British slowworms by H.A. BAYLIS (1928) and Polish lizards by S. MARKOWSKI (1933) [18, 19]. Further studies of slowworms' helminths were carried out in Russia [26, 40, 41, 43], Ukraine

[15, 30, 31, 38, 44, 45], Georgia [15, 38, 44], Turkey [33, 35, 36], Czech Republic [20, 21] Poland [22-24], Great Britain [25], Belarus [27-29], Romania [32] and Armenia [46]. Works by V.P. Sharpilo (1974, 2003) are devoted to the endemic nematode species of the Caucasus fauna *Neoxysomatium caucasicum*, which first described from slowworm [37, 39].

In total, there are 21 species of parasites in representatives of genus *Anguis* in the Western Palearctic: 2 trematodes, 1 cestoda, 17 nematodes and 1 acanthocephalan (Table 1).

Phylum Plathelminthes Schneider, 1873

Class Cestoda Rudolphi, 1808

Family Mesocestoididae Perrier, 1897

Mesocestoides sp., tetrathyridium. Cestodes of the genus *Mesocestoides* usually parasitize at the larval stage as a rule in the body cavity of reptiles. Lizards can be intermediate or paratenic (reservoir) host of this parasite. The definitive

hosts of this parasite are birds of prey or mammals. The development cycle of mesocestoides involves 3 levels of hosts: first intermediate (apparently, arthropoda), second intermediate (amphibians, reptiles or rodents) and definitive (birds of prey or mammals) [47, 48].

Other reported reptilian hosts: *Lacerta agilis* [49, 50]; *Lacerta schreiberi* [51]; *Lacerta trilineata* [52]; *Lacerta viridis* [53]; *Apathya cappacocica* [54]; *Phoenicolacerta laevis* [55]; *Anatololacerta danfordi* [56]; *Phrynocephalus mystaceus* [57]; *Podarcis bocagei* [58]; *Podarcis hispanicus* [58]; *Podarcis muralis* [59]; *Podarcis pityusensis* [60]; *Psammodromus hispanicus* [61, 62]; *Eremias arguta* [63]; *Eremias velox* [63]; *Coronella austriaca* [64]; *Zamenis longissimus* [65]; *Psammophis lineolatus* [63].

Class Trematoda Rudolphi, 1808

Family Brachylaimidae Joyeux et Foley, 1930

Brachylaima sp. J. LEWIN (1990), according to the figure, noted a young parasite with an undeveloped reproductive system, which may be a consequence of the parasite's development in a non-specific host. All other authors recorded *Brachylaima* sp., metacercariae, in the slowworm and other reptile species. Finding *Brachylaima* sp. in legless lizards should be considered accidental. Terrestrial gastropods are intermediate hosts of trematodes of genus *Brachylaima*, while birds and mammals are the definitive hosts [66]. Parasite infestation occurs when lizards feed on land mollusks, which are intermediate hosts of these trematodes. Apparently, reptiles occur to be abortive hosts for *Brachylaima* spp.

Other reported reptilian hosts: *Lacerta agilis* [49]; *Lacerta trilineata* [52]; *Lacerta strigata* [15, 38]; *Podarcis bocagei* [51]; *Podarcis carbonelli* [67]; *Podarcis lilfordi* [60, 68]; *Podarcis pityusensis* [68], *Chalcides ocellatus* [69].

Family Pleurogenidae Looss, 1899

Pleurogenoides medians (OLSSON 1876). It is a common parasite of anurans and caudates. The first intermediate hosts of the trematode are gastropods, the second intermediate hosts are the larvae and adult insects, that live near water (dragonflies, caddisflies, mayflies, alderflies, dipterans), and crustaceans (*Asellus aquaticus*, amphipods) [70]. The invasion of the slowworms

by *Pleurogenoides medians*, apparently, happened accidentally by eating additional hosts of the trematode.

Other reported reptilian hosts: *Lacerta agilis* [15, 20, 49, 50]; *Lacerta trilineata* [52]; *Zootoca vivipara* [15]; *Podarcis hispanica* [62, 71].

Phylum Acanthocephala (Rudolphi, 1808)

Class Palaeacanthocephala (Meyer, 1931)

Family Echinorhynchidae Cobbold, 1876

Acanthocephalus ranae (SCHRANK 1788). It is a common parasite of amphibians, mainly anurans. Crustaceans, such as *Asellus aquaticus* and wood lice [72, 73] serve as intermediate hosts of the genus *Acanthocephalus*. Amphibians become infected by accidental swallowing the intermediate hosts of the parasite.

Other reported reptilian hosts: *Natrix natrix* [27, 31, 74].

Phylum Nematoda Potts, 1932

Class Secernentea Chitwood, 1958

Family Rhabdiasidae Railliet 1916

Entomelas entomelas (Dujardin, 1845). A common parasite of legless lizards of genus *Anguis* and *P. apodus*. As a rule, studies indicate 2 species of the genus *Entomelas* TRAVASSOS 1930: *Entomelas dujardini* (МАУРАС 1916) and *Entomelas entomelas* [75]. M.R. BAKER (1980) changed description of *Entomelas* genus and synonymized *Entomelas dujardini* with *E. entomelas*. Life style of the parasite is characterized by heterogony. Invasive larvae are obtained from the free-living generation. Infection of reptiles with invasive larvae occurs both directly from the external environment and through food objects (mollusks and earthworms), which are paratenic hosts of nematodes [76, 77].

Other reported reptilian hosts: *Pseudopus apodus* [78].

Rhabdias bufonis (SCHRANK 1788). It is a common parasite of amphibians. The development of nematodes of the family Rhabdiasidae occurs without the participation of intermediate hosts and is characterized by the alternation of free-living and parasitic generations. In most species of rhabdiasids, invasive larvae develop in the free-living generation [77]. Infection of amphibians occurs through percutaneous penetration of invasive parasite larvae from the soil.

No other reported reptilian hosts.

***Rhabdias fuscovenosa* (RAILLIET 1899).** It is a common parasite of colubrid snakes. Invasive larvae infect snakes orally. Unlike nematodes of amphibians, invasive larvae from *Rhabdias* species, that parasitize on reptiles, develop from parasitic eggs [79]. Eggs, which are laid by female parasites, enter the digestive tract through the trachea and excrete outside, where larvae reach the invasive stage [80]. Reptile infection with nematodes happens orally, so paratenic hosts may be involved [77, 80].

Other reported reptilian hosts: *Lacerta viridis* [21]; *Natrix natrix* [15, 20, 31, 32, 40, 42, 43, 63, 78, 81-85]; *Natrix tessellata* [15, 20, 32, 38, 42, 43, 63, 81, 84, 85]; *Coronella austriaca* [15], *Elaphe dione* [15], *Zamenis longissimus* [34]; *Eryx tataricus* [15], *Vipera ursinii* [15], *Vipera berus* [15, 32, 40, 86].

***Rhabdias* sp.** Nematodes from slowworms in Ukraine were not identified to the species level [31].

Other reported reptilian hosts: *Coronella austriaca* [31].

Family Rhabditidae Oerley 1880

Rhabditidae sp. Nematodes of this family are common parasites among insects and mollusks. According to R. JONES ET AL. (2011), the finding of this parasite in the slowworms' faeces may indicate post-cyclic parasitism (the parasite lives in the slowworm's body because of its food objects - soil invertebrates) [25]. We assume that the record of data on nematodes as an *Anguis*' parasite is a mistake. Other researchers weren't able to detect this parasite during dissections of slowworms.

No other reported reptilian hosts.

Family Molineidae Skrjabin et Schulz, 1937

***Oswaldocruzia filiformis* (GOEZE 1782).** It is a common parasite of Palearctic amphibians and reptiles. The development of the parasite occurs without the participation of intermediate hosts. Invasion of the hosts happens orally through the contact with invasive parasite larvae on land [87].

Apparently, all information about *Oswaldocruzia filiformis* in reptiles must be reviewed. In recent

studies, new species were described and identification key for genus *Oswaldocruzia* Travassos 1917 was created. Three species of this genus were also reviewed: *Oswaldocruzia filiformis* (GOEZE 1782), *Oswaldocruzia bialata* (MOLIN 1860) and *Oswaldocruzia skrjabini* Travassos, 1937 [87-95]. R.S. SVITIN (2017) described new species for reptiles: *Oswaldocruzia lacertica* (sand lizard) and *Oswaldocruzia lisnykiensis* (*Anguis*) [30]. Still, mostly, parasitological studies of amphibians and reptiles name *Oswaldocruzia filiformis* species.

SVITIN and GOROBCHISHIN (2015) assume that all *Oswaldocruzia* spp. in snakes should be considered to be cases of post-cyclic parasitism [96].

Other reported reptilian hosts: *Lacerta agilis* [15, 20, 28, 32, 40, 42, 43, 50, 65, 86, 97]; *Lacerta viridis* [20, 21, 33, 53, 59, 98]; *Podarcis tauricus* [33]; *Darevskia rudis* [99]; *Pseudopus apodus* [15]; *Takydromus amurensis* [15]; *Zootoca vivipara* [20, 26, 28, 40, 43, 100]; *Coronella austriaca* [15, 31]; *Natrix natrix* [20, 23, 27, 31, 49, 63, 64, 96, 101]; *Natrix tessellata* [15, 63]; *Telescopus fallax* [15]; *Vipera ammodytes* [15]; *Vipera berus* [20, 26, 27, 40, 43, 96, 102, 103].

***Oswaldocruzia lisnykiensis* Svitin 2017.** It is a common parasite of lizards of genus *Anguis* [30]. V.P. SHARPILO (1976) thinks that the phenomenon of host-specificity among the parasitic worms of reptiles is extremely rare. Apparently, as other *Oswaldocruzia* spp., they have direct life cycle without the participation of intermediate hosts.

No other reported reptilian hosts.

Family Cosmocercidae Railliet, 1916

***Oxysomatium brevicaudatum* (ZEDER 1800).** It is a common parasite of amphibians and, less commonly, reptiles. It seems that, as other cosmocercid species, they have direct development cycle, without the participation of intermediate hosts. Eggs develop in the environment. Infection of hosts with invasive larvae most likely happens orally.

Other reported reptilian hosts: *Natrix natrix* [15, 33]; *Vipera berus* [15]; *Zamenis longissimus* [34].

***Oxysomatium* sp.** Nematodes of the genus *Oxysomatium* (not identified to the species) were

found in only one of the examined slowworms in Armenia [46].

No other reported reptilian hosts.

***Neoxysomatium caucasicum* Sharpilo, 1974.** It is a common parasite of lizards of genus *Anguis*. Life cycle has not been studied, but should be direct. Infection of slowworms with invasive larvae happens orally from the environment.

No other reported reptilian hosts.

***Cosmocerca ornata* (Dujardin, 1845).** It is a common parasite of amphibians. As other nematodes of *Cosmocercidae* family, they have direct development cycle, without the participation of intermediate hosts [77].

Our studies have shown that invasive larvae (L3) penetrate the amphibians through the eyes, in the near-surface layer of water. On the conjunctiva of the lower eyelid, they survive third and fourth moltings. Young nematodes migrate to third part of posterior intestine through the oropharyngeal cavity and gastrointestinal tract [104]. Therefore, we consider the finding of *Cosmocerca ornata* in slowworms to be a mistake. The researchers probably dealt with the *Cosmocerca commutata* (DIESING 1851), a parasite of terrestrial amphibians.

No other reported reptilian hosts.

***Aplectana acuminata* (SCHRANK 1788).** A common parasite of anurans. It seems that, as other cosmocercid species, they have direct life cycle, without the participation of intermediate hosts. Infection of hosts with invasive larvae happens orally from the environment.

Other reported reptilian hosts: *Natrix natrix* [15].

Family Protostrongylidae Leiper, 1926

Protostrongylidae sp. Nematodes of *Protostrongylidae* family have an indirect life cycle. The development of parasites requires participation of intermediate hosts, terrestrial molluscs, in which larvae become invasive [105, 106]. Infection of slowworms with protostrongylids occurs by consuming molluscs.

No other reported reptilian hosts.

Family Spirocercidae Chitwood et Wehr, 1932

***Ascarops strongylina* (RUDOLPHI 1819), juv.** It is a widespread parasite of pigs, boars and live-stock [15]. The parasite has an indirect life cycle. Nematode eggs fall into the environment with host's faeces. Coprophagous beetles are intermediate hosts of the parasite. After that, the larvae become infectious, and infect pigs when they ingest infected dung beetles [107]. Reptiles serve as paratenic hosts of the parasite.

Other reported reptilian hosts: *Lacerta agilis* [15, 27, 50]; *Zootoca vivipara* [15, 28]; *Dolichophis jugularis* [108]; *Platyceps najadum* [108]; *Dolichophis caspius* [15]; *Coronella austriaca* [27]; *Gloydus halys* [109]; *Malpolon monspessulanus* [108]; *Natrix tessellata* [15], *Natrix natrix* [15, 27]; *Coronella austriaca* [15], *Vipera aspis* [110]; *Vipera berus* [15, 27]; *Vipera latastei* [110]; *Emys orbicularis* [15].

***Physocephalus sexalatus* (Molin, 1860), juv.** It is a widespread parasite of domestic pigs and boars [77]. The life cycle is the same with that of *Ascarops strongylina*. At larval stage, it is a widespread parasite of reptiles, birds, insectivores, bats and rodents, which serve as paratenic hosts of the parasite. Infection of reptiles with this parasite occurs after eating intermediate hosts of the parasite – coprophagous beetles [107].

Other reported reptilian hosts: *Laudakia caucasica* [15, 38], *Laudakia erythrogastra* [109], *Laudakia lehmanni* [109], *Phrynocephalus mystaceus* [109], *Pseudopus apodus* [15], *Eremias velox* [15, 38], *Lacerta strigata* [15, 38], *Zootoca vivipara* [15], *Lacerta agilis* [15, 38], *Lacerta saxicola* [15, 38], *Telescopus fallax* [15], *Natrix natrix* [15], *Natrix tessellata* [15], *Podarcis tauricus* [15], *Dolichophis caspius* [15, 38], *Platyceps najadum* [15], *Zamenis longissimus* [15], *Coronella austriaca* [15], *Vipera berus* [15], *Emys orbicularis* [15], *Agrionemys horsfieldii* [15].

Family Physalopteridae Railliet, 1893

***Abbreviata* sp.** Nematodes of genus *Abbreviata* Travassos (1919) parasitize the digestive tract of reptiles. These parasites have an indirect life cycle. The intermediate hosts of physalopterids are arthropods [77]. Infection of reptiles occurs by arthropods consumption.

Other reported reptilian hosts: *Lacerta agilis* [21]; *Lacerta viridis* [21]; *Coronella austriaca* [21].

Unidentified larvae of suborder Spirurina Railliet et Henry, 1915

Agamospirura minuta (SHARPILO 1963), juv. Intermediate hosts, arthropods [77], are involved in the life cycle of spiruride nematodes. Reptiles are paratenic hosts. J. LEWIN (1990) considers erroneous defining of *Agamospirura* larvae, found in slowworms by V.P. SHARPILO (1964, 1976), as Spirurina. J. Lewin thinks that the larvae of *Protostrongylidae* sp. are similar to *Agamospirura* spp., which was found in reptiles by V.P. Sharpilo.

Other reported reptilian hosts: *Lacerta viridis*, *Coronella austriaca*, *Natrix natrix*, *Natrix tessellata*, *Vipera berus* [15, 38, 44].

Agamospirura sp., juv. Larvae of spirurian nematodes, found in slowworms in Ukraine, were not identified to the species level [31].

Other reported reptilian hosts: *Lacerta agilis* [31], *Natrix natrix* [31]; *Coronella austriaca* [31], *Vipera berus* [31].

Out of 21 parasites that were found in legless lizards, the majority of helminths are «random» parasites of slowworms (10 species): *Brachylaima* sp., *Pleurogenoides medians*, *Acanthocephalus ranae*, *Cosmocerca ornata*, *Aplectana acuminata*, *Rhabdias bufonis*, *Rhabdias fuscovenosa*, *Rhabdias* sp., *Abbreviata* sp., and *Rhabditidae* sp. These helminth species are common parasites of other vertebrates (mainly, amphibians and snakes).

Most of parasites (15 species) of genus *Anguis* were discovered in adult form. Six species of helminthes parasitize legless lizards in larval stage. The finding of larval form of parasites in slowworms involves that they are paratenic hosts of these helminthes.

Relatively high rates of invasion were noted in slowworms with such parasites: *Neoxysomatium caucasicum*, *Oxysomatium* sp., *Cosmocerca ornata*, *Aplectana acuminata*, *Physocephalus sexalatus*, juv., *Rhabdias fuscovenosa*, *Rhabdias bufonis*, *Rhabdias* sp., *Rhabditidae* sp., *Protostrongylidae* sp., juv. Low rates of invasion and occasional discoveries were recorded in slow-

worms with such parasites: *Mesocestoides* sp., tetrathyridium, *Brachylaima* sp., *Pleurogenoides medians*, *Acanthocephalus ranae*, *Abbreviata* sp., *Ascarops strongylina*, juv., *Agamospirura minuta*, juv., *Agamospirura* sp., juv.

Nematodes predominate in helminth fauna of slowworms. Cestodes, trematodes, and acanthocephalans are rarely found in legless lizards. Helminth fauna of lizards of the genus *Anguis* characterizes the ecology of these reptiles. Infection of legless lizards with helminths happens, firstly, because of slowworms living in a damp forest floor (favorable environment for the development of nematodes' eggs and larvae with a direct development cycle), and secondly, because of lizards feeding on terrestrial invertebrates, which are intermediate and/or paratenic hosts of parasites.

The parasites with a direct life cycle prevail in helminth fauna of slowworms (12 species). The remaining 9 parasitic species are characterized by an indirect life cycle with intermediate hosts. Helminths of slowworms are better studied in Ukraine (9 recorded species), Belarus (8), Russia and Turkey (7 in each) and Poland (6). Helminth fauna of legless lizards in Czech Republic (5), Georgia (4), and Great Britain (3) is proven to be less diverse. Only one species was found in slowworms in Spain, Armenia, Azerbaijan, Romania and France. That is probably due to the small number of examined legless lizards.

There are only 5 common parasites for the lizards of genus *Anguis*: *Entomelas entomelas*, *Oswaldocruzia filiformis*, *Oswaldocruzia lisnykiensis*, *Oxysomatium brevicaudatum* and *Neoxysomatium caucasicum*. All of them are nematodes with a direct life cycle. *Entomelas entomelas*, *Oswaldocruzia filiformis* and *Oxysomatium brevicaudatum* are found in lizards in most studies. Most of the helminth species recorded in legless lizards are parasites of other vertebrates (mainly amphibians and snakes).

Overview has shown the need for further helminthological studies of slowworms in Palearctic, since these studies should expand list of parasite-host for legless lizards.

Table 1. Checklist of parasites of slow worm *Anguis fragilis*

Parasite	Infection site	Locality	Country	n	P (%)	I	Author of record
<i>Mesocestoides</i> sp., tetrathyridium	sw	Bieszczady Mountains	Poland	44	2,3	1	[24]
<i>Brachylaima</i> sp.	i	Bieszczady Mountains	Poland	44	4,6	1	[24]
		Trabzon, Bursa provinces	Turkey	15	6,7	1	[36]
<i>Pleurogenoides medians</i>	i	Kharkov region	Ukraine	17	5,9	25	[31]
<i>Acanthocephalus ranae</i>	i	Brest and Gomel regions	Belarus	19	21,1	1-3	[28]
<i>Entomelas entomelas</i>	bc	Pomerania	Poland	4	-	1-6	[19]
	l, p	Leningrad region	Russia	20	70,0	1-25	[26]
	l, p	-	Turkey	9	66,7	-	[33]
	l	Lagodekhi	Georgia	2	1 of 2	4	[38]
	l	Bohemia (Sumava)	Czech Republic	2	1 of 2	3	[20]
	o, p	Moravia	Czech Republic	10	90,0	1-50	[20]
	o, p	Kharkov region	Ukraine	17	52,9	1-17	[31]
	l	Bieszczady Mountains	Poland	1	+	8	[22]
	l, o, p, nc	Bieszczady Mountains	Poland	44	92,3	1-22	[24]
	l, o	-	Spain		+		[111]
	l, o, p	Samarskaya Luka	Russia	2	100	1-6	[43]
		Brest and Gomel regions	Belarus	30	36,7	1-10	[28; 29]
	-	-	Romania	2	1 of 2	3	[32]
	l	alikesir, Bursa, Duzce Sakarya provinces	Turkey	18	44,4	1-9	[35]
	l, o, p	Mordovia Nature Reserve	Russia	3	3 of 3	2-15	[40]
	l, o, p	«Smolny» National Park, Mordovia	Russia	24	91,7	1-16	[41]
l, o	Trabzon, Bursa provinces	Turkey	15	93,3	1-15	[36]	
<i>Rhabdias bufonis</i>	l	alikesir, Bursa, Duzce Sakarya provinces	Turkey	18	44,4	1-9	[35]
<i>Rhabdias fuscovenosa</i>	l	Southern Moravia	Czech Republic	7	28,6	>350	[21]
<i>Rhabdias</i> sp.	l	Kharkov region	Ukraine	17	64,7	1-10	[31]
Rhabditidae sp.	i	Wales	UK	100	83,0	1-2000*	[25]
<i>Oswaldocruzia filiformis</i>	i	-	Turkey	9	11,1	-	[33]
		Lagodekhi	Georgia	2	2 of 2	3-8	[33]
		Bohemia (Sumava)	Czech Republic	2	1 of 2	1	[20]
		Moravia	Czech Republic	10	20,0	1-6	[20]
		Kharkov region	Ukraine	17	52,9	1-10	[31]
		Krasnodar region	Russia	4	75,0	1-2	[112]
		Bieszczady Mountains	Poland	2	2 of 2	9-12	[23]
	d, s	Bieszczady Moun-	Poland	44	89,7	1-53	[24]

	i	tains					
		Samarskaya Luka	Russia	2	1 of 2	11	[43]
		Brest and Gomel regions	Belarus	19	10,5	1	[28]
		South Moravia	Czech Republic	7	14,3	2	[21]
		alikesir, Bursa, Duzce Sakarya provinces	Turkey	18	22,2	1-14	[35]
		Mordovia Nature Reserve	Russia	3	3 of 3	1-3	[40]
		«Smolny» National Park, Mordovia	Russia	24	20,8	1-2	[41]
<i>Oswaldocruzia lisnyiensis</i>	i	Kyiv, Chernivtsi, Volyn, Sumy, Lviv, Khmelnytskyi regions	Ukraine			+	[30]
<i>Oxysomatium brevicaudatum</i>	i	Pomerania	Poland	4	100	1-6	[19]
		-	Turkey	9	11,1	-	[33]
		Bohemia (Sumava)	Czech Republic	2	2 of 2	5-18	[20]
		Moravia	Czech Republic	10	90,0	1-67	[20]
		Kharkov region	Ukraine	17	58,8	1-7	[31]
	c	Bieszczady Mountains	Poland	44	61,5	1-42	[24]
	i	alikesir, Bursa, Duzce Sakarya provinces	Turkey	18	83,3	1-8	[35]
		Southern Moravia	Czech Republic	7	42,9	3-10	[21]
		Brest and Gomel regions	Belarus	30	20,0	1-8	[28; 29]
		Samarskaya Luka	Russia	2	2 of 2	8-27	[43]
		«Smolny» National Park, Mordovia	Russia	24	100,0	1-41	[41]
Trabzon, Bursa provinces		Turkey	15	93,3	1-60	[36]	
		Wales	UK	100	38,0	1-686*	[25]
<i>Oxysomatium</i> sp.	i	-	Armenia	1	+	14	[45]
<i>Neoxysomatium caucasicum</i>	i	Lagodekhi	Georgia	2	2 of 2	10-13	[33; 37]
		Krasnodar region	Russia	4	75,0	28-29	[112]
		Suhumi, Chiatura	Georgia			+	[37]
		Lankaran and Astaran regions	Azerbaijan			+	[37]
		Dagestan	Russia	12	-	10-35	[39]
<i>Cosmocerca ornata</i>	i	Kharkov region	Ukraine	17	5,9	1	[31]
		Brest and Gomel regions	Belarus	30	3,3	1	[28; 29]
		alikesir, Bursa, Duzce Sakarya provinces	Turkey	18	50,0	1-6	[35]
<i>Aplectana acuminata</i>	i	Leningrad region	Russia	20	95,0	1-43	[26]
		Kharkov region	Ukraine	17	5,9	2	[31]

Protostrongylidae sp., juv.	hp, sw, dw	Bieszczady Mountains	Poland	44	9,1	1–25	[24]
<i>Ascarops strongylina</i> , juv.	–	Brest and Gomel regions	Belarus	30	3,3	15	[28; 29]
<i>Physocephalus sexalatus</i> , juv.	–	Brest and Gomel regions	Belarus	19	21,1	4–112	[28]
<i>Abbreviata</i> sp.	–	Southern Moravia	Czech Republic	7	28,6	1–3	[21]
<i>Agamospirura minuta</i> , juv.	iw	Volyn, Khmelnytskyi regions	Ukraine	9	16,3	1–8	[44]
		Lagodekhi	Georgia	2	2 of 2	1–8	[15]
		Brest and Gomel regions	Belarus	19	5,3	1	[28]
<i>Agamospirura</i> sp., juv.		Kharkov region	Ukraine	17	11,8	2–3	[31]

* – Jones *et al.* (2011) examined faecal samples of slow worms

bc – body cavity; c – colon; d – duodenum; dw – duodenum wall; hp – hepatic parenchyma; i – intestine; iw – intestine wall; l – lung; nc – nasal cavity; o – oesophagus; p – pharynx; s – stomach; sw – stomach wall; juv. – third-stage juvenile

REFERENCES

- Gvozdik V., Jandzik D., Lymberakis P., Jablonski D., Moravec J. 2010. Slow worm *Anguis fragilis* (Reptilia: Anguidae) as a species complex: Genetic structure reveals deep divergences. *Molecular Phylogenetics and Evolution*. Vol. 55. Issue 2. P. 460–472.
- Jablonski, D., Jandzik, D., Mikulicek, P., Dzukic, G., Ljubisavljevic, K., Tzankov, N., Jelic, D., Thanou, E., Moravec, J., Gvozdik, V. 2016. Contrasting evolutionary histories of the legless lizards slow worms (*Anguis*) shaped by the topography of the Balkan Peninsula. *BMC Evolutionary Biology* 16: 99. DOI: 10.1186/s12862-016-0669-1.
- Renet, J., Lucente, D., Delaugerre, M., Gerriet, O., Deso, G., Abbat-Tista, C., Cimmaruta, R. 2018. Discovery of an Italian slow worm (*Anguis veronensis* Pollini, 1818) population on a Western Mediterranean Island confirmed by genetic analysis. *Acta Herpetologica*. 13(2): 165-169. DOI: 10.13128/Acta_Herpetol-23290.
- Gvozdik V., Benkovsky N., Crottini A., Bellati A., Moravec J., Romano A., Sacchi R., Jandzik D. 2013. An ancient lineage of slow worms, genus *Anguis* (Squamata: Anguidae), survived in the Italian Peninsula. *Molecular Phylogenetics and Evolution*. Vol. 69. Issue 3. P. 1077–1092.
- Koppitz, C. 2018. Are *Anguis graeca* (Bedriaga, 1881) and *Anguis cephallonica* Werner, 1894, sympatric in the lowlands of southern Laconia? *Herpetozoa* 30(3-4): 223-225.
- Psonis, N., Antoniou, A., Karameta, E., Leache, AD, Kotsakiozi, P., Darriba, D., Kozlov, A., Stamatakis, A., Poursanidis, D., Kukushkin, O., Jablonski, D., Crnobrnja-Isailovic, J., Gherghel, I., Lymberakis, P., Poulakakis, N. 2018. Resolving complex phylogeographic patterns in the Balkan Peninsula using closely related wall-lizard species as a model system. *Molecular Phylogenetics and Evolution* 125: 100-115. DOI: 10.1016/j.ympev.2018.03.021
- Beebee T., Griffiths R. 2000. *Amphibians and reptiles: A natural history of the British herpetofauna*. London: Harper Collins Publish. pp. 272.
- Szabo K, Voros J. 2014. Distribution and hybridization of *Anguis fragilis* and *A. colchica* in Hungary. *Amphibia-Reptilia* 35(1): 135-140. DOI: 10.1163/15685381-00002927
- Jablonski D, Najbar B, Grochowalska R, Gvozdik V, Strzala T. 2017. Phylogeography and postglacial colonization of Central Europe by *Anguis fragilis* and *Anguis colchica*. *Amphibia-Reptilia* 38(4): 562-569. DOI: 10.1163/15685381-00003133
- Pedersen, IL, Jensen, JK, Toft, S. 2009. A method of obtaining dietary data for slow worms (*Anguis fragilis*) by means of non-harmful cooling and results from a Danish population. *Journal of Natural History*

- 43(15-16): 1011-1025. DOI: 10.1080/00222930802702464
11. Fish, A.C.M. 2016. Observations on felt and corrugated roof sheeting as materials for constructing coverboards to assess slow worm (*Anguis fragilis*) and common lizard (*Zootoca vivipara*) populations. *Herpetological Bulletin* 135: 4-6.
 12. Tuniyev B.S., Lotiev K.Yu., Tuniyev S.B., Gabaev V.N., Kidov A.A. 2017. Amphibians and reptiles of South Ossetia. *Nature Conservation Research* 2(2): 1-23. <http://dx.doi.org/0.24189/ncr.2017.002>.
 13. Bykhovskaya-Pavlovskaya I.E. 1985. *Fish Parasites: A Guide to Study*. Leningrad: Nauka. 121 p. (in Russian)
 14. Anikanova V.S., Bugmyrin S.V. & Ieshko E.P. 2007. *Methods for collection and study of helminths in small mammals*. Petrozavodsk: Karelian Research Centre of RAS. 145 p. (in Russian)
 15. Sharpilo V.P. 1976. *Parasitic worms of the reptilian fauna of the USSR*. Kyiv: Naukova Dumka. 376 p. (in Russian).
 16. Dujardin F. 1845. *Histoire naturelle des helminthes ou vers intestinaux*. Paris: Librairie Encyclopédique de Roret. 645 pp.
 17. Muhling P. 1898. *Die Helminthenfauna der Wirbeltiere Ostpreussens*. *Archiv für Naturgeschichte*. 1: 1-118.
 18. Baylis H.A. 1928. Records of some parasite worms from British vertebrates. *Annals and Magazine of Natural History*. Ser. 10(1). P. 329-343.
 19. Markowski S. 1933. Contributions a l'étude de la fauna helminthologique de la presqu'île de Hel. *Fragmenta Faunistica Musei Zoologici Polonici*. 2: 107-111.
 20. Moravec F. 1963. Prispěvek k poznání helmintofauny našich plazů. *Spisy přírodovědecké fakulty J. E. Purkyně v Brně*. 19: 353-388.
 21. Borkovcova M., Kopriva J. 2005. Parasitic helminths of reptiles (Reptilia) in South Moravia (Czech Republic). *Parasitology Research*. 95 (1): 77-78.
 22. Bertman M., Okulewicz A. 1984. *Entomelas dujardini* (Maupas, 1916) (Nematoda) u padalca – *Anguis fragilis* L. *Wiadomości Parazytologiczne*. 30: 341-343. (In Polish)
 23. Bertman M., Okulewicz A. 1987. *Padalec – Anguis fragilis* L. i zaskroniec zwyczajny – *Natrix natrix* (L.) nowymi krajowymi żywicielami *Oswaldocruzia filiformis* (Goeze, 1782) Travassos 1917 (Nematoda). *Wiadomości Parazytologiczne*. 33: 209-212. (In Polish)
 24. Lewin J. 1990. Parasitic worms in a slow-worm (*Anguis fragilis* L.) population from the Bieszczady Mountains (Poland). *Acta Parasitologica Polonica*. 35(3): 207-215.
 25. Jones R., Brown D.S., Harris E., Jones J., Symondson W.O.C., Bruford M.W., Cable J. 2012. First record of *Neoxysomatum brevicaudatum* through the non-invasive sampling of *Anguis fragilis*: complementary morphological and molecular detection. *Journal of Helminthology*. 86: 125-129.
 26. Markov G.S. 1952. Parasite fauna of reptiles of the Leningrad region. *Scientific papers of the Leningrad State University* 28(141): 217-229 (in Russian).
 27. Shimalov V.V., Shimalov V.T. 2000. Helminth fauna of snakes (Reptilia, Serpentes) in Belorussian Polesye. *Parasitology Research*. 86: 340-341.
 28. Shimalov V.V., Shimalov V.T., Shimalov A.V. 2000. Helminth fauna of lizards (Reptilia, Sauria) in the southern part of Belarus. *Parasitology Research*. 86: 343.
 29. Shimalov V. V. 2010. Helminth fauna of reptiles in the Republic of Belarus. *Parazitologiya*. 44 (1): 22-29 (in Russian).
 30. Svitin R. 2017. Two new species of *Oswaldocruzia* (Nematoda, Molineidae) parasitising lizards in Ukraine. *Zootaxa*. 4263 (2): 358-368.
 31. Shevchenko N.N. 1963. Helminth fauna of reptiles of the Seversky Donets Valley. Helminths of humans, animals and plants and ways to fight them. Moscow: Nauka, pp. 112-114 (in Russian).
 32. Mihalca A.D., Gherman C., Ghira I., Cozma V. 2007. Helminth parasites of reptiles (Reptilia) in Romania. *Parasitology Research*. 101: 491-492. DOI: 10.1007/s00436-007-0486-y
 33. Schad G.A., Kuntz R.E., Wells W.H. 1960. Nematode parasites from Turkish vertebrates: an annotated list. *Canadian Journal of Zoology*. 38: 949-963.

34. Dusen S., Ugurtas I.H., Altunel F.N. 2010a. Nematode parasites of the smooth snake, *Coronella austriaca* Laurenti, 1768 and the aesculapian snake, *Zamenis longissimus* (Laurenti, 1768) (Ophidia: Colubridae), collected from Northwestern Turkey. *North-Western Journal of Zoology*. 6 (1): 86–89.
35. Dusen S., Ugurtas I.H., Aydogdu A. 2010b. Nematode parasites of the two limbless lizards: Turkish worm lizard, *Blanus trauchi* (Bedriaga, 1884) (Squamata: Amphisbaenidae), and slow worm, *Anguis fragilis* Linnaeus 1758 (Squamata: Anguidae), from Turkey. *Helminthologia*. 47(3): 158–163.
36. Sumer N., Birlik S., Yildirimhan H.S. 2019. Morphological and molecular taxonomy of helminths of the slow worm *Anguis fragilis* (Linnaeus) (Squamata: Anguidae) from Turkey. *Biharean Biologist*. Vol. 13 – in press.
37. Sharpilo V.P. 2003. Rare and locally distributed palearctic species of the reptile parasitic worms: *Oxysomatium caucasicum* (Nematoda, Cosmocercidae). *Vestnik Zoologii*. 37 (5): 69–72. (in Russian).
38. Sharpilo V.P. 1962. On the study of the helminth fauna of reptiles in the Transcaucasia. *Collection of works of the Zoological Museum*. 31: 63–69 (in Russian).
39. Sharpilo V.P. 1974. A new member of the genus *Neoxysomatium* Ballesteros Marquez, 1945 (Nematoda, Cosmocercidae) a parasite of slow worms of the Caucasus. *Parazitologiya*. 8 (2): 112–114 (in Russian).
40. Kirillov A.A., Ruchin A.B., Fayzulin A.I. & Chihlyayev I.V. 2015. Helminths of reptiles of Mordovia: advance information. *Proceedings of the Mordovia State Nature Reserve* 14: 243–255 (in Russian).
41. Kirillov A.A., Kirillova N. Yu. & Ruchin A.B. 2018. Helminth fauna of *Anguis fragilis* L. (Anguidae) of Smolny National Park. *Proceedings of the Mordovia State Nature Reserve*. 21: 279–283 (in Russian).
42. Kirillov A.A., Kirillova N. Yu. 2018. Overview of helminths in reptiles of the National Park «Samarskaya Luka» (Russia). *Nature Conservation Research*. 3(Suppl. 1): 73–82. DOI: 10.24189/ncr.2018.049 (in Russian).
43. Kirillov A.A. 2000. Helminth fauna of reptiles of the Samara region. *Proceedings of Samara Scientific Center of the Russian Academy of Sciences*. 3: 324–329. (in Russian)
44. Sharpilo V.P. 1964. Larval forms of nematodes parasites of reptiles of the fauna of the Ukrainian SSR. *Works of Ukrainian Scientific Society of Parasitologists*. Kyiv: Naukova Dumka, pp. 112–124. (in Russian)
45. Shevchenko N.N., Belinisova L.K. 1964. About some new helminth species of reptiles of the Seversky Donets Valley in Kharkov region. *Questions of Genetics and Zoology*. Kharkov: Kharkov University, pp. 47–55. (in Russian)
46. Sargsyan N., Danielyan F., Arakelyan M. 2014. Seven new species of helminths for reptiles from Armenia. *Acta Parasitologica*. 59 (3): 442–447.
47. Rausch R.L. 1994. Family Mesocostoididae Fuhrmann, 1907. In: Khalil L.F., Jones A., Bray R.A. (eds.). *Keys to the cestode parasites of vertebrates*. CAB International, Wallingford, pp. 309–314.
48. Skirnisson K., Jouet D., Ferte H., Nielsen O.K. 2016. Occurrence of *Mesocostoides canislagopodis* (Rudolphi, 1810) (Krabbe, 1865) in mammals and birds in Iceland and its molecular discrimination within the *Mesocostoides* species complex. *Parasitology Research*. 115 (7): 2597–2607.
49. Lewin J. 1992a. Parasites of the sand lizard (*Lacerta agilis* L.) in Poland. *Acta Parasitologica Polonica*. 37: 19–24.
50. Sharpilo V.P., Biserkov V., Kostadinova A., Behnke J.M., Kuzmin Y.I. 2001. Helminths of the sand lizard *Lacerta agilis* (Reptilia, Lacertidae), in the Palaeartic: faunal diversity and spatial patterns of variation in the composition and structure of component communities. *Parasitology*. 123: 389–400.
51. Roca V., Ferragut M.V. 1989. Helminthofauna del lagarto verdinegro, *Lacerta schreiberi* Bedriaga, 1878 (Reptilia: Lacertidae) del Sistema Central (España). *Revista iberica de parasitologia*. 49: 291–300. (In Spanish)
52. Yildirimhan H.S., Bursey C.R., Altunel F.N. 2011. Helminth parasites of the Balkan green lizard *Lacerta trilineata* Bedriaga

- 1886 from Bursa, Turkey. Turkish Journal of Zoology. 35: 519–535.
53. Biserkov V., Kostadinova A. 1998. Intestinal helminth communities in the green lizard *Lacerta viridis* from Bulgaria. Journal of Helminthology. 72: 267–271.
54. Birlik S., Yildirimhan H.S., Sumer N., Ilgaz C., Kumlutas Y., Guglu O., Durmus S.H. 2015. The helminth fauna of *Apathya cappadocica* (Werner, 1902) (Anatolian Lizard) (Squamata: Lacertidae) from Turkey. Helminthologia. 52 (4): 310–315.
55. Birlik S., Yildirimhan H.S., Sümer N., Kumlutas Y., Ilgaz C., Durmus S.H., Guclu O., Candan K. 2016. Helminth fauna of Lebanon lizard, *Phoenicolacerta laevis* (Gray, 1838) (Squamata: Lacertidae) from Southern Turkey. Helminthologia. 53 (3): 262–269.
56. Gurelli G., Gocmen B., Cetin-Dogan T., Alpagut-Keskin N. 2007. First record of *Mesocestoides* spp. Vaillant, 1863. Tetrathyridia (Cestoidea: Cyclophyllidae) in Anatolian lizard, *Anatololacerta danfordi* (Gunther, 1876) in Turkey. North-Western Journal of Zoology. 3: 96–106.
57. Ikromov E.K., Cho M.R. 2004. On new representatives of the helminth fauna of reptiles (Testudines and Sauria) in Uzbekistan. Journal of Asia-Pacific Entomology. 7: 13–17.
58. Roca V., Lopez-Balaguer E., Hornero M.J. 1989. Helminthofauna de *Podarcis hispanica* (Steindachner, 1870) *Podarcis bocagei* (Seoane, 1884) (Reptilia: Lacertidae) en el Cuadrante Noroccidental de la Peninsula Iberica. Revista iberica de parasitologia. 49: 127–135. (in Spanish)
59. Kirin D. 2002a. New data on the helminth fauna of *Lacerta viridis* Laurenti, 1768, and *Podarcis muralis* (Laurenti, 1768) (Reptilia: Lacertidae) in Bulgaria. Acta Zoologica Bulgarica. 54: 43–48.
60. Roca V., Hornero M.J. 1994. Helminth infracommunities of *Podarcis pityusensis* and *Podarcis lilfordi* (Sauria: Lacertidae) from the Balearic Islands (western Mediterranean basin). Canadian Journal of Zoology. 72: 658–664.
61. Roca V., Lluch J., Navarro P. 1986b. Contribucion al conocimiento de la helminthofauna de los herpetos ibericos. V. Parasitos de *Psammotromus algirus* (L., 1758) Boulenger, 1887, *Psammotromus hispanicus* Fitzinger, 1926 y *Acanthodactylus erythrurus* (Schinz, 1833) Mertens, 1925 (Reptilia: Lacertidae). Boletin de la Real Sociedad Espanola de Historia Natural. (Biol.). 81: 69–78.
62. Roca V., Lluch J. 1988. L'helminthofauna des lacertidae (Reptilia) de la zone thermomediterraneenne de l'est de l'Espagne. Aspects ecologiques. Vie Milieu. 38: 201–205. (in Spanish)
63. Markov G.S., Ivanov V.P., Nikulin V.P. & Chernobay V.F. 1962. Helminth fauna of reptiles of the Volga delta and Caspian steppes. Pr. of Astrakhan Reserve 6: 145–172 (in Russian).
64. Biserkov V.Y. 1996. New records of platyhelminth parasites from snakes in Bulgaria. Comptes rendus de l'Académie bulgare des Sciences. 49: 73–75.
65. Lewin J. 1992b. Parasites of *Lacerta vivipara* Jacquin, 1787 in Poland. Acta Parasitologica Polonica. 37: 79–82.
66. Pojmanska T. Family Brachylaimidae Joyeux & Foley, 1930. In : Gibson D.I., Jones A. and Bray R.A. (eds.). Keys to the Trematoda. Vol. 1. CABI Publishing, Wallingford, UK and The Natural History Museum, London, 2002. pp. 544.
67. Galdon M.A., Roca V., Barbosa D., Carretero M.A. 2006. Intestinal helminth communities of *Podarcis bocagei* and *Podarcis carbonelli* (Sauria: Lacertidae) in NW Portugal. Helminthologia. 43: 37–41.
68. Hornero M.J., Roca V. 1992. Helminthofauna de *Podarcis lilfordi* (Gunther, 1874) (Sauria, Lacertidae) de los islotes de Menorca (Islas Baleares, Mediterraneo Occidental). Miscellanea Zoológica. Vol. 16: 1–6. (in Spanish)
69. Incedogan S., Yildirimhan H.S., Bursley C.R. 2014. Helminth parasites of the ocellated skink, *Chalcides ocellatus* (Foskal, 1775) (Scincidae) from Turkey. Comparative Parasitology. 81: 260–269.
70. Sudarikov V. E., Shigin A. A., Kurochkin Yu. V. et al. 2002. Metacercariae of trematode parasites of freshwater hydrobionts in Cen-

- tral Russia. Moscow: Nauka. 298 p. (in Russian).
71. Roca V., Lluh J., Navarro P. 1986a. Contribucion al conocimiento de la helminthofauna de los herpetos ibericos. 1. Parasitos de Lacertidae: *Lacerta lepida* Daudin, 1802 y *Podarcis hispanica* Steindachner, 1870. *Revista iberica de parasitologia*. 46: 129–136. (in Spanish)
 72. Dezfuli B.S. 2000. Host-parasite interface between *Asellus aquaticus* (Isopoda) and larvae of *Acanthocephalus anguillae* (Acanthocephala). *Folia parasitologica*. 47 (2): 154–156.
 73. Vakarenko A. G. & Lisitsina I. I. 2001. Cystacanth of *Acanthocephalus falcatus* (Acanthocephala, Echinorhynchidae) from the New Intermediate Host Woodlouse *Ligidium hypnorum* (Isopoda, Ligiidae). *Vestnik zoologii*. 35 (6): 67–69 (in Russian)
 74. Bertman, M. 1993. *Diplodiscus subclavatus* (Pallas, 1760) (Trematoda) and *Acanthocephalus ranae* (Schrank, 1788) (Acanthocephala) in grass snake – *Natrix natrix* (L.). *Wiadomosci Parazytologiczne*. 39: 405–406 (In Polish)
 75. Baker M. R. Revision of *Entomelas Travassos*, 1930 (Nematoda: Rhabdiasidae) with a review of genera in the family. *Systematic Parasitology*. 1980. Vol. 1 (2). P. 83–90.
 76. Moravec F. 1974. Description of the free-living males and females of *Entomelas entomelas* (Dujardin, 1845) (Nematoda: Rhabdiasidae). *Scripta Facultatis Scientiarum Naturalium Universitatis Purkynianae Brunensis*. 4: 97–100.
 77. Anderson R.C. 2000. *Nematode Parasites of Vertebrates. Their Development and Transmission*. 2nd Edition. Wallingford, Oxon: CABI Publishing. 650 p.
 78. Halajian A., Bursey C.R., Goldberg S.R., Mohammad Ali Gol S. 2013. Helminth Parasites of the European Glass Lizard, *Pseudopus apodus* (Squamata: Anguillidae), and European Grass Snake, *Natrix natrix* (Serpentes: Colubridae), from Iran. *Comparative Parasitology*. 80 (1): 151–156.
 79. Kuzmin Yu. 2013. Review of Rhabdiasidae (Nematoda) from the Holarctic. *Zootaxa*. 3639 (1): 1–76.
 80. Chu T. 1936. Studies on the life history of *Rhabdias fuscovenosa* var. *catanensis* (Rizzo 1902). *Journal of Parasitology*. 22: 140–160.
 81. Yildirimhan H.S., Bursey C.R., Goldberg S. R. 2007. Helminth parasites of the grass snake *Natrix natrix* and the dice snake *Natrix tessellata* (Serpentes: Colubridae) from Turkey. *Comparative Parasitology*. 74 (2): 343–354.
 82. Bakiev A., Kirillov A., Mebert K. 2011. Diet and parasitic helminths of Dice snakes from the Volga Basin, Russia. *Mertensiella*. 18: 325–329.
 83. Ruchin A.B. & Kirillov A.A. 2012. The helminthofauna of the grass snake *Natrix natrix* L. from Republic of Mordovia. *Biological Sciences of Kazakhstan*. 4: 344–348 (in Russian)
 84. Reshetnikov A.N., Sokolov S.G., Chikhlyayev I.V., Faizulin A.I., Kirillov A.A., Kuzovenko A.E., Protasova E.N., Skomorokhov M.O. 2013. Direct and indirect interactions between an invasive Alien Fish (*Perccottus glenii*) and two native semi-aquatic snakes. *Copeia*. 1: 103–110.
 85. Yossefi M.R., Nikzad R., Nikzad M., Mousapour A., Ramazanpour S., Rahimi M.T. 2014. High helminthic infection of the European grass snake *Natrix natrix* and the dice snake *Natrix tessellata* (Serpentes: Colubridae) from Iran. *Asian Pacific Journal of Tropical Disease*. 4 (Supp. 1): S263–S267.
 86. Shevchenko, N.N., Barabashova, V.N. 1958. Helminth fauna of *Lacerta agilis* L. and *Vipera berus* L. in the Kharkov area. *Works on Helminthology*. P. 389–394. (in Russian)
 87. Hendrikx W.M.L. 1983. Observations on the routes of infection of *Oswaldocruzia filiformis* (Nematoda: Trichostrongylidae) in Amphibia. *Zeitschrift für Parasitenkunde*. 69 (1): 119–126.
 88. Ben Slimane B., Durette-Desset M.C. 1991. Quatre nouvelles especes du genre *Oswaldocruzia* Travassos, 1917 (Nematoda: Trichostrongyloidea) parasites d'Amphibiens d' Equateur. *Revue Suisse de Zoologie*. 100: 113–136.
 89. Ben Slimane B., Durette-Desset M.C. 1996. Two new *Oswaldocruzia* (Nematoda, Trichostrongylina, Molineoidea) parasites of

- amphibians from Cameroon. *Helminthologia*. 33: 143–147.
90. Ben Slimane B., Durette-Desset M.C. 1997. Revision du genre *Oswaldocruzia* (Nematoda, Trichostrongylida, Molineoidea) en zone nearctique avec description de cinq nouvelles espèces. *Zoosystema*. 19: 61–79.
91. Durette-Desset M.C., Batcharov A., Ben Slimane B., Chabaud A.G. 1993. Some *Oswaldocruzia* (Nematoda: Trichostrongyloidea) parasites of Amphibia in Bulgaria. Re-description of *Oswaldocruzia bialata* (Molin, 1860). *Helminthologia*. 30 (2): 99–104.
92. Ben Slimane B., Durette-Desset M.C., Chabaud A.G. 1995a. Phenomenes de speciation chez les *Oswaldocruzia* (Nematoda: Trichostrongylina, Molineoidea) parasites de Lezards de Guadeloupe. *Parasite*. 2: 373–383.
93. Ben Slimane B., Lluch J., Durette-Desset M.C. 1995b. Two new species of the genus *Oswaldocruzia* Travassos, 1917 (Nematode: Trichostrongylina: Molineidae) parasitizing Spanish amphibians. *Research and Review in Parasitology*. 55 (4): 209–215.
94. Ben Slimane B., Chabaud A.G., Durette-Desset M.C. 1996a. Les Nematodes *Trichostrongylina* parasites d'Amphibiens et de Reptiles: problemes taxonomiques, phyletiques et biogeographiques. *Systematic parasitology*. 35 (3): 179–206.
95. Ben Slimane B., Guerero R., Durette-Desset M.C. 1996b. *Oswaldocruzia venezuelensis* n.sp. (Nematoda, Trichostrongyloidea, Molineoidea) a parasite of *Bufo marinus* from Venezuela. *Folia Parasitologica*. 43 (4): 297–300.
96. Svitin R. & Gorobchishyn V. 2015. Postcycle parasitism of nematodes of the genus *Oswaldocruzia* Travassos, 1917 parasitizing several species of snakes from the territory of Ukraine. *Bulletin of Taras Shevchenko national university of Kyiv* 1 (18): 68–70 (in Ukrainian)
97. Okulewicz A. 1976. *Oswaldocruzia filiformis* (Goeze, 1782) Travassos 1917 – Nowy pasożyt jaszczurki zwinki (*Lacerta agilis* L.). *Wiadomosci Parazytologiczne*. 12: 297–301. (in Polish)
98. Marconcini A., Triantafyllou G. 1970. Note helmintologiche sulla fauna selvatica in Italia. *Annali della Facolta di Medicina Veterinaria di Pisa*. 22: 217–229. (in Italian)
99. Roca V., Jorge F., Ilgaz C., Kumlutas Y., Durmus S.H., Carretero M.H. 2016. Intestinal parasites of unisexual and bisexual lizards *Darevskia* spp. (Lacertidae) from Northeastern Anatolia. *Helminthologia*. 53 (3): 298–303.
100. Sanchis V., Roig J.M., Carretero M.A., Roca V., Llorente G.A. 2000. Host-parasite relationships of *Zootoca vivipara* (Sauria: Lacertidae) in the Pyrenees (North Spain). *Folia Parasitologica*. 47: 118–122.
101. Kirin D. 2002b. New records of the helminth fauna from grass snake *Natrix natrix* L., 1758 and dice snake *Natrix tessellata* Laurenti, 1768 (Colubridae: Reptilia) in South Bulgaria. *Acta Zoologica Bulgarica*. 54: 49–53.
102. Markov G.S., Mozgovoï A.A. 1969. Causes of the low level of helminth infection in *Vipera berus* in the northern European part of its distribution area. *Proceedings of the Helminthological Laboratory of Academy of Sciences of the USSR*. 20: 91–96 (in Russian)
103. Novokhatskaya O.V. 2008. On the occurrence of the nematode *Oswaldocruzia filiformis* (Strongylida: Molineidae) in Karelia. *Parazitologiya*. 42: 204–209. (in Russian)
104. Kirillov A.A., Kirillova N.Yu. 2016. Analysis of the reproductive structure of the hemipopulation of the *Cosmocerca ornata* (Dujardin, 1845) (Nematoda: Cosmocercidae) in marsh frogs of different ages. *Inland Water Biology*. 9 (3): 310–318.
105. Hoberg E.P., Polley L., Jenkins E.J., Kutz S.J., Veitch A.M., Elkin B.T. 2008. Integrated approaches and empirical models for investigation of parasitic diseases in northern wildlife. *Emerging Infectious Diseases Journal*. 14: 10–17.
106. Kafle P., Sullivan J., Verocai G.G., Kutz S.J. 2017. Experimental Life-Cycle of *Varestrongylus eleguneniensis* (Nematoda: Protostrongylidae) in a Captive Reindeer (*Rangifer tarandus tarandus*) and a Muskox

- (*Ovibos moschatus moschatus*). *Journal of Parasitology*. 103 (5): 584–587. doi: 10.1645/17-19.
107. Roepstorff A., Nansen P. 1998. Epidemiology, diagnosis, and control of helminth parasites of swine. FAO of the UN, Rome. pp. 161.
108. Biserkov V.Y. 1995. New records of nematodes and acanthocephalans from snakes in Bulgaria. *Comptes rendus de l'Academie bulgare des sciences: sciences mathematiques et naturelles*. 48 (11-12): 87–89.
109. Bogdanov O.P., Lutta A.S., Markov G.S., Ryzhikov K.M. 1969. New data about parasite fauna of *Ancystrodon halys* (Reptilia, Crotalidae). *Zoological Journal*. 68: 179–183. (in Russian)
110. Santos X., Martinez-Freiria F., Pleguezuelos J.M., Roca V. 2006. First helminthological data on Iberian vipers: Helminth communities and host-parasite relationships. *Acta Parasitologica*. 51: 130–135.
111. Cordero del Campillo M., Castanon-Ordonez L., Reguera-Feo A. 1994. *Indice-catalogo de zooparasitos ibericos*. 2 edicion. Secretariado de Publicaciones, Universidad de Leon. Leon. 650 pp.
112. Markov G.S., Lukina G.P., Markova L.I. & Mozgina A.A. 1964. To the parasitic fauna of reptiles of the North Caucasus. *Memoirs of Volgograd State Pedagogical University* 16: 99–105. (in Russian)