



## Medicinal Plants as Potent Power for Malaria Control: Review

Mathalaimuthu Baranitharan<sup>1\*</sup>, Vairakannu Tamizhazhagan<sup>2</sup>,  
Kalimuthu Kovendan<sup>1</sup>

<sup>1</sup> Department of Zoology, Annamalai University, Annamalainagar-608 002, Tamilnadu, India.

<sup>2</sup> Department of Zoology, Syed Ammal Arts and Science College, Ramanathapuram- 623513, Tamilnadu, India.

### ABSTRACT

Mosquito vectors are living organisms that can transmit infectious diseases between human from animals to humans. It is bloodsucking insect that ingest disease-producing microorganisms during a blood meal from an infected host. This review aims to explain some medicinal plants that exhibit mosquitocidal agents on genus *Anopheles* mosquito species. Currently, the utilization of artificial chemicals to regulate insects and mosquitoes raises many considerations associated with environment and human health. Another aim is to use natural merchandise that possess sensible effectiveness, simple handiness, fewer adverse environmental impacts, and are environmentally friendly have crystal rectifier to the multiplied interest in plant origin chemical pesticides. It's supported original articles obtained by looking on major databases. Our literature review disclosed that 96 medicinal plants used in the all parts (leaves, root, bark and bud). This article envisaged to review the reports of ovicidal, oviposition deterrent, larvicidal, pupicidal and repellent activities, which might be employed in vector-borne disease management programs.

**Keywords:** Mosquitocidal Activity, Medicinal Plants, *Anopheles* Species.

**HOW TO CITE THIS ARTICLE:** Mathalaimuthu Baranitharan, Vairakannu Tamizhazhagan, Kalimuthu Koven-dan; Medicinal Plants as Potent Power for Malaria Control: Review, Entomol Appl Sci Lett, 2019, 6 (1): 28-44.

**Corresponding author:** Mathalaimuthu Baranitharan

**E-mail** ✉ [bharanitharan2011@gmail.com](mailto:bharanitharan2011@gmail.com)

**Received:** 21/11/2018

**Accepted:** 18/03/2019

### INTRODUCTION

#### Malaria

Mosquitoes have been accounted as the well-spring of different diseases influencing human. Including around 3500 species, mosquitoes are discovered past the tropical and subtropical locales of the world [1]. Malaria in people is brought on by contamination with one or a greater amount of a few types of *Plasmodium* (i.e., *Plasmodium falciparum*, *Plasmodium vivax*, *Plasmodium ovale*, *Plasmodium malariae*, and incidentally other *Plasmodium* species) parasites. The parasite is transmitted by the bite of an infective female *Anopheles* mosquito. *P. falciparum* and *P. vivax* species cause the most contaminations around the world. *P. falciparum* is the operators that most usually cause serious and possibly lethal malaria. As indicated by the most as of late accessible data, an expected 207

million clinical cases and 627,000 (0.3%) deaths were reported worldwide in 2012, for the most part among children matured five years living in sub-Saharan Africa. *P. vivax* and *P. ovale* have torpid liver stages, which can reactivate and cause malaria a while for years after the underlying disease. *P. malariae* can bring about durable diseases and if untreated or insufficiently treated, can continue asymptotically in the human host for quite a long time, even a lifetime.

#### Malaria disease worldwide

*Anopheles stephensi* transmits of malaria fever in urban district of India. Among 53 Anopheline species present in India, 9 are vectors of malaria fever [2] and malaria is still the most critical reason for mortality with around a few million new cases emerging each year. A more effective approach to decrease mosquito populace is to focus on the hatchlings. The main arrangement is to keep the ailment conveying mosquito from

reproducing and gnawing people. In this situation, mosquito vector control is pivotal. *An. subpictus* Grassi is distributed throughout India, China, Afghanistan, Borneo, Malaysia, Sri Lanka, Philippines, Indonesia, and Java. It is a dominant species in Haryana and Uttaranchal states. Though it is a non-vector species, same infected specimens with malaria parasite have been reported from India, Indonesia, and Java. *An. subpictus* is a significant secondary vector in Sri Lanka [3, 4]. *An. subpictus* is recognized as the secondary vector of malaria in South East Asia, with a large number of cases being reported from India. India contributes 77% of the world's population [5, 6]. Members of the *An. gambiae* complex are the most important vectors of malaria in sub-Saharan Africa of which *An. gambiae* and *An. arabiensis* are the most widely distributed and most efficient vectors [7]. Mosquito larvae and pupae are normally focused on utilizing organophosphates, insect development controllers and microbial control operators. Inside lingering showering and insect spray treated bed net are likewise utilized to decrease transmission of malaria in tropical nations. Notwithstanding, engineered chemicals have solid negative consequences for human wellbeing and nature, and incite resistance in various mosquito species [8, 9].

#### **Eruption of malaria disease in the world**

Mosquitoes have been accounted for as the well-spring of different diseases influencing human, and are discovered in the tropical and subtropical locales of the world. In Asia, the mosquito is the most important vector of malaria, with 200 to 450 million infections annually worldwide; it sources up to 2.7 million deaths. WHO Malaria Report estimates that 3.3 billion people were at the peril of malaria in 2010, although of all geographical regions, populations living in sub-Saharan Africa (SSA) have the highest peril of obtaining malaria; among 216 million episodes of malaria in 2010, approximately 81%, or 174 million occasions, were noticed from the African region. There were estimated malaria demises of 655,000 in 2010, of which 91% were from Africa [7]. Malaria afflicts 36% of the world people i.e. 2020 million in 107 countries and territories position in the tropical and subtropical regions.

As indicated by the most recent evaluations, there were around 198 million cases with malaria fever in 2013 and an expected 584,000 passing. Most passing's happen among children living in Africa. Malaria death rates among children in Africa have been decreased by an expected percentage of 58% since 2000. Globally, 4.28 million deaths have been averted due to malaria control efforts between 2001 and 2013, of which 3.92 million (92%) are in children under 5 years of age in sub-Saharan Africa. They account for 20% of the 20 million child deaths; it has been estimated that it has been averted in sub-Saharan Africa between 2001 and 2013 through the overall reduction in child mortality rates [3]. In 2013, around 6.3 million children died before their fifth birthday, at a rate of around 17,000 per day. The risk of dying before age five varies enormously depending on where a child is born. In Luxembourg, the under-five mortality pace is fair 2 deaths per 1,000 live births; in Angola it is 167 per 1,000. Since 1990, 223 million children have died before their fifth birthday [9]. Malaria deaths dropped by 58% in African children under five years old from 2000-2013 [3]. As per the most recent evaluations from WHO, there were 214 million new instances of malaria worldwide in 2015 and there were an expected 438,000 malaria passing. Children under five are especially helpless to malaria ailment, disease and demise. Malaria slaughtered an expected 306,000 under-fives comprehensively, incorporating 292,000 kids in the African region [4]. Malaria incidence rate has declined by 41% since 2000, and 2010. Mortality has fallen by 61% since 2000 and 29% since 2015. Seventeen countries eliminated malaria between 2000 and 2015, with a further 13 countries "approaching elimination" [10]. According to the reports, India has 6% of all malaria cases in the world, 6% of the deaths, and 51% of the cases in global *P. vivax* cases. The report estimates the total cases in India as 1.31 million and the deaths as 194 [11]. History uncovers that the occurrence of malaria in India happened in the 1945s with an expected 75 million cases and 800,000 passed for each year. This makes female mosquito a perfect transmitter of different blood borne pathogens and operators of destroying human diseases. The year's malaria has been considered as one of the main sources of

deaths in India (<http://nvbdcp.gov.in/malaria-new.html>) (Table 1).

#### **Medicinal plants based mosquito control**

Medicinal plants might be elective wellsprings of mosquito control specialists since they constitute a rich wellspring of bioactive compounds that are biodegradable and conceivably suitable to control mosquitoes. Pesticides of organic inception might serve as suitable option for bio-control procedures later on [12]. The use of synthetic chemicals for management of mosquitoes raises many issues connected to the environment and human health [13]. Eco-friendly tools to manage mosquito young instar populations in an IPM framework are urgently needed. Natural products are usually most preferred due to their less harmful nature and fast biodegradability. Plant products have been used traditionally by human communities in different rural areas worldwide against insect vectors and parasites [14]. Most of the mosquito control programs target the larval stage in their breeding sites with larvicides. Monoterpenes such as  $\alpha$ -pinene, limonene, terpinolene, citronellol, citronellal, camphor and thymol are regular constituents of a number of essential oils of medicinal plants. The chemicals derived from plants have been projected as weapons in future mosquito control program as they are shown to act as general toxicant, growth and reproductive inhibitors, repellents and oviposition-deterrent [15]. Table 2 gives definite surveys of plant items reported for mosquitocides and insecticides tested against malarial species.

#### **CONCLUSION**

Plants as option for wellspring of larvicidal, ovicidal, oviposition deterrent, pupicidal and repellent activity reported in various ethnobotanical reviews. All the tried plants had diverse scope of larvicidal, ovicidal, oviposition deterrent, pupicidal and repellent action which might be utilized as traditional mosquito control operators. On the premise of the present examination results we presume that therapeutic plants extracts contains strong mosquitocidal bioactive standards which might be required further sanitizations to have its manufactured analogs, which will be done in future.

#### **ACKNOWLEDGEMENTS**

The authors are appreciative to superior power for stipend of money related help with Award of University Research Fellowship and Professor and Head, Department of Zoology, Annamalai University for the laboratory facilities provided. We acknowledge the members of the University Grants Commission (UGC), New Delhi, for their financial assistance (Ref. No. F 42-597/2013SR).

#### **REFERENCES**

1. Kalita B, Bora S, Sharma AK (2013) Plant essential oils as mosquito repellent-a review. *Int J Res Dev Pharm L Sci* 3:741-747
2. Baranitharan M, Dhanasekaran S, Murugan K, Kovendan K, Gokulakrishnan J, Benelli G (2017) *Coleus aromaticus* leaf extract fractions: A source of novel ovicides, larvicides and repellents against *Anopheles*, *Aedes* and *Culex* mosquito vectors? *Proc Safe Environ Prot* 106: 23-33.
3. World Malaria Report (2014) Malaria fact sheet December. WHO Press, 38.
4. World Health Organization (2015) Malaria fact sheet December. WHO Press, 9.
5. UNICEF (2013) Committing to Child Survival: A Promise Renewed - Progress Report, p.4.
6. IGME (2013) Levels and trends in child mortality: Report. New York, the United Nations Interagency Group for Child Mortality Estimation.
7. World Health Organization (2012) Handbook for integrated vector management. WHO; Geneva.
8. World Health Organization (2014) World Malaria Report. Malaria Fact sheet December. WHO Press, p.40.
9. UNICEF (2014) Committing to Child Survival: A Promise Renewed - Progress Report, p.5, 13, 16.
10. World Health Organization (2016) World Malaria Report. Geneva.
11. World Health Organization (2017) World malaria report, Geneva.
12. Veerakumar K, Govindarajan M (2014) Adulticidal properties of synthesized

- silver nanoparticles using leaf extracts of *Feronia elephantum* (Rutaceae) against filariasis, malaria, and dengue vector mosquitoes. Parasitol Res doi:10.1007/s00436-014-4077-4
13. Benelli G (2015) Research in mosquito control: current challenges for a brighter future. Parasitol Res 2801-2805.
  14. Baranitharan M, Dhanasekaran S, Murugan K, Kovendan K, Gokulakrishnan J (2016) Chemical composition and laboratory investigation of *Melissa officinalis* essential oil against human malarial vector mosquito, *Anopheles stephensi* L. (Diptera: Culicidae). J Coast Life Med 4: 969-973. <https://doi.org/10.12980/jclm.4.201616-174>
  15. Dhanasekaran S, Baranitharan M, Muthulingam M, Senthilmurugan S, Jeyasankar A, Barbara Sawicka (2018) Studies on the impact of medicinal plants in relation to malaria vector control against *Anopheles stephensi*. Innoriginal Int J Sci 5(3): 12-14.
  16. Bagavan A, Abdul Rahuman A (2010) Evaluation of larvicidal activity of medicinal plant extracts against three mosquito vectors. Asia Pac J Tro Med 29-34.
  17. Kovendan K, Murugan M, Mahesh Kumar P, Thiyagarajan P, John William S (2013) Ovicidal, repellent, adulticidal and field evaluations of plant extract against dengue, malaria and filarial vectors. Parasitol Res 112: 1205-1219.
  18. Krishnappa K, Elumalai K (2014) Mosquitocidal activity of indigenous plants of Western Ghats, *Achras sapota* Linn. (Sapotaceae) and *Cassia auriculata* L. (Fabaceae) against a common malarial vector, *Anopheles stephensi* Liston (Culicidae: Diptera). J Coast Lif Med 2: 402-410.
  19. Krishnappa K, Elumalai K, Dhanasekaran S, Gokulakrishnan J (2012) Larvicidal and repellent properties of *Adansonia digitata* against medically important human malarial vector mosquito *Anopheles stephensi* (Diptera: Culicidae). J Vector Borne Dis 49: 86-90.
  20. Kamaraj C, Abdul Rahuman A, Bagavan A, Abdul Zahir A, Elango G, Kandan P, et al (2010) Larvicidal efficacy of medicinal plant extracts against *Anopheles stephensi* and *Culex quinquefasciatus* (Diptera: Culicidae). Trop Biomed 27: 211-219.
  21. Dinesh D, Murugan K, Madhiyazhagan P, Panneerselvam C, Mahesh Kumar P, Nicoletti M, et al (2015) Mosquitocidal and antibacterial activity of green-synthesized silver nanoparticles from *Aloe vera* extracts: towards an effective tool against the malaria vector *Anopheles stephensi*? Parasitol Res 114: 1519-1529.
  22. Santhosh SB, Yuvarajan R, Natarajan D (2015) *Annona muricata* leaf extract-mediated silver nanoparticles synthesis and its larvicidal potential against dengue, malaria and filariasis vector. Parasitol Res 114: 3087-3096.
  23. Balu Selvakumar, Gokulakrishnan J, Elanchezhian K, Deepa J (2015) Mosquito larvicidal, ovicidal and pupicidal activities of *Annona reticulata* linn (Annonaceae) against *Aedes aegypti* (Linn.), *Anopheles stephensi* Liston and *Culex quinquefasciatus* (Say) (Diptera: Culicidae). Int J Rec Sci Res 6: 2690-2696.
  24. Younoussa Lame, Elias Nchiwan Nukenine, Danga Yinyang Simon Pierre, Ajaegbu Eze Elijah, Charles Okechukwu Esimone (2015) Laboratory Evaluations of the Fractions Efficacy of *Annona senegalensis* (Annonaceae) Leaf Extract on Immature Stage Development of Malarial and Filarial Mosquito Vectors. Arthropod-Borne Dis 9: 226-237.
  25. Vijaya kumar S, Panagal Mani, John Bastin TMM, Arun kumar R, Ravikumar G (2011) Larvicidal, oviposition deterrent and repellent activity of *Annona squamosa* extracts against hazardous mosquito vectors. Int J Pharm Tech 3: 3143-3155.
  26. Kuppan Velu, Deven Elumalai, Periaswamy Hemalatha, Arumugam Janaki, Muthu Babu, Maduraiveeran Hemavathi, et al (2015) Evaluation of silver nanoparticles toxicity of *Arachis hypogaea* peel extracts and its larvicidal activity against malaria and dengue

- vectors. Environ Sci Pollut Res DOI10.1007/s11356-015-4919-3.
27. Panneerselvam C, Murugan K, Kovendan K, Mahesh Kumar P (2012) Mosquito larvicidal, pupicidal, adulticidal, and repellent activity of *Artemisia nilagirica* (Family: Compositae) against *Anopheles stephensi* and *Aedes aegypti*. Parasitol Res 111: 2241-2251.
  28. Murugan K, James Pitchai J, Madhiyazhagan P, Nataraj T, Nareshkumar A, Jiang-Shiou Hwang, et al (2014) Larvicidal, repellent and smoke toxicity effect of neem products against malarial vector, *Anopheles stephensi*. Int J Pure Appl Zool 2: 71-83.
  29. Nganjiwa JI, Pukuma MS, Philimon J, Ekundayo TM (2015) Evaluation of larvicidal properties of leaves and root ethanolic extracts of some plants herbs against fourth instar mosquito larvae. Int J Sci Res Pub 5: 1-7.
  30. Kovendan K, Murugan K, Prasanna Kumar K, Panneerselvam C, Mahesh Kumar P, Amerasan D, et al (2012) Mosquitocidal properties of *Calotropis gigantea* (Family: Asclepiadaceae) leaf extract and bacterial insecticide, *Bacillus thuringiensis*, against the mosquito vectors. Parasitol Res 111: 531-544.
  31. Subarani S, Selvi Sabhanayakam, Kamaraj C, Elango G, Mohamed Abdul Kadir (2013) Efficacy of larvicidal and pupicidal activity of *Catharanthus roseus* aqueous and solvent extracts against *Anopheles stephensi* Liston and *Culex quinquefasciatus* Say (Diptera: Culicidae). Asia Pac J Trop Med 625-630.
  32. Jagruti H, sukhthankar, hemanth Kumar, Godinho MHS, Ashwani Kumar (2014) Larvicidal activity of methanolic leaf extracts of plant, *Chromolaena odorata* L. (Asteraceae) against vector mosquitoes. Int J Moquit Res 1: 33-38.
  33. Muthukumaran U, Govindarajan M (2015) Mosquito larvicidal potential of silver nanoparticles synthesized using *Chlomelia asiatica* (Rubiaceae) against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae). Parasitol Res 114: 989-999.
  34. Arokiyaraj S, Dinesh Kumar V, Elakya V, Kamala T, Sung Kwon Park, Ragam M, et al (2015) Biosynthesized silver nanoparticles using floral extract of *Chrysanthemum indicum* L.-potential for malaria vector control. Environ Sci Pollut Res 22: 9759-9765.
  35. Krishnappa K, Mathivanan T, Elumalai, A, Jeyasankar A, Dhanasekaran S, Elumalai K (2013) Evaluation of *Cissus quadrangularis* and *Combretum ovalifolium* Medicinal Plants Extracts against Medically Important Human Malarial Vector Mosquito *Anopheles stephensi* Liston (Diptera:Culicidae). Int J Inter Dis Res Rev 1: 11-18.
  36. Ajithadas Aruna, Vijayalakshmi K, Karthikeyan V (2014) Larvicidal activity of Methanolic Extract of the leaves of *Citrullus lanatus*. Int J Advanc Pharm Biol Chem 3: 717-722.
  37. Murugan K, Mahesh Kumar P, Kovendan K, Amerasan D, Subrmaniam J, Jiang-Shiou Hwang (2012) Larvicidal, pupicidal, repellent and adulticidal activity of *Citrus sinensis* orange peel extract against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae). Parasitol Res 111: 1757-1769.
  38. Govindarajan M (2010) Chemical composition and larvicidal activity of leaf essential oil from *Clausena anisata* (Willd.) Hook. F. Ex. Benth (Rutaceae) against three mosquito species. Asia Pac J Trop Med 874-877.
  39. Arivoli S, Samuel T (2011) Larvicidal efficacy of *Cleistanthus collinus* (Roxb.) (Euphorbiaceae) leaf extracts against vector mosquitoes (Diptera: Culicidae). Asia Pac J Trop Biomed S281-S283.
  40. Kovendan K, Murugan K (2011) Effect of Medicinal Plants on the Mosquito Vectors from the Different Agroclimatic Regions of Tamil Nadu, India. Adv Environ Biol 5: 335-344.
  41. Rahul B Salunkhe, Satish V Patil, Chandrashekhar D Patil, Bipinchandra K Salunke (2011) Larvicidal potential of silver nanoparticles synthesized using fungus *Cochliobolus lunatus* against *Aedes aegypti* (Linnaeus, 1762) and *Anopheles*

- stephensi* Liston (Diptera; Culicidae). Parasitol Res 109: 823-831.
42. Baranitharan M, Dhanasekaran S, Mahesh Babu S, Sridhar N, Krishnappa K (2014) Larvicidal activity of *Coleus aromaticus* benth (Lamiaceae) leaf extracts against three vector mosquitoes. Int J Res Bio Sci 4: 55-59.
  43. Baranitharan M, and Dhanasekaran S (2014) Mosquito larvicidal properties of *Commiphora caudata* (Wight & Arn.) (Bursaceae) against *Aedes aegypti* (Linn.), *Anopheles stephensi* (Liston), *Culex quinquefasciatus* (Say). Int J Curr Microbiol App Sci 3: 262-268.
  44. Elangovan A, Dhanasekaran S, Anandan A, Krishnappa K, Gokulakrishnan J, Elumalai K (2012) Mosquitocidal activities of *Corchorus capsularis* L (Malvaceae) against a common malarial vector, *Anopheles stephensi* (Liston) and a dengue vector *Aedes aegypti* (L) (Diptera: Culicidae). Int J Recen Scien Res 3: 564-568.
  45. Karunamoorthi K, Ilango K (2010) Larvicidal activity of *Cymbopogon citratus* (DC) Stapf. and *Croton macrostachyus* Del. against *Anopheles arabiensis* Patton, a potent malaria vector. Eur Rev Med Pharmacol Sci 14: 57-62.
  46. Singha S, Chandra G (2011) Mosquito larvicidal activity of some common spices and vegetable waste on *Culex quinquefasciatus* and *Anopheles stephensi*. Asia Pac J Trop Med 288-293.
  47. Murugan K, Dinesh D, Jenil Kumar P, Panneerselvam C, Subramaniam J, Madhiyazhagan P, et al (2015) *Datura metel*-synthesized silver nanoparticles magnify predation of dragonfly nymphs against the malaria vector *Anopheles stephensi*. Parasitol Res 114: 4645-4654.
  48. Swathi S, Muruganantham G, Ghosh SK, Pradeep AS (2012) Larvicidal and repellent activities of ethanolic extract of *Datura stramonium* leaves against mosquitoes. Int J Pharmac Phytoche Res 4: 25-27.
  49. Dohutia C, Bhattacharyya DR, Sharma SK, Mohapatra PK, Bhattacharjee K, Gogoi K, et al (2015) Larvicidal activity of few select indigenous plants of North East India against disease vector mosquitoes (Diptera: Culicidae). Trop Biomed 32: 17-23.
  50. Srinivasan R, Natarajan D, Shivakumar MS, Vinuchakkaravarthy T, Velmurugan D (2015) Bioassay guided isolation of mosquito larvicidal compound from acetone leaf extract of *Elaeagnus indica* Servett Bull and its in-silico study. Industrial Crop Produc 76: 394-401.
  51. Elumalai K, Dhanasekaran S, Anandan A, Krishnappa K, Gokulakrishnan J, Elangovan A (2012) Larvicidal, ovicidal and pupicidal activity of *Eranthemum roseum* (Vahl) R. Br. against malarial vector mosquito, *Anopheles stephensi* (Liston) (Diptera: Culicidae). Int J Curr Agricul Scie 2: 28-33.
  52. Govindarajan M, Sivakumar R, Amsath A, Niraimathi S (2012) Larvicidal efficacy of botanical extracts against two important vector mosquitoes. Eur Rev Med Pharmacol Sci 16: 386-392
  53. Prathibha KP, Raghavendra BS Vijayan VA (2010) Evaluation of larvicidal effect of *Euodia ridleyi* Hochr. leaf extract against three mosquito species at Mysore. Res J Biol Sci 5: 452-455.
  54. Agalya Priyadarshini K, Murugan K, Panneerselvam C, Ponarulselvam S, Jiang-Shiou Hwang, Marcello Nicoletti (2012) Biolarvicidal and pupicidal potential of silver nanoparticles synthesized using *Euphorbia hirta* against *Anopheles stephensi* Liston (Diptera: Culicidae). Parasitol Res 111: 997-1006.
  55. Elangovan A, Dhanasekaran S, Anandan A, Krishnappa K, Gokulakrishnan J, Elumalai K (2012) Larvicidal and ovicidal activities of *Exacum pedunculatum* (Linn.) (Gentianaceae) against a common malarial vector, *Anopheles stephensi* Liston (Diptera : Culicidae). Int J Recen Scien Res 3: 559-563.
  56. Govindarajan M, Sivakumar R, Amsath A, Niraimathi S (2011) Mosquito larvicidal properties of *Ficus benghalensis* L. (Family: Moraceae) against *Culex tritaeniorhynchus* Giles and *Anopheles*

- subpictus* Grassi (Diptera: Culicidae). Asia Pac J Trop Med 505-509.
57. Krishnappa K, Dhanasekaran S, Elumalai K (2012) Larvicidal, ovicidal and pupicidal activities of *Gliricidia sepium* (Jacq.) (Leguminosae) against the malarial vector, *Anopheles stephensi* Liston (Culicidae: Diptera). Asia Pac J Trop Med 598-604.
  58. Dhanasekaran S, Krishnappa K, Anandan A, Elumalai KJ (2013) Larvicidal, ovicidal and repellent activity of selected indigenous medicinal plants against malarial vector *Anopheles stephensi* (Liston), dengue vector *Aedes aegypti* (Linn.), Japanese encephalitis vector, *Culex tritaeniorhynchus* (Giles.) (Diptera: Culicidae). Agri Technol 9: 29-47.
  59. Gopiesh Khanna V, Kannabiran K, Rajakumar G, Abdul Rahuman A, Santhoshkumar T (2011) Biolarvicidal compound gymnemagenol isolated from leaf extract of miracle fruit plant, *Gymnema sylvestri* (Retz) Schult against malaria and filariasis vectors. Parasitol Res 109: 1373-1386.
  60. Bashir Ahmad, Javid Ali (2013) Evaluation of Larvicidal Activity of *Hippophae rhamnoides* L. Leaves Extracts on *Aedes aegypti* and *Anopheles stephensi* (Diptera: Culicidae). Middle-East J Sci Res 13: 703-709.
  61. Suman TY, Radhika Rajasree SR, Jayaseelan C, Regina Mary R, Gayathri S, Aranganathan S, et al (2015) GC-MS analysis of bioactive components and biosynthesis of silver nanoparticles using *Hybanthus enneaspermus* at room temperature evaluation of their stability and its larvicidal activity. Environ Sci Pollut Res DOI 10.1007/s11356-015-5468-5.
  62. Zewdneh Tomass, Mamuya Hadis, Asegid Taye, Yalemtehey Mekonnen, Beyene Petros (2011) Larvicidal effects of *Jatropha curcas* L. against *Anopheles arabiensis* (Diptera: Culicidae). CNCS Mekelle University 3: 52-64.
  63. Kaliyaperumal Karunamoorthi, Askual Girmay, Samuel Fekadu (2014) Larvicidal efficacy of Ethiopian ethnomedicinal plant *Juniperus procera* essential oil against Afrotropical malaria vector *Anopheles arabiensis* (Diptera: Culicidae). Asia Pac J Trop Biomed 4: S99-S106.
  64. Babita R. Malik, Mukesh K. Malik, Balakrishnan N, Suresh B (2014) Evaluation of larvicidal activity of the different extracts against important species of mosquito: *Anopheles stephensi*. J Parasitol Vec Biol 6: 11-15.
  65. Arivoli S, John Ravindran K, Tennyson S (2012) Larvicidal Efficacy of Plant Extracts against the Malarial Vector *Anopheles stephensi* Liston (Diptera: Culicidae). World J Med Scien 7: 77-80.
  66. Kovendan K, Shanthakumar S, Praseeja C, Mahesh Kumar P, Murugan K, Vincent S (2014) Mosquitocidal properties of *Morinda citrifolia* L. (Noni) (Family: Rubiaceae) leaf extract and *Metarhizium anisopliae* against malaria vector, *Anopheles stephensi* Liston. (Diptera: Culicidae). Asia Pac J Trop Dis 4: S173-S180.
  67. Subramaniam J, Murugan K, Kovendan K (2012) Larvicidal and pupicidal efficacy of *Momordica charantia* leaf extract and bacterial insecticide, *Bacillus thuringiensis* against malarial vector, *Anopheles stephensi* Liston. (Diptera: Culicidae). J Biopest 5: 163-169.
  68. Kovendan K, Murugan K, Shanthakumar S, Vincent S, Jiang-Shiou Hwang (2012) Larvicidal activity of *Morinda citrifolia* L. (Noni) (Family: Rubiaceae) leaf extract against *Anopheles stephensi*, *Culex quinquefasciatus*, and *Aedes aegypti*. Parasitol Res 111: 1481-1490.
  69. Prabhu K, Murugan K, Nareshkumar A, Ramasubramanian N, Bragadeeswaran S (2011) Larvicidal and repellent potential of *Moringa oleifera* against malarial vector, *Anopheles stephensi* Liston (Insecta: Diptera: Culicidae). Asia Pac J Trop Biomed 124-129
  70. Krishnamoorthy S, Chandrasekaran M, Adaikala Raj G, Jayaraman M, Venkatesalu V (2015) Identification of chemical constituents and larvicidal activity of essential oil from *Murraya exotica* L. (Rutaceae) against *Aedes aegypti*,

- Anopheles stephensi* and *Culex quinquefasciatus* (Diptera: Culicidae). Parasitol Res 114: 1839-1845.
71. Santhoshkumar T, Abdul Rahuman A, Rajakumar G, Marimuthu S, Bagavan A, Jayaseelan C, et al (2011) Synthesis of silver nanoparticles using *Nelumbo nucifera* leaf extract and its larvicidal activity against malaria and filariasis vectors. Parasitol Res 108: 693-702
  72. Adewole A, Oderinde A, Bankole O, Faparusi F, Oyede RT (2013) Larvicidal activities of three plant extracts of common wire weed (*Sida Acuta*), Catnip (*Nepeta Cataria*) and Neem (*Azadirachta Indica*) against the larva of mosquito (*Anopheles Gambiae*). Acedem J Med Plant 1: 37-40.
  73. Mathath Roni, Kadarkarai Murugan, Chellasamy Panneerselvam, Jayapal Subramaniam, Jiang-Shiou Hwang (2013) Evaluation of leaf aqueous extract and synthesized silver nanoparticles using *Nerium oleander* against *Anopheles stephensi* (Diptera: Culicidae). Parasitol Res 112: 981-990.
  74. Adaikala Raj G, Chandrasekaran M, Krishnamoorthy S, Jayaraman M, Venkatesalu V (2015) Phytochemical profile and larvicidal properties of seed essential oil from *Nigella sativa* L. (Ranunculaceae), against *Aedes aegypti*, *Anopheles stephensi*, and *Culex quinquefasciatus* (Diptera: Culicidae). Parasitol Res 114: 3385-3391.
  75. Imam TS, Tajuddeen UM (2013) Qualitative phytochemical screening and larvicidal potencies of ethanolic extracts of five selected macrophyte species against *Anopheles* mosquitoes (Diptera: Culicidae). J Res Environ Scie Toxicol 2: 121-125.
  76. Maurya P, Sharma P, Mohan L, Mohan Verma M, Srivastava C (2012) Larvicidal efficacy of *Ocimum basilicum* extracts and its synergistic effect with neonicotinoid in the management of *Anopheles stephensi*. Asia Pac J Trop Dis 110-116.
  77. Pearl Ugochi Ofoegbu, Nkechi Esther Onyedineke, Chukwunonso Nwokeji, Ngozi Georgewill Esie, Nathaniel Isibor (2013) Laboratory evaluation of ethanolic and methanolic extracts of *Ocimum gratissimum* against larva of *Anopheles gambiae* and non-target organisms. Mum Ent Zool 8: 185-190.
  78. Gokulakrishnan J, Baranitharan M, Dhanasekaran S, Kavikuyil R, Abirami R, Deepa J, et al (2015) Mosquito Larvicidal Properties of *Ocimum sanctum* Linn. (Lamiaceae) against *Aedes aegypti* (Linn.), *Anopheles stephensi* (Liston), *Culex quinquefasciatus* (Say). Life Scie Arch 1: 46-52.
  79. Kovendan K, Murugan K, Vincent S, Donald R Barnard (2012) Mosquito larvicidal properties of *Orthosiphon thymiflorus* (Roth) Sleesen. (Family: Labiatae) against mosquito vectors, *Anopheles stephensi*, *Culex quinquefasciatus* and *Aedes aegypti* (Diptera: Culicidae). Asia Pac J Trop Med 299-305.
  80. Balu Selvakumar, Gokulakrishnan J, Elumalai K, Dhanasekaran S, Anandan A, Krishnappa K (2012) Mosquito larvicidal activity of *Oxystelma esculentum* plant extracts against *Anopheles stephensi* (Diptera: Culicidae). Int J Rec Sci Res 3: 321-324.
  81. Balu Selvakumar, Gokulakrishnan J, Elanchezhian K, Deepa J (2015) Mosquitocidal activities of Indian medicinal plant *Pavonia odorata* willd (Malvaceae) against selected vector mosquitoes (Diptera: Culicidae). Int J Curr Advan Res 4: 221-227.
  82. Gokulakrishnan J, Baranitharan M, Dhanasekaran S, Deepa J, Selvakumar B, Thushimenan S (2016) Laboratory evaluation of *Pedaliium murex* L. extracts on the South East India disease vector mosquitoes (Diptera: Culicidae). Int J Zool Appl Biosci 1: 7-14.
  83. Samidurai K (2012) Mosquito larvicidal and ovicidal properties of *Pemphis acidula* Frost. (Lythraceae) against *Culex tritaeniorhynchus* Giles and *Anopheles subpictus* Grassi (Diptera: Culicidae). Asia Pac J Trop Biomed 1862-1866.
  84. Chandrashekhar D Paril, Hemant P Borase, Satish V Patil, Rahul B Salunkhe,



- Bipinchandra K Salunke (2012) Larvicidal activity of silver nanoparticles synthesized using *Pergularia daemia* plant latex against *Aedes aegypti* and *Anopheles stephensi* and nontarget fish *Poecilia reticulata*. *Parasitol Res* 111: 555-562.
85. Kamaraj C, Abdul Rahuman A, Baganvan A, Elango G, Abdul Zahir A, Santhoshkumar T (2011) Larvicidal and repellent activity of medicinal plant extracts from Eastern Ghats of South India against malaria and filariasis vectors. *Asia Pac J Trop Med* 6:698-705.
86. V Tamizhazhagan, K Pugazhendy, M Baranitharan, Humphrey Agevi (2017). Pest potential of *Pisonia alba* extracts and fractions against mosquito-borne disease (Diptera: Culicidae). *International Journal of Pharma Sciences and Research*.8, 5; 52-58
87. Govindarajan M, Rajeswary M, Sivakumar R (2013) Larvicidal and ovicidal efficacy of *Pithecellobium dulce* (Roxb.) Benth. (Fabaceae) against *Anopheles stephensi* Liston and *Aedes aegypti* Linn. (Diptera: Culicidae). *Indian J Med Res* 138: 129-134.
88. Rawani A, Banerjee A and Chandra G (2012) Mosquito larvicidal and biting deterrence activity of bud of *Polianthus tuberosa* plants extract against *Anopheles stephensi* and *Culex quinquefasciatus*. *Asia Pac J Trop Dis* 200-204.
89. Deepa M, Palanisamy K, Krishnappa K, Elumalai K (2014) Mosquitocidal activity of *Polygala arvensis* Willd against *Aedes aegypti* (Linn.), *Anopheles stephensi* (Liston.) and *Culex quinquefasciatus* (Say.) (Diptera: Culicidae). *Int J Mosquito Res* 1: 30-34.
90. Lakshmanan, S., S. Thushimenan, and V. Tamizhazhagan (2017). "Antifeedant, larvicidal and oviposition deterrent activity of *Pongamia pinnata* and *Ceiba pentandra* against pod borer larvae of *Helicoverpa armigera* (Noctuidae: Lepidoptera)." *Indo American Journal of Pharmaceutical Sciences* 4.2 180-185
91. Guna Ranjan Kolli, Balakrishnan, Vijayan, Raja Sundararajan (2013) Evaluation of larvicidal activity of *Pongamia pinnata* extracts against three mosquito vectors. *Asia Pac J Trop Biomed* 3: 853-858.
92. Tyagi Varun, Yadav Ruchi, Sharma Ajay Kumar, Tyagi Vivek, Yadav Shweta, Vijay Veer, et al (2013) Larvicidal activity of leaf extract of some weeds against malaria vector *Anopheles stephensi*. *Int J Malari Res Rev* 1: 35-39.
93. Woquan Sama, Edith O Ajaiyeoba, Mohammed I Choudhary (2014) Larvicidal properties of Simalikalactone from *Quassia africana* (Simaroubaceae) Baill and Baill, on the malaria vector, *Anopheles gambiae*. *African J Tradit Complement Altern Med* 11: 84-88.
94. Awad Khalafalla Taha, Huda Elfatih Osman, Omar Ahmed A Sidahmed (2011) Larvicidal effects of some plant extracts against *Anopheles arabiensis* Patton larvae (Diptera : Culicidae). *J Scien Technol* 12: 67-73.
95. Baranitharan M, Dhanasekaran S, Gokulakrishnan J, Krishnappa K, Deepa J (2015) Mosquito larvicidal properties of *Sesamum indicum* L. against *Aedes aegypti* (Linn.), *Anopheles stephensi* (Liston), *Culex quinquefasciatus* (Say) (Diptera: Culicidae). *Life Sci Arch* 1: 72-77.
96. Govindarajan M (2010) Larvicidal and repellent activities of *Sida acuta* Burm. F. (Family: Malvaceae) against three important vector mosquitoes. *Asia Pac J Trop Med* 3: 691-695.
97. Premalatha S, Elumalai K, Jeyasankar A (2013) Mosquitocidal properties of *Solanum trilobatum* L. (Solanaceae) leaf extracts against three important human vector mosquitoes (Diptera: Culicidae). *Asia Pac J Trop Med* 854-858.
98. Elijah Ajaegbu Eze, Simon Pierre Yinyang Danga, Festus Basden Chiedu Okoye (2014) Larvicidal activity of the leaf extracts of *Spondias mombin* Linn. (Anacardiaceae) from various solvents against malaria, dengue and filarial vector mosquitoes (Diptera: Culicidae). *J Vector Borne Dis* 51: 300-306.
99. Jayaseelan C, Abdul Rahuman A, Rajakumar G, Vishnu Kirthi A, Santhoshkumar T, Marimuthu S, et al

- (2011) Synthesis of pediculocidal and larvicidal silver nanoparticles by leaf extract from heartleaf moonseed plant, *Tinospora cordifolia* Miers. *Parasitol Res* 109: 185-194.
100. Devan Elumalai, Patheri Kunyil Kaleena1, Mujeera Fathima, Naresh Kumar (2013) Phytochemical screening and larvicidal activity of *Tridax procumbens* (L) against *Anopheles stephensi* (Liston), *Aedes aegypti* (L) and *Culex quinquefasciatus* (Say). *Int J Bioscience Res* 2: 1-14.
101. Alina Hellert, Gaurav Sharma, Kaushal Kumar, Veena Agrawal (2012) Exploration of larvicidal activity of *Vernonia anthelmintica* (L.) wild seed crude extracts in different solvents against malaria (*Anopheles stephensi*) and dengue (*Aedes aegypti*) vectors. *Malaria J* 11: 1-3.
102. Ombito O Japheth, Matasyoh C Josphat, Vulule M John (2014) Chemical composition and larvicidal activity of *Zanthoxylum gillettii* essential oil against *Anopheles gambiae*. *African J Biotechnol* 13: 2175-2180.

**Table 1.** Incidence of Malaria surveillance data in India

Years	Total malaria cases million	<i>P. falciparum</i> million	Death due to malaria (Thousand)
1961	49,151	-	-
1962	59375	-	-
1963	87306	-	-
1964	112942	-	-
1965	99,667	-	-
1966	148012	-	-
1967	278214	-	-
1968	274634	-	-
1969	347975	-	-
1970	694017	-	-
1971	1322398	-	-
1972	1428649	-	-
1973	1930273	-	-
1974	3167658	-	-
1975	5166142	-	-
1976	6.47	0.75	59
1984	2.18	0.65	247
1985	1.86	0.54	213
1986	1.79	0.64	323
1987	1.66	0.62	188
1988	1.85	0.68	209
1989	2.05	0.76	268
1990	2.02	0.75	353
1991	2.12	0.92	421
1992	2.13	0.88	422
1993	2.21	0.85	354
1994	2.51	0.99	1122
1995	2.93	1.14	1151
1996	3.04	1.18	1010
1997	2.66	1.01	879
1998	2.22	1.03	664
1999	2.28	1.14	1048
2000	2.03	1.05	932
2001	2.09	1.01	1005
2002	1.84	0.90	973
2003	1.87	0.86	1006
2004	1.92	0.89	949
2005	1.82	0.81	963
2006	1.79	0.84	1707
2007	1.51	0.74	1311
2008	1.53	0.77	1055
2009	1.56	0.84	1144
2010	1.60	0.83	1018
2011	1.31	0.67	754
2012	1.07	0.53	519
2013	0.88	0.46	440
2014	1.10	0.72	561
2015	1.17	0.78	384
2016	1.09	0.71	331
2017	1.31	0.53	194
2018 (up to July)	172643	92436	14

Sources data (<http://nvbdcp.gov.in/malaria-new.html>)

**Table 2.** Plant reported for larvicidal, ovicidal, oviposition deterrent, pupicidal and repellent activity

Plant species	Family	Plant parts	Instars larvae / eggs / pupae / adults	Bioactivity	LC <sub>50</sub> and LC <sub>90</sub> values	References
<i>Abrus precatorius</i>	Fabaceae	Seed	IV instars	Larvicidal	Ethyl acetate extract was 19.31 and 71.71 ppm, respectively	Bagavan and Rahman [16]
<i>Acalypha alnifolia</i>	Euphorbiaceae	Leaf	NL	Ovicidal, repellent, adulticidal	Adulticidal activity of methanol extract was 274.76 and 495.88 ppm, respectively	Kovendan et al. [17]
<i>Achras sapota</i>	Sapotaceae	Leaf	III instars	Larvicidal, ovicidal, pupicidal and repellent	Methanol extract was 39.54 and 98.53 ppm, respectively	Krishnappa and Elumalai [18]
<i>Adansonia digitata</i>	Bornbaceae	Leaf	III instars	Larvicidal and repellent	Methanol extract was 78.18 and 155.42 ppm, respectively	Krishnappa et al. [19]
<i>Adhatoda vasica</i>	Acanthaceae	Leaf	IV instars	Larvicidal	Acetone extract was 18.20 and 96.33 ppm, respectively	Kamaraj et al. [20]
<i>Aloe vera</i>	Asphodelaceae	leaves	I, II, III, IV instars	Larvicidal, pupicidal	Ethanol extract of LC <sub>50</sub> value was 48.79, 59.09, 70.88, 83.58 ppm, respectively	Dinesh et al. [21]
<i>Annona muricata</i>	Annonaceae	Leaf	NL	Larvicidal	Aqueous extract of LC <sub>50</sub> value was 61.38 and 156.55 ppm, respectively	Santhosh et al. [22]
<i>Annona reticulata</i>	Annonaceae	leaf	III instars	Larvicidal, ovicidal and pupicidal	Methanol extract of LC <sub>50</sub> value was 74.36 and 93.85 ppm, respectively	Baluselvakumar et al. [23]
<i>Annona senegalensis</i>	Annonaceae	Leaf	IV instars	Larvicidal, ovicidal and pupicidal	N-hexane fraction of LC <sub>50</sub> value was 298.8 and 572.9 ppm, respectively	Younoussa et al. [24]
<i>Annona squamosa</i>	Annonaceae	Leaf	IV instars	Larvicidal, oviposition deterrent and repellent	Petroleum ether extract was 118.4 and 213.10 ppm, respectively	Vijaya kumar et al. [25]
<i>Arachis hypogaea</i>	Fabaceae	Leaves	IV instar	Larvicidal	Aqueous extract of LC <sub>50</sub> value was 71.57 and 214.62 ppm, respectively	KuppanVelu et al. [26]
<i>Artemisia nilagirica</i>	Compositae	Leaves	I, II, III, IV instars	Larvicidal, pupicidal, adulticidal, repellent	Methanol extract was 272.50, 311.40, 361.51, 442.51 and 590.07, 688.81, 789.34, 901.59 ppm, respectively	Panneerselvam et al. [27]
<i>Azadirachta indica</i>	Meliaceae	Leaf, pod	IV instars	Larvicidal, repellent and smoke toxicity effect	Azadirachtin was 0.299 and 1.061 ppm, respectively	Murugan et al. [28]
<i>Balanites aegyptiaca</i>	Balanitaceae	Roots	IV instars	Larvicidal	Ethanol extract of LC <sub>50</sub> value was 6.61 ppm, respectively	Nganjiwa et al. [29]
<i>Calotropis gigantean</i>	Asclepiadaceae	Leaf	I, II, III, IV instars	Larvicidal, pupicidal	Methanol extract was 73.77, 89.64, 121.69,	Kovendan et al. [30]

					155.49 and 199.31, 245.07, 373.85, 415.31 ppm, respectively	
<i>Catharanthus roseus</i>	Apocynaceae	Leaves	IV instar	Larvicidal, pupicidal	Aqueous extract was 68.62 and 184.85 ppm, respectively	Subarani et al. [31]
<i>Chlomolaena adorata</i>	Asteraceae	Leaf	III, IV instars	Larvicidal	Methanolic extract was 1613 and 8306 ppm, respectively	Jagruti et al. [32]
<i>Chomelia asiatica</i>	Rubiaceae	Leaf	III instars	Larvicidal	Aqueous extract of LC <sub>50</sub> value was 90.17 and 165.18 ppm, respectively	Muthu kumaran and Govindarajan [33]
<i>Chrysanthemum indicum</i>	Asteraceae	flower	I, II, III, IV instars	Larvicidal	Aqueous extract was 78.22, 110.63, 181.85, 298.79 and 417.30, 531.31, 675.76, 973.54 ppm, respectively	Arokiyaraj et al. [34]
<i>Cissus quadrangularis</i>	Vitaceae	Leaf	IV instars	Larvicidal and ovicidal	Methanol extract was 37.48 and 95.93 ppm, respectively	Krishnappa et al. [35]
<i>Citrullus lanatus</i>	Cucurbitaceae	Leaf	III, IV instars	Larvicidal	Methanolic extract was 84.23 and 989.39 ppm, respectively	Ajithadas aruna et al. [36]
<i>Citrus sinensis</i>	Rutaceae	Leaf	I, II, III, IV instars	Larvicidal	Ethanol extract was 182.24, 227.93, 291.69, 398.00 and 452.44, 544.72, 659.31, 858.92 ppm, respectively	Murugan et al. [37]
<i>Clausena anisata</i>	Rutaceae	Leaf oil	III instars	Larvicidal	Essential oil from 119.59 and 209.96 ppm, respectively	Govindarajan [38]
<i>Cleistanthus collinus</i>	Euphorbiaceae	Leaf	III instars	Larvicidal	Ethyl acetate extract was 399.72 and 1251.76 ppm, respectively	Arivoli and Samuel [39]
<i>Clerodendron inerme</i>	Verbinaceae	Leaf	I, II, III, IV instars	Larvicidal	Crude extract was 55.04, 63.33, 73.05, 80.16 and 125.50, 137.16, 153.55, 156.93 ppm, respectively	Kovendan and Murugan [40]
<i>Cochliobolus lunatus</i>	Pleosporaceae	Leaves	II, III, IV instars	Larvicidal	Plant extract was 1.17, 1.30, 141 and 2.99, 3.13, 3.29 ppm, respectively	Rahul et al. [41]
<i>Coleus aromaticus</i>	Lamiaceae	Leaf	III instars	Larvicidal	Ethyl acetate extract was 28.88 and 65.35 ppm, respectively	Barani tharan et al. [42]
<i>Commiphora caudata</i>	Bursaceae	Leaf	III instars	Larvicidal	Ethyl acetate extract was 96.04 and 104.44 ppm, respectively	Barani tharan and Dhanasekaran [43]
<i>Corchorus capsularis</i>	Malvaceae	Leaf	III instars	Larvicidal, ovicidal	Methanol extract was 176.19 and 334.56 ppm, respectively	Elangovan et al. [44]
<i>Cymbopogon citratus</i>	Poaceae	Leaf	III instars	Larvicidal	Methanol extract was 74.02 and 158.20 ppm, respectively	Karunamoorthi and Ilango [45]
<i>Cuminum cyminum</i>	Apiaceae	Vegetables	III instars	Larvicidal (24, 48 and 72	Chloroform; methanol (1:1) extract of 0.15,	Singha, and Chandra [46]

				hours)	0.09, 0.04 and 1.21, 0.23, 0.12 ppm, respectively	
<i>Datura metel</i>	Solanaceae	Leaves	I, II, III, IV instars	Larvicidal	Aqueous extract was 34.69, 42.16, 47.16, 55.22 ppm and 105.47, 118.82, 133.25, 150.11 ppm, respectively	Murugan et al. [47]
<i>Datura stramonium</i>	Solanaceae	Leaf	IV instars	Larvicidal, repellent	Ethanol extract of LD <sub>50</sub> and LD <sub>90</sub> values was 16.07 and 41.95 dose, respectively	Swathi et al. [48]
<i>Derris elliptica</i>	Fabaceae	Root, shoot	III instars	Larvicidal	Petroleum ether extract was 0.307 and 0.58 ppm, respectively	Dohutia et al. [49]
<i>Elaeagnus indica</i>	Elaeagnaceae	Leaf	IV instar	Larvicidal	Acetone extract of LC <sub>50</sub> value was 3.30 and 6.94 ppm, respectively	Srinivasan et al. [50]
<i>Eranthemum roseum</i>	Acanthaceae	Leaf	III instars	Larvicidal, oviducidal, pupicidal	Acetone extract was 121.65 and 237.38 ppm, respectively	Elumalai et al. [51]
<i>Ervatamia coronaria</i>	Apocynaceae	Leaf	III instar	Larvicidal	Methanol extract was 86.47 and 159.59 ppm, respectively	Govindarajan et al. [52]
<i>Euodia ridleyi</i>	Rutaceae	Leaf		Larvicidal	Ethyl acetate extract was 120.0 and 178.20 ppm, respectively	Prathibha et al. [53]
<i>Euphorbia hirta</i>	Euphorbiaceae	Leaves	I, II, III, IV instars	Larvicidal, pupicidal	Methanol extract was 121.51, 145.40, 169.11, 197.40 and 236.44, 293.75, 331.42, 371.34 ppm, respectively	Agalya Priyadarshini et al. [54]
<i>Exacum pedunculatum</i>	Gentianaceae	Leaf	IV instars	Larvicidal and oviducidal	Ethanol extract was 121.24 240.57 ppm, respectively	Elangovan et al. [55]
<i>Ficus benghalensis</i>	Moraceae	Leaf	III instars	Larvicidal	Methanol extract of 56.66 and 100.88 ppm, respectively	Govindarajan et al. [56]
<i>Gliricidia sepium</i>	Leguminosae	Leaf	III instars	Larvicidal, oviducidal and pupicidal	Ethanol extract was 121.79 and 231.98 ppm, respectively	Krishnappa et al. [57]
<i>Gnetum ula</i>	Gnetaceae	Leaf	III instars	Larvicidal, oviducidal and repellent	Ethanol extract of LC <sub>50</sub> value was 82.86 ppm, respectively	Dhanasekaran et al. [58]
<i>Gymnema sylvestris</i>	Asclepiadaceae	Leaves	IV instar	Larvicidal	Petroleum ether extract of LC <sub>50</sub> value was 166.28 ppm, respectively	Gopiesh Khanna et al. [59]
<i>Hippophae rhamnoides</i>	Elaeagnaceae	Leaf	IV instars	Larvicidal (24 hours only)	Ethanol extract was 1494.30 and 2605.78 ppm, respectively	Bazhir and Javid [60]
<i>Hybanthus enneaspermus</i>	Violaceae	Leaf	IV instar	Larvicidal	Aqueous extract of LC <sub>50</sub> value was 117.83, respectively	Suman et al. [61]
<i>Jatropha curcas</i>	Euphorbiaceae	Leaf	III instars	Larvicidal	Methanol extract was 92.09 and 241.09 ppm, respectively	Zewdneh Tomass et al. [62]
<i>Juniperus</i>	Cupressaceae	Oil	III instars	Larvicidal	Essential oil was 14.42	Kaliyaperumal et

<i>procera</i>					and 24.65 ppm, respectively	al. [63]
<i>Lantana camara</i>	Verbenaceae	Flowers	III, IV instars	Larvicidal	Petroleum ether extract was 126.7 and 248.9 ppm, respectively	Babita et al. [64]
<i>Leucas aspera</i>	Lamiaceae	Leaf	III instars	Larvicidal	Ethyl acetate extract was 352.84 and 1033.60 ppm, respectively	Arivoli et al. [65]
<i>Melissa officinalis</i>	Lamiaceae	Oil	III instar	Larvicidal, ovicidal and repellent	Citronellal compound was 85.44 and 159.73 ppm	Barani tharan et al. [14]
<i>Metarhizium anisopliae</i>	Clavicipitaceae	Leaves	I, II, III, IV instars	Larvicidal, pupicidal	Ethanol extract was 1.40, 3.99, 5.56, 8.77 and 13.84, 17.62, 22.20, 25.71 ppm, respectively	Kovendan et al. [66]
<i>Momordica charantia</i>	Cucurbitaceae	Leaf	I, II, III, IV instars	Larvicidal, pupicidal	Methanol extract was 93.45, 123.74, 167.17, 216.15 and 454.96, 573.31, 630.66, 722.25 ppm, respectively	Subramaniam et al. [67]
<i>Morinda citrifolia</i>	Rubiaceae	Leaf	III instar	Larvicidal	Methanol extract was 261.96 and 505.06 ppm, respectively	Kovendan et al. [68]
<i>Moringa oleifera</i>	Moringaceae	Leaf	I, II, III and IV instars	Larvicidal, pupicidal and repellent	Methanol extract was 57.7, 63.9, 72.4, 78.9 and 125.9, 133.0, 139.8, 143.2 ppm, respectively	Prabhu et al. [69]
<i>Murraya exotica</i>	Rutaceae	Oil	IV instars	Larvicidal	Essential oil of LC <sub>50</sub> value was 56.3 and 107.8 ppm, respectively	Krishnamoorthy et al. [70]
<i>Nelumbo nucifera</i>	Nelumbonaceae	Leaves	IV instar	Larvicidal	Methanol extract was 8.89 and 28.65 ppm, respectively	Santhoshkumar et al. [71]
<i>Nepeta cataria</i>	Lamiaceae	Leaf	IV instars	Larvicidal	Methanol extract of LC <sub>50</sub> value was 0.93 ppm, respectively	Adewole et al. [72]
<i>Nerium oleander</i>	Apocynaceae	Leaf	I, II, III, IV instars	Larvicidal, pupicidal	Extract was 20.60, 24.90, 28.22, 33.55 and 41.62, 50.33, 57.78, 33.99 ppm, respectively	Mathath Roni et al. [73]
<i>Nigella sativa</i>	Ranunculaceae	Seed	IV instars	Larvicidal	Essential oil of LC <sub>50</sub> value was 88.1 and 272.4 ppm, respectively	Adaikala Raj et al. [74]
<i>Nymphaea lotus</i>	Nymphaeaceae	Leaf	III instars	Larvicidal	Ethanol extract of LC <sub>50</sub> value was 62.8 ppm, respectively	Iman and Tajuddeen [75]
<i>Ocimum basilicum</i>	Lamiaceae	Leaf	III instars	Larvicidal	Methanol extract of 0.01 and 3.13 ppm, respectively	Maurya et al. [76]
<i>Ocimum gratissimum</i>	Lamiaceae	Leaf	Instars larvae	Larvicidal	Ethanol extract was 60.9 and 464.4 mg/ml, respectively	Pearl et al. [77]
<i>Ocimum sanctum</i>	Lamiaceae	Leaf	III instars	Larvicidal	Methanol extract was 115.32 and 209.25 ppm, respectively	Gokulakrishnan et al. [78]
<i>Orthosiphon</i>	Labiatae	Leaf	III instars	Larvicidal	Methanol extract of	Kovendan et al.

<i>thymiflorus</i>					118.74 and 377.09 ppm, respectively	[79]
<i>Oxystelma esculentum</i>	Apocynaceae	Leaf	III instars	Larvicidal	Methanol extract was 63.84 and 122.48 ppm, respectively	Balu et al. [80]
<i>Pavonia odorata</i>	Malvaceae	Leaf	III instars	Larvicidal and repellent	Methanol extract was 58.22 and 239.82 ppm, respectively	Balu et al. [81]
<i>Pedaliium murex</i>	Pedaliaceae	Leaf	III instar	Larvicidal, ovicidal and repellent	Methanol extract was 111.66 and 200.01 ppm	Gokulakrishnan et al. [82]
<i>Pemphis acidula</i>	Lythraceae	Leaf	III instars	Larvicidal and ovicidal	Methanol extract of 22.1 and 43.71 ppm, respectively	Samidurai [83]
<i>Pergularia daemia</i>	Asclepiadaceae	Leaves	I, II, III, IV	Larvicidal	Crude aqueous extract was 81.47 and 159.51 ppm, respectively	Chandrashekhar et al. [84]
<i>Piper nigrum</i>	Piperaceae	Seed	III instars	Larvicidal and repellent	Ethyl acetate was 24.54 and 108.03 ppm, respectively	Kamaraj et al. [85]
<i>Pisonia alba</i>	Nyctaginaceae	Leaves	III instars	Larvicidal and ovicidal	Petroleum ether 1.0 ppm 2.0	Tamizhazhagan et al [86]
<i>Pithecellobium dulce</i>	Fabaceae	Leaf, seed	III instars	Larvicidal and ovicidal	Methanol extract was 145.43 and 251.23 ppm, respectively	Govindarajan et al. [87]
<i>Polianthes tuberosa</i>	Agavaceae	Bud	I, II, III and IV instars	Larvicidal and repellent (24 hours only)	Crude plant extract of 0.18, 0.51, 0.05, 0.30 and 1.38, 0.51, 0.29, 7.03 ppm, respectively	Rawani et al. [88]
<i>Polygala arvensis</i>	Polygalaceae	Leaf	III instars	Larvicidal, ovicidal and repellent	Methanol extract was 46.37 and 189.82 ppm, respectively	Deepa et al. [89] and Lakshmanan et al.[90]
<i>Pongamia pinnata</i>	Fabaceae	Bark	IV instars	Larvicidal	Methanol extract was 151.7 and 299.4 ppm, respectively	Guna et al. [91]
<i>Prosopis juliflora</i>	Fabaceae	Leaf	III instars	Larvicidal	Leaf extracts was 37.55 and 514.35 ppm, respectively	Tyagi Varun et al. [92]
<i>Quassia africana</i>	Simaroubaceae	Leaf, stem, roots	IV instars	Larvicidal	Methanol extract of LC <sub>50</sub> value was 17.58 ppm, respectively	Woquan Sama et al. [93]
<i>Ricinus communis</i>	Euphorbiaceae	Leaf	III instars	Larvicidal	Ethanol extract was 282.70 and 501.23 ppm, respectively	Awad et al. [94]
<i>Sesamum indicum</i>	Pedaliaceae	Leaf	III instar	Larvicidal	Methanol extract was 338.27 and 538.50 ppm, respectively	Barani tharan et al. [95]
<i>Sida acuta</i>	Malvaceae	Leaf	III instars	Larvicidal and repellent	Methanol extract was 38.64 and 74.78 ppm, respectively	Govindarajan [96]
<i>Solanum trilobatum</i>	Solanaceae	Leaf		Toxicity	116.64 ppm	Premalatha et al. [97]
<i>Spondias mombin</i>	Anacardiaceae	Leaf	IV instars	Larvicidal	Hexane fraction was 92.2 and 245.37 ppm, respectively	Elijah et al. [98]



<i>Tinospora cordifolia</i>	Menispermaceae	Leaf	IV instars	Larvicidal	Aqueous extract of LC <sub>50</sub> was 53.93 ppm, respectively	Jayaseelan <i>et al.</i> [99]
<i>Tridax procumbens</i>	Compositae	Whole	IV instars	Larvicidal (24 hours only)	Methanol extract was 57.991 and 172.280 ppm, respectively	Devan Elumalai <i>et al.</i> [100]
<i>Vernonia anthelmintica</i>	Asteraceae	Seed	III, IV instars	Larvicidal	Ethanol extract was 1.95 and 10.49 ppm, respectively	Alina <i>et al.</i> [101]
<i>Zanthoxylum gillettii</i>	Rutaceae	Oil	III instars	Larvicidal	Essential oil LC <sub>50</sub> value was 57.73 and 140.24 ppm, respectively	Ombito <i>et al.</i> [102]